arXiv:math/0701091v1 [math.AG] 3 Jan 2007

UNIVERSITÀ DEGLI STUDI DI ROMA
"LA SAPIENZA"

Facoltà di Scienze Matematiche Fisiche e Naturali
Dottorato di Ricerca in Matematica
XVIII Ciclo

Differential Graded Lie Algebras and Deformations of Holomorphic Maps

Dottoranda
Donatella Iacono

Relatore
Prof. *Marco Manetti*

This thesis is devoted
to the memory
of my grandmother Mina.

Contents

Introduction 5

Chapter I. Functors of Artin rings 15
  I.1. Generalities on functors of Artin rings 15
  I.2. Deformation functor of complex manifolds 24
  I.3. DGLAs and deformation functor 28

Chapter II. Deformation of complex manifolds 41
  II.1. Differential forms 42
  II.2. Kähler manifolds 43
  II.3. Holomorphic fiber bundle and Dolbeault's cohomology 44
  II.4. The Kodaira-Spencer algebra KS_X 47
  II.5. Contraction map and holomorphic Lie derivative 47
  II.6. Induced maps f_* and f^* by a holomorphic map f 52
  II.7. Deformations of complex manifolds 55

Chapter III. Deformation functor of a pair of morphisms of DGLAs 63
  III.1. Functors $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$ 63
  III.2. Extended Deformation Functors 73

Chapter IV. Deformations of holomorphic maps 87
  IV.1. Deformation functor $\mathrm{Def}(f)$ of holomorphic maps 87
  IV.2. Infinitesimal deformations of holomorphic maps 92

Chapter V. Semiregularity maps 101
  V.1. Deformations with fixed codomain 102
  V.2. Deformations of a map with fixed domain 109
  V.3. Deformations with fixed domain and codomain 109
  V.4. Deformations of the inclusion map 110

Bibliography 115

Introduction

In the last fifty years deformation theory has played an important role in algebraic and complex geometry.

The study of small deformations of the complex structures of complex manifolds started with the works of K. Kodaira and D.C. Spencer [**17**] and M. Kuranishi [**19**].

Then A. Grothendieck [**12**], M. Schlessinger [**32**] and M. Artin [**1**] formalized this theory translating it into a functorial language. The idea was that with an infinitesimal deformation of a geometric object we can associate a deformation functor of Artin rings: that is, a functor from the category **Art** of local Artinian $\mathbb{C}$-algebras (with residue field $\mathbb{C}$) to the category **Set** of sets, that satisfies some extra conditions (see Definition I.1.10).

A modern approach to deformation theory is via differential graded Lie algebras (DGLA for short).

The guiding principle is the idea due to P. Deligne, V. Drinfeld, D. Quillen and M. Kontsevich (see [**18**]) that "in characteristic zero every deformation problem is controlled by a differential graded Lie algebra".

Inspired by this principle, the aim of this thesis is to follow the modern approach to study the infinitesimal deformations of holomorphic maps of compact complex manifolds.

More precisely, a DGLA is a differential graded vector space with a structure of graded Lie algebra, plus some compatibility conditions between the differential and the bracket (of the Lie structure) (see Definition I.3.5).

Moreover, using the solutions of the Maurer-Cartan equation $dx + \dfrac{1}{2}[x,x] = 0$ it is well known how we can associate with a DGLA L a deformation functor Def_L. Written in detail:

$$\operatorname{Def}_L : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\operatorname{Def}_L(A) = \frac{\{x \in L^1 \otimes m_A \mid dx + \frac{1}{2}[x,x] = 0\}}{gauge},$$

where m_A is the maximal ideal of A and the gauge equivalence is induced by the gauge action $*$ of $\exp(L^0 \otimes m_A)$ on the set of solutions of the Maurer-Cartan equation (see Definition I.3.30).

Then the idea of the principle is that we can define a DGLA L (up to quasi-isomorphism) from the geometrical data of the problem, such that the deformation functor Def_L is isomorphic to the deformation functor associated with the geometric problem.

We note that it is easier to study a deformation functor associated with a DGLA but, in general, it is not an easy task to find the right DGLA (up to quasi-isomorphism) associated with the problem.

A first example, in which the associated DGLA is well understood, is the case of deformations of complex manifolds.

Let X be a compact complex manifold. Then X is obtained by gluing together a finite number of polydisks in $\mathbb{C}^n$. The fundamental idea of K. Kodaira and D.C. Spencer is that “a deformation of X is regarded as the gluing of the same polydisks via different identifications” (see [**16**, pag. 182]) .

Translating it into a functorial language we define, for each $A \in$ **Art**, an infinitesimal deformation of X over $\mathrm{Spec}(A)$ as a commutative diagram

$$\begin{tikzcd}
X \arrow[r, "i"] \arrow[d] & X_A \arrow[d, "\pi"] \\
\mathrm{Spec}(\mathbb{C}) \arrow[r, "a"] & \mathrm{Spec}(A),
\end{tikzcd}$$

where π is a proper flat holomorphic map and X coincides with the restriction of X_A over the closed point of $\mathrm{Spec}(A)$ (see Definition I.2.4). Moreover, we can give the notions of isomorphism and of trivial deformation ($X_A \cong X \times \mathrm{Spec}(A)$).

Thus we can define the functor associated with the infinitesimal deformations of X:

$$\mathrm{Def}_X : \mathbf{Art} \to \mathbf{Set},$$

$$\mathrm{Def}_X(A) = \left\{ \begin{array}{c} \text{isomorphism classes of} \\ \text{infinitesimal deformations} \\ \text{of } X \text{ over } \mathrm{Spec}(A) \end{array} \right\}.$$

Therefore, following the idea of the principle, we are looking for a DGLA L such that $\mathrm{Def}_L \cong \mathrm{Def}_X$.

Let Θ_X be the holomorphic tangent bundle of X and consider the sheaf $\mathcal{A}_X^{0,*}(\Theta_X)$ of the differential forms of $(0,*)$-type, with coefficients in Θ_X.

Then we define the Kodaira-Spencer algebra $A_X^{0,*}(\Theta_X)$ of X as the graded vector space of global sections of the sheaf $\mathcal{A}_X^{0,*}(\Theta_X)$.

Considering the (opposite) Dolbeault differential and the bracket of vector fields, it is possible to endow $A_X^{0,*}(\Theta_X)$ with a natural structure of differential graded Lie algebra (see Definition II.4.1).

The DGLA $A_X^{0,*}(\Theta_X)$ controls the problem of infinitesimal deformations of X (see Theorem II.7.3):

THEOREM (A). *Let X be a complex compact manifold and $A_X^{0,*}(\Theta_X)$ its Kodaira-Spencer algebra. Then there exists an isomorphism of functors*

$$\mathrm{Def}_{A_X^{0,*}(\Theta_X)} \longrightarrow \mathrm{Def}_X .$$

In this case the correspondence is clear between the solutions of the Maurer-Cartan equation and the infinitesimal deformations of X, such that the gauge equivalence corresponds to the isomorphism of deformations. In particular, a solution of the Maurer-Cartan equation is gauge equivalent to zero if and only if it induces a trivial deformation of X.

The next natural problem is to investigate the embedded deformations of a submanifold in a fixed manifold.

Very recently, M. Manetti in [**24**] studies this problem using the approach via DGLA.

In his work, more than proving the existence of a DGLA which controls this geometric problem, M. Manetti develops some algebraic tools related to the DGLA.

More precisely, he describes a general construction to define a new deformation functor associated with a morphism of DGLAs.

Given a morphism of differential graded Lie algebras

$$h : L \longrightarrow M$$

he defines the functor

$$\mathrm{Def}_h : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{Def}_h(A) = \frac{\{(l,m) \in (L^1 \otimes m_A) \times (M^0 \otimes m_A) |\ dx + \frac{1}{2}[x,x] = 0 \text{ and } e^m * h(l) = 0\}}{gauge},$$

where this gauge equivalence is a generalization of the previous one (see Remark III.1.13). This new functor is an extension of the functor associated with a single DGLA: by choosing $h = M = 0$, Def_h reduces to Def_L.

Moreover, using path objects (see Example I.3.14) he shows that for every choice of L, M and h there exists a DGLA H such that $\mathrm{Def}_H \cong \mathrm{Def}_h$.

In particular, this implies that if a deformation functor associated with a geometric problem is isomorphic to Def_h, for some $h : L \longrightarrow M$, then automatically we know the existence of a DGLA that controls the problem and we have an explicit description of it.

This is what M. Manetti does in [**24**]: by suitably choosing L, M and h he proves that there exists an isomorphism between the functor Def_h and the functor associated with the infinitesimal deformations of a submanifold in a fixed manifold. Actually, let X be a compact complex manifold and Z a submanifold. The infinitesimal emebedded deformations of Z can be interpreted as the deformations of the inclusion map $i : Z \hookrightarrow X$ inducing the trivial deformation on X.

Consider the Kodaira-Spencer DGLA $A_X^{0,*}(\Theta_X)$ of X and the differential graded Lie subalgebra $A_X^{0,*}(\Theta_X(-log\, Z))$ defined by the following exact sequence

$$0 \longrightarrow A_X^{0,*}(\Theta_X(-log\, Z)) \longrightarrow A_X^{0,*}(\Theta_X) \longrightarrow A_Z^{0,*}(N_{Z|X}) \longrightarrow 0,$$

where $N_{Z|X}$ is the normal bundle of Z in X (see Section II.5.1).

We have already observed that the DGLA $A_X^{0,*}(\Theta_X)$ controls the infinitesimal deformations of X, whereas the DGLA $A_X^{0,*}(\Theta_X(-log\, Z))$ controls the infinitesimal deformations of the pair $Z \subset X$ (each solution of the Maurer-Cartan equation in $A_X^{0,*}(\Theta_X(-log\, Z))$ defines a deformation of both Z and X).

Fix $M = A_X^{0,*}(\Theta_X)$, $L = A_X^{0,*}(\Theta_X(-log\, Z))$ and h the inclusion:

$$h : A_X^{0,*}(\Theta_X(-log\, Z)) \hookrightarrow A_X^{0,*}(\Theta_X).$$

Then it is clear how we can associate with each element $(l, m) \in \mathrm{Def}_h$ a deformation of Z in X, with X fixed: the infinitesimal deformation of Z is the one corresponding to the Maurer-Cartan solution $l \in A_X^{0,*}(\Theta_X(-log\, Z))$ and it induces a trivial deformation of X, since we are requiring that $h(l)$ is gauge equivalent to zero in $A_X^{0,*}(\Theta_X)$.

These new ideas developed by M. Manetti are of fundamental importance for this thesis which, in some sense, can be considered a generalization of them. Actually, we extend these techniques to study not only the deformations of an inclusion but, in general, the deformations of holomorphic maps.

These deformations were first studied from the classical point of view (no DGLA) by E. Horikawa [**14**] and [**15**], M. Namba [**27**] and by Z. Ran [**28**].

More precisely, let $f : X \longrightarrow Y$ be a holomorphic map of compact complex manifolds.

There are several aspects of deformations of f: we can deform just the map f fixing both X and Y, we can allow to deform f and X or Y or, more in general, we can deform everything: the map f, X and Y.

The infinitesimal deformations of f, with fixed domain and target, can be interpreted as infinitesimal deformations of the graph of f in the product $X \times Y$, with $X \times Y$ fixed (see Section V.3). Therefore, we are considering infinitesimal deformations of a submanifold in a fixed manifold and so the DGLA approach to this case is implicitly included in M. Manetti's work [**24**].

Next, we turn our attention to the general case in which we deform f, X and Y.

In a functorial language, for each $A \in \mathbf{Art}$, we define an infinitesimal deformation of f over $\operatorname{Spec}(A)$ as a commutative diagram

$$\begin{tikzcd}
X_A \arrow[rr, "F"] \arrow[dr, "\pi"'] & & Y_A \arrow[dl, "\mu"] \\
& S, &
\end{tikzcd}$$

where $S = \operatorname{Spec}(A)$, (X_A, π, S) and (Y_A, π, S) are infinitesimal deformations of X and Y respectively, and F is a holomorphic map that restricted to the fibers over the closed point of S coincides with f.

In this case too we can give the notions of isomorphism and of trivial deformation.

Then we can define the functor of infinitesimal deformations of a holomorphic map $f : X \longrightarrow Y$:

$$\operatorname{Def}(f) : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\operatorname{Def}(f)(A) = \left\{ \begin{array}{c} \text{isomorphism classes of} \\ \text{infinitesimal deformations of} \\ f \text{ over } \operatorname{Spec}(A) \end{array} \right\}.$$

Let Γ be the graph of f in $X \times Y$. An infinitesimal deformation of the map f can be interpreted as an infinitesimal deformation of Γ in $X \times Y$, such that the induced deformation of the product $X \times Y$ is a product of deformations of X and Y. In general not all the deformations of a product are products of deformations, as it was showed by K. Kodaira and D.C Spencer (see Remark II.7.5).

Consider the Kodaira-Spencer algebra $A^{0,*}_{X\times Y}(\Theta_{X\times Y})$ of the product $X \times Y$ and the differential graded Lie subalgebra $A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma))$ defined by the following exact sequence

$$0 \longrightarrow A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma)) \longrightarrow A^{0,*}_{X\times Y}(\Theta_{X\times Y}) \longrightarrow A^{0,*}_{\Gamma}(N_{\Gamma|X\times Y}) \longrightarrow 0,$$

where $N_{\Gamma|X\times Y}$ is the normal bundle of the graph Γ in $X \times Y$ (see Section II.5.1).

As before, we know that $A^{0,*}_{X\times Y}(\Theta_{X\times Y})$ controls the infinitesimal deformations of $X \times Y$ and $A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma))$ controls the infinitesimal deformations of the pair $\Gamma \subset X \times Y$ (each solution of the Maurer-Cartan equation in $A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma))$ defines a deformation of both Γ and $X \times Y$).

Fix $M = A^{0,*}_{X\times Y}(\Theta_{X\times Y})$, $L = A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma))$ and h the inclusion:

$$h : A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma)) \hookrightarrow A^{0,*}_{X\times Y}(\Theta_{X\times Y}).$$

In the general case, it does not suffice to consider just the DGLA L or the morphism $h : L \longrightarrow M$, since they have no control on the induced deformations on $X \times Y$.

Therefore, we need to define a new functor: the deformation functor associated with a pair of morphisms.

Given morphisms of differential graded Lie algebras $h : L \longrightarrow M$ and $g : N \longrightarrow M$:

$$\begin{tikzcd} & L \arrow[d, "h"] \\ N \arrow[r, "g"] & M, \end{tikzcd}$$

we define the functor

$$\mathrm{Def}_{(h,g)} : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{Def}_{(h,g)}(A) = \{(x, y, e^p) \in (L^1 \otimes m_A) \times (N^1 \otimes m_A) \times \exp(M^0 \otimes m_A)|$$

$$dx + \frac{1}{2}[x, x] = 0,\ dy + \frac{1}{2}[y, y] = 0,\ g(y) = e^p * h(x)\}/gauge,$$

where this gauge equivalence is an extension of the previous ones (see Definition III.1.12).

This functor is a generalization of the previous ones: by choosing $N = 0$ and $g = 0$, $\mathrm{Def}_{(h,g)}$ reduces to Def_h; by choosing $N = M = 0$ and $h = g = 0$, $\mathrm{Def}_{(h,g)}$ reduces to Def_L.

Consider the DGLA $A^{0,*}_X(\Theta_X) \times A^{0,*}_Y(\Theta_Y)$ and the morphism $g = (p^*, q^*) : A^{0,*}_X(\Theta_X) \times A^{0,*}_Y(\Theta_Y) \longrightarrow A^{0,*}_{X\times Y}(\Theta_{X\times Y})$, where p and q are the natural projections of the product $X \times Y$ on X and Y, respectively: $p : X \times Y \longrightarrow X$ and $q : X \times Y \longrightarrow Y$.

We note that the solutions $n = (n_1, n_2)$ of the Maurer-Cartan equation in $N = A^{0,*}_X(\Theta_X) \times A^{0,*}_Y(\Theta_Y)$ correspond to infinitesimal deformations of both X (induced by n_1) and Y (induced by n_2). Moreover the image $g(n)$ satisfies the Maurer-Cartan equation in $M = A^{0,*}_{X\times Y}(\Theta_{X\times Y})$ and so it is associated with an infinitesimal deformation of $X \times Y$, that is exactly the one obtained as product of the deformations of X (induced by n_1) and of Y (induced by n_2).

Therefore, this g gives exactly the control on the deformations of the product that we are looking for.

Let $M = A^{0,*}_{X\times Y}(\Theta_{X\times Y})$, $L = A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma))$, $h : L \longrightarrow M$ the inclusion, $N = A^{0,*}_X(\Theta_X) \times A^{0,*}_Y(\Theta_Y)$ and $g = (p^*, q^*) : N \longrightarrow M$. Then we are in the following situation:

$$\begin{tikzcd}
 & A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma)) \arrow[d, hook, "h"] \\
A^{0,*}_X(\Theta_X)\times A^{0,*}_Y(\Theta_Y) \arrow[r, "g=(p^*,q^*)"] & A^{0,*}_{X\times Y}(\Theta_{X\times Y}).
\end{tikzcd}$$

In conclusion, each deformation of the map f corresponds to a Maurer-Cartan solution $l \in A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma))$, such that $h(l)$ induces a deformation of $X \times Y$ isomorphic to a deformation induced by $g(n)$, for some Maurer-Cartan solution $n \in A^{0,*}_X(\Theta_X) \times A^{0,*}_Y(\Theta_Y)$ (that is, $h(l)$ and $g(n)$ are gauge equivalent in $A^{0,*}_{X\times Y}(\Theta_{X\times Y})$).

Therefore, $\mathrm{Def}_{(h,g)}$ encodes all the geometric data of the problem and the following theorem is quite obvious (see Theorem IV.2.5).

Theorem (B). *Let $f : X \longrightarrow Y$ be a holomorphic map of compact complex manifold. Then, with the notation above, there exists an isomorphism of functors*

$$\mathrm{Def}_{(h,g)} \cong \mathrm{Def}(f).$$

This theorem holds for the general case of infinitesimal deformations of f, the other cases are obtained as specializations of it.

For example, the deformations of f with fixed domain or fixed target, are obtained by considering $N = A^{0,*}_Y(\Theta_Y)$ or $N = A^{0,*}_X(\Theta_X)$, respectively.

In particular, using path objects, for each choice of $h : L \longrightarrow M$ and $g : N \longrightarrow M$, we are able to find a differential graded Lie algebra $H_{(h,g)}$ such that $\mathrm{Def}_{\mathrm{H}_{(h,g)}} \cong \mathrm{Def}_{(h,g)}$ (see Theorem III.2.36).

Therefore, we give an explicit description (more than the existence) of a DGLA that controls the deformations of holomorphic maps (Theorem IV.2.6).

Finally, we apply these techniques to study the obstructions to deform holomorphic maps.

The idea is the following: if we have an infinitesimal deformation of a geometric object, then we want to know if it is possible to extend it.

More precisely, let $F : \mathbf{Art} \longrightarrow \mathbf{Set}$ be a deformation functor. A (complete) obstruction space for F is a vector space V, such that for each surjection $B \longrightarrow A$ in **Art** and each element $x \in F(A)$, there exists an obstruction element $v_x \in V$, associated with x, that is zero if and only if x can be lifted to $F(B)$ (for details see Section I.1.1).

Therefore, we would like to control this obstruction space and know when the associated obstruction element is zero.

In general, we just know a vector space that contains these elements but we have no explicit description of which elements actually are obstructions. Among other things, if W is another vector space which contains V, then also W is an obstruction space for F. Then, in some sense we are looking for the "smallest" obstruction space (see Remark I.1.26).

For example, the obstructions of the functor associated with a DGLA L are naturally contained in $H^2(L)$ (see Section I.3.5), but we do not know which classes in $H^2(L)$ are indeed obstructions.

In the case of a complex compact manifold X, an obstruction space for the deformation functor Def_X is the second cohomology vector space $H^2(X, \Theta_X)$ of the holomorphic tangent bundle Θ_X of X (Theorem I.2.9).

If X is also Kähler, then A. Beauville, H. Clemens [**5**] and Z. Ran [**29**] [**30**] proved that the obstructions are contained in a subspace of $H^2(X, \Theta_X)$ defined as the kernel of a well defined map. This is the so-called "Kodaira's principle" (see for example [**5**, Theorem 10.1], [**22**, Corollary 3.4] or [**7**, Corollary 12.6], [**29**, Theorem 0] or [**30**, Corollary 3.5]).

In the case of embedded deformations of a submanifold Z in a fixed manifold X, then the obstructions are naturally contained in the first cohomology $H^1(Z, N_{Z|X})$ of the normal bundle $N_{Z|X}$ of Z in X. In this case too, if X is Kähler, it is possible to define a map on $H^1(Z, N_{Z|X})$, called the "semiregularity map", that contains the obstructions in the kernel. The idea of this map is due to S. Bloch [**3**] and it is also studied, using the DGLA approach, by M. Manetti [**24**, Theorem 0.1 and Section 9].

In the case of deformations of a holomorphic map $f : X \longrightarrow Y$ with fixed codomain, it was proved by E. Horikawa in [**14**] (see Theorem IV.1.10) that the obstructions are contained in the second hypercohomology group $\mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right)$.

Using the approach via DGLA that we have explained, we can give an easy proof of this theorem (Proposition V.1.1) but, maybe most important, we can improve it in the case of Kähler manifolds (Corollary V.1.5).

Actually, let $n = dimX$, $p = dimY - dimX$ and $\mathcal{H}$ be the space of harmonic forms on Y of type $(n+1, n-1)$. By Dolbeault's theorem and Serre's duality we obtain the equalities $\mathcal{H}^\nu = (H^{n-1}(Y, \Omega_Y^{n+1}))^\nu = H^{p+1}(Y, \Omega_Y^{p-1})$.

Using the contraction $\lrcorner$ of vector fields with differential forms (see Sections II.5 and V.1.2), for each $\omega \in \mathcal{H}$ we can define the following map

$$A_X^{0,*}(f^*\Theta_Y) \xrightarrow{\lrcorner\omega} A_X^{n,*+n-1}$$

$$\lrcorner\omega(\phi f^*\chi) = \phi f^*(\chi \lrcorner \omega) \in A_X^{n,p+n-1} \qquad \forall\ \phi f^*\chi \in A_X^{0,p}(f^*\Theta_Y).$$

By choosing ω such that $f^*\omega = 0$, we get the following commutative diagram (see Corollary V.1.5)

$$\begin{tikzcd}
A_X^{0,*}(f^*\Theta_Y) \arrow[r, "\lrcorner\omega"] & A_X^{n,*+n-1} \\
A_X^{0,*}(\Theta_X) \arrow[u, "f_*"] \arrow[r] & 0. \arrow[u]
\end{tikzcd}$$

Then, for each ω we get a morphism

$$\mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right) \longrightarrow H^n(X, \Omega_X^n),$$

which composed with the integration on X gives

$$\sigma : \mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

Using σ we get the following theorem (see Corollary V.1.5).

Theorem (C). *Let $f : X \longrightarrow Y$ be a holomorphic map of compact Kähler manifolds. Let $p = dimY - dimX$. Then the obstruction space to the infinitesimal deformations of f with fixed Y is contained in the kernel of the map*

$$\sigma : \mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

The structure of this work is as follows.

Chapter I contains the basic material about deformation functors. In Section I.1 we define the deformation functors of Artin rings, tangent and obstruction spaces and some related properties.

Section I.2 is devoted to studying the deformation functor Def_X of the infinitesimal deformations of a compact complex manifold X.

In the last Section I.3 we introduce the differential graded Lie algebras (DGLAs) and two associated functors: the Maurer-Cartan functor MC_L and the deformation functor Def_L (for each DGLA L).

In *Chapter II* we fix the notation about complex manifolds. We recall the notions of differential forms (Section II.1) of Čech and Dolbeault cohomology (Section II.3) and some properties of Kähler manifolds (Section II.2). We also study the maps f_* and f^* induced by a holomorphic map f (Section II.6). In particular, in Section II.4, we define the Kodaira-Spencer differential graded Lie algebra $A_X^{0,*}(\Theta_X)$ associated with a complex manifold X.

Section II.7 contains the proof of theorem A (Theorem II.7.3): we prove the existence of an isomorphism $\mathrm{Def}_{A_X^{0,*}(\Theta_X)} \cong \mathrm{Def}_X$ and so the Kodaira-Spencer algebra $A_X^{0,*}(\Theta_X)$ controls the infinitesimal deformation of X.

Chapter III is the technical bulk of this thesis. In Section III.1.2 we define the Maurer-Cartan functor $\mathrm{MC}_{(h,g)}$ and the deformation functor $\mathrm{Def}_{(h,g)}$ associated with a pair of morphisms of DGLAs $h : L \longrightarrow M$ and $g : N \longrightarrow M$. Sections III.1.3 and III.1.4 are devoted to the study of some properties of these functors as for example tangent and obstruction spaces.

In Section III.2 we introduce the extended deformation functors to prove the existence of a DGLA $H_{(h,g)}$ such that $\mathrm{Def}_{(h,g)} \cong \mathrm{Def}_{H_{(h,g)}}$ (Theorem III.2.36).

In *Chapter IV* we study the infinitesimal deformations of holomorphic maps.

Section IV.1 is devoted to defining the deformation functor $\mathrm{Def}(f)$ of infinitesimal deformations of a holomorphic map $f : X \longrightarrow Y$ of compact complex manifolds.

In Section IV.2, we prove theorem B, i.e., the existence of a pair of morphisms of DGLAs $h : A^{0,*}_{X\times Y}(\Theta_{X\times Y}(-log\,\Gamma)) \hookrightarrow A^{0,*}_{X\times Y}(\Theta_{X\times Y})$ and $g = (p^*, q^*) : A^{0,*}_X(\Theta_X) \times A^{0,*}_Y(\Theta_Y) \longrightarrow A^{0,*}_{X\times Y}(\Theta_{X\times Y})$, such that $\mathrm{Def}_{(h,g)} \cong \mathrm{Def}(f)$ (see Theorem IV.2.5).

Chapter V contains examples and applications of the techniques described before. In Section V.1 we study the infinitesimal deformations of holomorphic maps with fixed codomain and Section V.1.2 contains the main result about the semiregularity map (Corollary V.1.5).

Then we study infinitesimal deformations of a holomorphic map with fixed domain and codomain (Section V.3) and the infinitesimal deformations of an inclusion (Section V.4).

Acknowledgments. It is a pleasure for me to show my deep gratitude to the advisor of this thesis Prof. Marco Manetti, an excellent helpful professor who supports and encourages me every time. I'm indebted with him for many useful discussions, advices and suggestions. Several ideas of this work are grown under his influence. I wish also to thank him for having introduced me to the exciting subject of deformation theory.

I can not forget here my graduate study advisor Prof. Marialuisa J. de Resmini, who encouraged me to continue my studies and “sent” me to a summer school where I had my first meeting with Algebraic Geometry. I am very grateful for her improvements to the final version of this thesis.

I am also in debt with all those people with whom I have useful mathematical discussions or who make my staying in Rome pleasant.

In this set there certainly are Paolo Antonini, Sara Azzali, Francesco Esposito, Elena Martinengo, Silvia Montarani and Antonio Rapagnetta.

I wish also to thank three not mathematical friends of mine Arianna, Federica and Patrizia who support me in many and different ways.

This thesis is dedicated to my family Maria Teresa, Giovanni Michele, Enrico and especially to (the memory of) my grandmother Mina.

CHAPTER I

Functors of Artin rings

In this chapter we collect some definitions and main properties of deformation functors.

In the first section, we introduce the notions of functor of Artin rings, of deformation functors and we define the tangent and obstruction spaces.

Section I.2 is devoted to the study of the deformation functor Def_X of infinitesimal deformation of complex manifolds.

In Section I.3 we introduce the fundamental notions of differential graded Lie algebra (DGLA) L and of deformation functor associated with a DGLA Def_L.

The main references for this chapter are [**6**], [**20**], [**23**], [**31**] and [**32**].

I.1. Generalities on functors of Artin rings

Let $\mathbb{K}$ be a fixed field of characteristic zero.

Consider the following categories:

- **Set**: the category of sets in a fixed universe with $\{*\}$ a fixed set of cardinality 1;
- $\mathbf{Art} = \mathbf{Art}_{\mathbb{K}}$: the category of local Artinian $\mathbb{K}$-algebras with residue field $\mathbb{K}$ ($A/m_A = \mathbb{K}$);
- $\widehat{\mathbf{Art}} = \widehat{\mathbf{Art}_{\mathbb{K}}}$: the category of complete local Noetherian $\mathbb{K}$-algebras with residue field $\mathbb{K}$ ($A/m_A = \mathbb{K}$).

  For each $S \in \widehat{\mathbf{Art}}$ we also consider:
- $\mathbf{Art}_S$: the category of local Artinian S-algebras with residue field $\mathbb{K}$ (for such an element A, the structure morphism $S \longrightarrow A$ induces a trivial extension of the residue field $\mathbb{K}$);
- $\widehat{\mathbf{Art}_S}$: the category of complete local Noetherian S-algebras with residue field $\mathbb{K}$.

Remark I.1.1. We note that $\mathbf{Art}_S \subset \widehat{\mathbf{Art}_S}$. Moreover, by morphisms in a category of local objects we mean local morphisms and we often use the notation $A \in \mathbf{C}$ instead of $A \in ob(\mathbf{C})$, when $\mathbf{C}$ is a category.

If $\beta : B \longrightarrow A$ and $\gamma : C \longrightarrow A$ are morphisms in $\widehat{\mathbf{Art}_S}$ (in $\mathbf{Art}_S$, respectively), then

$$B \times_A C = \{(b, c) \in B \times C \mid \beta(b) = \gamma(c)\}$$

is the *fiber product* of β and γ and $B \times_A C \in \widehat{\mathbf{Art}_S}$ (in $\mathbf{Art}_S$, respectively).

Definition I.1.2. A *small extension* in $\widehat{\mathbf{Art}_S}$ (in $\mathbf{Art}_S$, respectively) is a short exact sequence

$$e: \quad 0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0,$$

where α is a morphism in $\widehat{\mathbf{Art}_S}$ (in $\mathbf{Art}_S$, respectively) and the kernel J is an ideal of B annihilated by the maximal ideal m_B, $m_B \cdot J = (0)$. This implies that J is $\mathbb{K}$-vector space.

A small extension is called *principal* if J is a one dimensional vector space ($J \cong \mathbb{K}$).

We will often say that a morphism $\alpha : B \longrightarrow A$ is a small extension, meaning that $0 \longrightarrow \ker(\alpha) \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$ is a small extension.

Remark I.1.3. Every surjective morphism in $\mathbf{Art}_S$ can be expressed as a finite composition of small extensions.

Actually, let $B \longrightarrow A$ be a surjection with kernel J:

$$0 \longrightarrow J \longrightarrow B \longrightarrow A \longrightarrow 0.$$

Since B is a local artinian ring, its maximal ideal m is nilpotent: there exists $n_0 \in \mathbb{N}$ such that $m^n = 0$ for each $n \geq n_0$; in particular, $m^n J = 0$.

Therefore, it is sufficient to consider the sequence of small extensions

$$0 \longrightarrow m^n J/m^{n+1}J \longrightarrow B/m^{n+1}J \longrightarrow B/m^n J \longrightarrow 0.$$

Remark I.1.4. In view of Remark I.1.3, it will be enough to check the surjection for small extensions instead of verifying the surjection for any morphism in $\mathbf{Art}_S$.

Definition I.1.5. A *functor of Artin rings* is a covariant functor $F : \mathbf{Art}_S \longrightarrow \mathbf{Set}$, such that $F(\mathbb{K}) = \{*\}$ is the one point set.

The functors of Artin rings $F : \mathbf{Art}_S \longrightarrow \mathbf{Set}$ and their natural transformations build a category denoted by $\mathbf{Fun}_S$. A natural transformation of functors $\gamma : F \longrightarrow G$ is an *isomorphism of functors* if and only if $\gamma(A) : F(A) \longrightarrow G(A)$ is bijective for each $A \in \mathbf{Art}_S$.

Example I.1.6. The trivial functor $F(A) = \{*\}$, for every $A \in \mathbf{Art}_S$.

Example I.1.7. Let $R \in \widehat{\mathbf{Art}_S}$. We define

$$h_R : \mathbf{Art}_S \longrightarrow \mathbf{Set},$$

such that

$$h_R(A) = \mathrm{Hom}_S(R, A).$$

$F \in \mathbf{Fun}_S$ is called *pro-representable* if it is isomorphic to h_R, for some $R \in \widehat{\mathbf{Art}_S}$.

If we can choose $R \in \mathbf{Art}_S$ then F is called *representable*.

Let $\mathbb{K}[\varepsilon]$, with $\varepsilon^2 = 0$, be the ring of dual numbers over $\mathbb{K}$. $\mathbb{K}[\varepsilon] = \mathbb{K} \oplus \mathbb{K}\varepsilon$ is a $\mathbb{K}$-vector space of dimension 2 and has a trivial S-algebra structure (induced by $S \longrightarrow \mathbb{K} \longrightarrow \mathbb{K}[\varepsilon]$).

Definition I.1.8. The set $t_F := F(\mathbb{K}[\varepsilon])$ is called the *tangent space* of $F \in \mathbf{Fun}_S$.

Let

$$\eta : F(B \times_A C) \longrightarrow F(B) \times_{F(A)} F(C) \tag{1}$$

be the map induced by the fiber product in $\mathbf{Art}_S$:

$$\begin{tikzcd}
B \times_A C \arrow[r] \arrow[d] & C \arrow[d] \\
B \arrow[r] & A.
\end{tikzcd}$$

Definition I.1.9. A functor F is called *homogeneous* if η is an isomorphism whenever $B \longrightarrow A$ is surjective.

Definition I.1.10. A functor F is called a *deformation functor* if

i) η is surjective whenever $B \longrightarrow A$ is surjective;
ii) η is an isomorphism whenever $A = \mathbb{K}$.

Remark I.1.11. The deformation functors will play an important role in this work.

In particular, we will study the following four deformation functors:

1) the functor Def_X of infinitesimal deformation of complex manifolds, in section I.2;
2) the functor Def_L associated with a differential graded Lie algebra L, in Section I.3;
3) the functor $\mathrm{Def}_{(h,g)}$ associated with a pair of morphisms of differential graded Lie algebras $h : L \longrightarrow M$ and $g : N \longrightarrow M$, in section III.1;
4) the functor Def_f associated with the infinitesimal deformations of a holomorphic map f, in section IV.1.

Example I.1.12. Let X be an algebraic scheme over $\mathbb{K}$ (separated of finite type over $\mathbb{K}$). Define the following functor

$$\mathrm{Def}_X : \mathbf{Art} \longrightarrow \mathbf{Set},$$

where $\mathrm{Def}_X(A)$ is the set of isomorphism classes of commutative diagram:

$$\begin{tikzcd}
X \arrow[r, "i"] \arrow[d] & X_A \arrow[d, "p_A"] \\
\mathrm{Spec}(\mathbb{K}) \arrow[r] & \mathrm{Spec}(A).
\end{tikzcd}$$

where i is a closed embedding and p_A is a flat morphism. Def_X is a deformation functor (see [**32**, Section 3.7] or [**31**, Prop. III.3.1]).

Proposition I.1.13. *Let F be a deformation functor. Then t_f has a natural structure of $\mathbb{K}$-vector space and every natural transformation of deformation functors $\phi : F \longrightarrow G$ induces a linear map between tangent spaces.*

Proof. Since $F(\mathbb{K})$ is just one point and the morphism η defined in (1) is an isomorphism for $A = \mathbb{K}$, we have $F(\mathbb{K}[\varepsilon]) \times F(\mathbb{K}[\varepsilon]) \cong F(\mathbb{K}[\varepsilon] \times_{\mathbb{K}} \mathbb{K}[\varepsilon])$.

Consider the map

$$+ : \mathbb{K}[\varepsilon] \times_{\mathbb{K}} \mathbb{K}[\varepsilon] \longrightarrow \mathbb{K}[\varepsilon],$$
$$(a + b\varepsilon, a + b'\varepsilon) \longmapsto a + (b + b')\varepsilon.$$

Then using the previous isomorphism, the map $+$ induces the addition on $F(\mathbb{K}[\varepsilon])$:

$$F(\mathbb{K}[\varepsilon]) \times F(\mathbb{K}[\varepsilon]) \xrightarrow{\cong} F(\mathbb{K}[\varepsilon] \times_{\mathbb{K}} \mathbb{K}[\varepsilon]) \xrightarrow{F(+)} F(\mathbb{K}[\varepsilon]).$$

Analogously, for the multiplication by a scalar $k \in \mathbb{K}$ we consider the map:

$$k : \mathbb{K}[\varepsilon] \longrightarrow \mathbb{K}[\varepsilon],$$
$$a + b\varepsilon \longmapsto a + (kb)\varepsilon.$$

□

Remark I.1.14. In the previous proposition we just used the fact that $F(\mathbb{K})$ is one point and that η in (1) is an isomorphism for $A = \mathbb{K}$ and $C = \mathbb{K}[\varepsilon]$

Definition I.1.15. A morphism $\phi : F \longrightarrow G$ in $\mathbf{Fun}_S$ is:

- *unramified* if $\phi : t_F \longrightarrow t_G$ is injective;
- *smooth* if the map

  $$F(B) \longrightarrow G(B) \times_{G(A)} F(A)$$

  induced by the diagram

  $$\begin{tikzcd}
  F(B) \arrow[r] \arrow[d, "\phi"'] & F(A) \arrow[d, "\phi"] \\
  G(B) \arrow[r] & G(A),
  \end{tikzcd}$$

  is surjective for every surjection $B \longrightarrow A$ in $\mathbf{Art}_S$;
- *étale* if ϕ is both smooth and unramified.

Remark I.1.16. If $\phi : F \longrightarrow G$ is smooth then, by taking $A = \mathbb{K}$ in the previous definition, we conclude that $\phi : F(B) \longrightarrow G(B)$ is surjective.

Remark I.1.17. If $\phi : F \longrightarrow G$ is an étale morphism, then it induces an isomorphism $\phi' : t_F \longrightarrow t_G$ on tangent spaces. Actually, ϕ is unramified and so ϕ' is injective. By hypothesis, ϕ is also smooth and so, applying Remark I.1.16 in the special case $B = \mathbb{K}[\varepsilon]$, ϕ' is surjective.

Definition I.1.18. A *functor F is smooth* if for every surjection $B \longrightarrow A$ of S-algebras $F(B) \longrightarrow F(A)$ is surjective (that is, the morphism $F \longrightarrow *$ is smooth).

Proposition I.1.19. *Let $\phi : F \longrightarrow G$ be an étale morphism of deformation functors. If G is homogeneous then ϕ is an isomorphism.*

Proof. Since ϕ is smooth, ϕ is surjective. Therefore, it is sufficient to prove the injectivity of ϕ using that G is homogeneous and ϕ is unramified.
The proof of this fact is taken from [**20**, Lemma 2.10]; it is given for the sake of completeness.
We prove it by induction on the length of A.

If $A = \mathbb{K}$ then $F(\mathbb{K}) = G(\mathbb{K}) = \{*\}$ and so the statement is obvious.

Let

$$0 \longrightarrow \mathbb{K}\varepsilon \longrightarrow B \longrightarrow A \longrightarrow 0,$$

be a principal small extension ($\varepsilon \cdot m_B = 0$). By induction $\phi : F(A) \longrightarrow G(A)$ is injective.

Consider the following isomorphism of S-algebras:

$$\varphi : B \times_{\mathbb{K}} \mathbb{K}[\varepsilon] \longrightarrow B \times_A B,$$

$$(b, \overline{b} + \beta\varepsilon) \longmapsto (b, b + \beta\varepsilon).$$

Since F is a deformation functor, $F(\varphi) : F(B) \times t_F \longrightarrow F(B) \times_{F(A)} F(B)$ is surjective.

Since G is also homogeneous, $G(\varphi) : G(B) \times t_G \longrightarrow G(B) \times_{G(A)} G(B)$ is an isomorphism. We note that $G(\varphi)(G(B) \times \{0\}) = \Delta$ diagonal.

Now, suppose that $\phi(\xi) = \phi(\eta) \in G(B)$ for ξ and $\eta \in F(B)$.

Since ϕ is injective on $F(A)$, $(\xi, \eta) \in F(B) \times_{F(A)} F(B)$.

Moreover, the surjectivity of $F(\varphi)$ implies the existence of an element $h \in t_F$ such that $F(\varphi)(\xi, h) = (\xi, \eta)$. Then $G(\varphi)(\phi(\xi), \phi(h)) = (\phi(\xi), \phi(\eta)) \in \Delta$ and so $\phi(h) = 0$ (G is an isomorphism).

By hypothesis, ϕ is unramified, therefore, $h = 0$ and so $\xi = \eta$. □

Remark I.1.20. Let F be a deformation functor and $f : B \longrightarrow A$ a surjection with $\ker f \cong \mathbb{K}$. Using the isomorphism $B \times_{\mathbb{K}} \mathbb{K}[\varepsilon] \cong B \times_A B$ of Proposition I.1.19, we obtain a commutative diagram

$$\begin{tikzcd}
F(B)\times t_F \arrow[r, "\pi"] \arrow[d, "\tau"'] & F(B) \arrow[d] \\
F(B) \arrow[r, "F(f)"'] & F(A).
\end{tikzcd}$$

π is the projection and τ defines an action of t_F on $F(B)$ which restricts to a transitive action on each fiber of $F(f)$ (see [**6**, Lemma 2.12]).

Corollary I.1.21. *Let F be a deformation functor, then $F = \{*\}$ if and only if $t_F = (0)$.*

Proof. One implication is obvious. Next, let F be a deformation functor such that $t_F = (0)$. We prove that $F(A) = \{*\}$ by induction on $dim_{\mathbb{K}}(A)$. If $A = \mathbb{K}$ then there is nothing to prove (by definition of functor of Artin rings). Next, let $\pi : B \longrightarrow A$ be a small extension and suppose that $F(A) = \{*\}$. By Remark I.1.20, $t_F = (0)$ acts transitively on the unique fiber $F(B)$ of the map $F(\pi)$. This implies $F(B) = \{*\}$. □

I.1.1. Obstruction theory.

Definition I.1.22. Let F be a functor of Artin rings; an *obstruction theory* for F is a pair (V, v_e) such that:

- V is a $\mathbb{K}$-vector space, called *obstruction space*;
- for every small extension in $\mathbf{Art}_S$

  $$e : 0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

  $v_e : F(A) \longrightarrow V \otimes_{\mathbb{K}} J$ is an obstruction map, satisfying the following properties:
  - If $\xi \in F(A)$ can be lifted to $F(B)$ then $v_e(\xi) = 0$.
  - For any morphism $\phi : e_1 \longrightarrow e_2$ of small extension, i.e.,

$$\begin{tikzcd}
e_1 : & 0 \arrow[r] & J_1 \arrow[r] \arrow[d, "\phi_J"] & B_1 \arrow[r] \arrow[d, "\phi_B"] & A_1 \arrow[r] \arrow[d, "\phi_A"] & 0 \\
e_2 : & 0 \arrow[r] & J_2 \arrow[r] & B_2 \arrow[r] & A_2 \arrow[r] & 0
\end{tikzcd} \tag{2}$$

    we have $v_{e_2}(\phi_A(a)) = (Id_V \otimes \phi_J)(v_{e_1}(a))$, for every $a \in F(A_1)$.

Definition I.1.23. An obstruction theory for a functor is *complete* if the lifting exists if and only if the obstruction vanishes.

Remark I.1.24. If F has (0) as complete obstruction space then F is smooth. In Proposition I.1.31 we will prove that the converse is also true for a deformation functor.

Remark I.1.25. Let $\psi : F \longrightarrow G$ be a natural transformation of functors and (V, v_e) an obstruction theory for G. Then $(V, v'_e := v_e \circ \psi)$ is an obstruction theory for F.

Actually, consider the small extension in $\mathbf{Art}_S$

$$e : 0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

and the map

$$v'_e : F(A) \xrightarrow{F(\psi)} G(A) \xrightarrow{v_e} V \otimes_{\mathbb{K}} J.$$

Let $\xi \in F(A)$ and suppose that it can be lifted to $\tilde{\xi} \in F(B)$. Therefore, $\psi(\xi) \in G(A)$ can be lifted to $\psi(\tilde{\xi}) \in G(B)$ and so $v'_e(\xi) = v_e(\psi(\xi)) = 0$.

Now, let $\phi : e_1 \longrightarrow e_2$ be a morphism of small extension as in (2). Then for each $a \in F(A_1)$ we have

$$(Id_V \otimes \phi_J)(v'_{e_1}(a)) = (Id_V \otimes \phi_J)(v_{e_1}(\psi(a))) =$$

$$v_{e_2}(\phi_A(\psi(a))) = v_{e_2}(\psi(\phi_A(a))) = v'_{e_2}(\phi_A((a))).$$

Moreover, if the morphism ψ is also smooth and (V, v_e) is complete for G, then (V, v'_e) is complete for F.

Indeed, suppose that $\xi \in F(A)$ is such that $0 = v'_e(\xi) = v_e(\psi(\xi))$. Then there exists $\eta \in G(B)$ that lifts $\psi(\xi) \in G(A)$ ((V, v_e) is complete for G). Consider the following diagram

$$\begin{tikzcd}
F(B) \arrow[r] \arrow[d] & F(A) \ni \xi \arrow[d] \\
\eta \in G(B) \arrow[r] & G(A) \ni \psi(\xi).
\end{tikzcd}$$

Since ψ is smooth, the map $F(B) \longrightarrow F(A) \times_{G(A)} G(B)$ is surjective and so there exists $\tilde{\xi} \in F(B)$ that lifts ξ.

Remark I.1.26. If V is complete a obstruction theory for a functor F, using embeddings of vector spaces we can construct infinitely many complete obstruction theories. Therefore, the goal is to find a "smallest" complete obstruction theory. The main results in this context is the following Theorem I.1.28. First of all, we give a definition.

Definition I.1.27. A morphism of obstruction theories $(V, v_e) \longrightarrow (W, w_e)$ is a linear map (of vector spaces) $\theta : V \longrightarrow W$ such that $w_e = \theta v_e$, for every small extensions e.

An obstruction theory (O_F, ob_e) for F is called *universal* if for any obstruction theory (V, v_e) there exists a unique morphism $(O_F, ob_e) \longrightarrow (V, v_e)$.

Theorem I.1.28. *(Fantechi, Manetti) Let F be a deformation functor. Then there exists a universal obstruction theory (O_F, ob_e) for F. Moreover, the universal obstruction theory is complete and every element of the vector space O_F is of the form $ob_e(\xi)$ for some principal extension*

$$e : \ 0 \longrightarrow \mathbb{K}\varepsilon \longrightarrow B \longrightarrow A \longrightarrow 0.$$

and some $\xi \in F(A)$.

Proof. See [**6**, Theorem 3.2 and Corollary 4.4]. □

Let $\phi : F \longrightarrow G$ be a morphism of deformation functors and (V, v_e), (W, w_e) obstructions theories for F and G, respectively. A linear map $\phi' : V \longrightarrow W$ is *compatible* with ϕ if $w_e\phi = \phi' v_e$ for every small extensions e.

Theorem I.1.29. *Let $\phi : F \longrightarrow G$ be a morphism of deformation functors and $\phi' : (V, v_e) \longrightarrow (W, w_e)$ a compatible morphism of obstruction theories. If (V, v_e) is complete, ϕ' injective and $t_F \longrightarrow t_G$ surjective, then ϕ is smooth.*

Proof. See [**20**, Proposition 2.17]. We have to prove that the map

$$F(B) \longrightarrow G(B) \times_{G(A)} F(A)$$

is surjective, for all small extensions

$$0 \longrightarrow \mathbb{K}\varepsilon \longrightarrow B \longrightarrow A \longrightarrow 0.$$

Let $(b', a) \in G(B) \times_{G(A)} F(A)$ and $a' \in G(A)$ their common image, that is, $b' \in G(B)$ lifts $a' \in G(A)$ and so $w_e(a') = 0$.

By hypothesis, ϕ' is injective and so $v_e(a) = 0$ $(0 = w_e(a') = w_e(\phi(a)) = \phi'(v_e(a)))$. Therefore, there exists $b \in F(B)$ that lifts a:

$$\begin{tikzcd}
b \in F(B) \arrow[r] \arrow[d] & F(A) \ni a \arrow[d] \\
b' \in G(B) \arrow[r] & G(A) \ni a'.
\end{tikzcd}$$

In general b does not lift b'. Let $b'' = \phi(b) \in G(B)$; then $(b'', b') \in G(B) \times_{G(A)} G(B)$.

As observed in the proof of Proposition I.1.19, we have an isomorphism $B \times_{\mathbb{K}} \mathbb{K}[\varepsilon] \cong B \times_A B$; since G is a deformation functor, there exists a surjective morphism

$$\alpha : G(B) \times t_G = G(B \times_A B) \longrightarrow G(B) \times_{G(A)} G(B).$$

Therefore, there exists $h \in t_G$ such that $(b'', \overline{b''} + h)$ is a lifting of $(b'', b'' + h = b')$.

By hypothesis, $t_F \longrightarrow t_G$ is surjective and so there exists a lifting $k \in t_F$ of $h \in t_G$. Taking $k + b \in F(B)$ produces a lifting of a that maps on b'.

□

Remark I.1.30. Let $\phi : F \longrightarrow G$ be a morphism of deformation functors and (O_F, v_e) and (O_G, w_e) their universal obstruction theories. Then $(O_G, w_e \circ \phi)$ is an obstruction theory for F. Thus, by Theorem I.1.28, there exists a morphism $o(\phi) : (O_F, v_e) \longrightarrow (O_G, w_e)$.

In conclusion, *every morphism of deformation functors induces a linear map both between tangent spaces and universal obstruction spaces.*

Next, we want to consider some useful properties between these morphisms.

PROPOSITION I.1.31. *Let* $\phi : F \longrightarrow G$ *be a morphism of deformation functors. Then* ϕ *is smooth if and only if* $t_F \longrightarrow t_G$ *is surjective and* $o(\phi) : O_F \longrightarrow O_G$ *is injective. In particular,* F *is smooth if and only if* $O_F = 0$.

PROOF. If $t_F \longrightarrow t_G$ is surjective and $o(\phi) : O_F \longrightarrow O_G$ is injective then ϕ is smooth by Theorem I.1.29.

Viceversa, suppose that ϕ is smooth, then by Remark I.1.16, $t_F = F(\mathbb{K}[\varepsilon]) \longrightarrow G(\mathbb{K}[\varepsilon]) = t_G$ is surjective. Let $B \longrightarrow A$ be a small extension and a an element of $F(A)$, with obstruction $x \in O_F$ such that $o(\phi)(x) = 0 \in O_G$. By Theorem I.1.28, O_G is complete and so $\phi(a) \in G(A)$ can be lifted to $b' \in G(B)$. Again, by Remark I.1.16, $F(B) \longrightarrow G(B)$ is surjective and so there exists $b \in F(B)$ that lifts a. Therefore, the obstruction of a is zero ($x = 0$) and this proves that $o(\phi)$ is injective. □

COROLLARY I.1.32. *A morphism of deformation functors* $\phi : F \longrightarrow G$ *is étale if and only if* $t_F \longrightarrow t_G$ *is bijective and* $o(\phi) : O_F \longrightarrow O_G$ *is injective.*

PROOF. If ϕ is étale, then by Remark I.1.17 $t_F \longrightarrow t_G$ is bijective. Since ϕ is also smooth, $o(\phi) : O_F \longrightarrow O_G$ is injective.

Conversely, by Proposition I.1.31 ϕ is smooth; by hypothesis, $t_F \longrightarrow t_G$ is injective, and so ϕ is also unramified. □

COROLLARY I.1.33. *Let* $\phi : F \longrightarrow G$ *be a morphism of deformation functors with* G *homogeneous. If* $t_F \longrightarrow t_G$ *is bijective and* $o(\phi) : O_F \longrightarrow O_G$ *is injective then* ϕ *is an isomorphism.*

PROOF. Put together Corollary I.1.32 and Proposition I.1.19. □

REMARK I.1.34. When G is not homogeneous, ϕ could be not injective and so we can just conclude the surjectivity of ϕ. Therefore, in these cases we prove the injectivity directly. This will happen in Theorems II.7.3 and IV.2.5.

COROLLARY I.1.35. *If* $\phi : F \longrightarrow G$ *is smooth then* $o(\phi) : O_F \longrightarrow O_G$ *is bijective.*

PROOF. By Proposition I.1.31, we have just to prove that $o(\phi) : O_F \longrightarrow O_G$ is surjective. Let $B \longrightarrow A$ be a small extension and $y \in O_G$ the obstruction to lifting $a' \in G(A)$ to $b' \in G(B)$. Since ϕ is smooth, there exists $a \in F(A)$, such that $\phi(a) = a'$, and $b \in F(B)$, such that $\phi(b) = b'$. Therefore, the obstruction $x \in O_F$ to lifting $a \in F(A)$ to $b \in F(B)$ is a lifting of $y \in O_G$.

□

I.2. Deformation functor of complex manifolds

In this section we study the infinitesimal deformation functor associated with a compact complex manifold. Thus we will work over the complex numbers and so $\mathbb{K} = \mathbb{C}$ and $\mathbf{Art} = \mathbf{Art}_{\mathbb{C}}$.

The main references are [**16**, Chapter 4], [**32**], [**31**, Chapter II].

Definition I.2.1. Let X be a compact complex manifold and $A \in \mathbf{Art}$. An *infinitesimal deformation* of X over $\operatorname{Spec}(A)$ is a commutative diagram of complex spaces

$$\begin{tikzcd}
X \arrow[r, "i"] \arrow[d] & X_A \arrow[d, "\pi"] \\
\operatorname{Spec}(\mathbb{C}) \arrow[r, "a"] & \operatorname{Spec}(A),
\end{tikzcd}$$

where π is a proper and flat holomorphic map, $a \in \operatorname{Spec}(A)$ is the closed point, i is a closed embedding and $X \cong X_A \times_{\operatorname{Spec}(A)} \operatorname{Spec}(\mathbb{C})$.

If $A = \mathbb{K}[\varepsilon]$ we call it a *first order* deformation of X.

Sometimes, for an infinitesimal deformation X_A over $\operatorname{Spec}(A)$, we also use the short notation $(X_A, \pi, \operatorname{Spec}(A))$.

Remark I.2.2. Let X_A be an infinitesimal deformation of X. We note that, by definition, it can be interpreted as a morphism of sheaves of algebras $\mathcal{O}_A \to \mathcal{O}_X$ such that $\mathcal{O}_A$ is flat over A and $\mathcal{O}_A \otimes_A \mathbb{C} \to \mathcal{O}_X$ is an isomorphism.

Given another deformation X'_A of X over $\operatorname{Spec}(A)$:

$$\begin{tikzcd}
X \arrow[r, "i'"] \arrow[d] & X'_A \arrow[d, "\pi'"] \\
\operatorname{Spec}(\mathbb{C}) \arrow[r, "a"] & \operatorname{Spec}(A),
\end{tikzcd}$$

we say that X_A and X'_A are *isomorphic* if there exists an isomorphism $\phi : X_A \longrightarrow X'_A$ over $\operatorname{Spec}(A)$, that induces the identity on X, that is, the following diagram is commutative

$$\begin{tikzcd}
 & X \arrow[dl, "i"'] \arrow[dr, "i'"] & \\
X_A \arrow[rr, "\phi"] \arrow[dr, "\pi"'] & & X'_A \arrow[dl, "\pi'"] \\
 & \operatorname{Spec}(A). &
\end{tikzcd}$$

We note that for every X we can always define the *infinitesimal product deformation*:

$$\begin{tikzcd}
X \arrow[r, "i"] \arrow[d] & X \times \operatorname{Spec}(A) \arrow[d] \\
\operatorname{Spec}(\mathbb{C}) \arrow[r, "a"] & \operatorname{Spec}(A).
\end{tikzcd}$$

Definition I.2.3. An infinitesimal deformation of X over $\operatorname{Spec}(A)$ is called *trivial* if it is isomorphic to the infinitesimal product deformation.

X is called *rigid* if every infinitesimal deformation of X over $\operatorname{Spec}(A)$ (for each $A \in \mathbf{Art}$) is trivial.

For every deformation X_A of X over $\operatorname{Spec}(A)$ and every morphism $A \longrightarrow B$ in $\mathbf{Art}$ ($\operatorname{Spec}(B) \longrightarrow \operatorname{Spec}(A)$), there exists an associated deformation of X over $\operatorname{Spec}(B)$, called *pull-back deformation*, induced by a basis change:

$$\begin{tikzcd}
X \arrow[r] \arrow[d] & X_A \times_{\operatorname{Spec}(A)} \operatorname{Spec}(B) \arrow[d] \\
\operatorname{Spec}(\mathbb{C}) \arrow[r] & \operatorname{Spec}(B).
\end{tikzcd}$$

Definition I.2.4. The *infinitesimal deformation functor* Def_X of a complex manifold X is defined as follows:

$$\operatorname{Def}_X : \mathbf{Art} \to \mathbf{Set},$$

$$A \longmapsto \operatorname{Def}_X(A) = \left\{ \begin{array}{c} \text{isomorphism classes of} \\ \text{infinitesimal deformations} \\ \text{of } X \text{ over } \operatorname{Spec}(A) \end{array} \right\}.$$

Proposition I.2.5. *Def_X is a deformation functor, i.e., it satisfies the conditions of Definition I.1.10.*

Proof. See [**32**, Section 3.7] or [**31**, Proposition III.3.1]. ☐

I.2.1. Tangent and obstruction spaces of Def_X. Let X be a compact complex manifold and Θ_X its holomorphic tangent bundle.

In this section we prove that the tangent space of Def_X is $\check{H}^1(X, \Theta_X)$ and that the obstruction space is naturally contained in $\check{H}^2(X, \Theta_X)$. First of all, we recall a useful lemma.

Lemma I.2.6. *Let B_0 be a $\mathbb{C}$-algebra and*

$$0 \longrightarrow J \longrightarrow B \longrightarrow A \longrightarrow 0$$

a small extension in $\mathbf{Art}$. Then there is a 1-1 correspondence

$$\left\{ \begin{array}{c} \textit{automorphisms of the trivial deformation } B_0 \otimes_{\mathbb{C}} B \\ \textit{inducing the identity on } B_0 \otimes_{\mathbb{C}} A \end{array} \right\} \longleftrightarrow \operatorname{Der}_{\mathbb{C}}(\mathrm{B}_0, \mathrm{B}_0) \otimes J$$

where the identity corresponds to the zero derivation, and the composition of automorphisms corresponds to the sum of derivations.

Proof. See [**31**, Lemma II.1.5] □

Remark I.2.7. Let U be a Stein open subset of a complex manifold X. Then the previous lemma is equivalent to saying that the following sequence is exact:

$$0 \longrightarrow \Gamma(U_i, \Theta_X) \otimes J \longrightarrow \operatorname{Aut}(\mathcal{O}_X(U_i) \otimes B) \longrightarrow \operatorname{Aut}(\mathcal{O}_X(U_i) \otimes A) \longrightarrow 0.$$

Moreover, we note that this is a central extension.

Theorem I.2.8. *Let X be a complex manifold. Then there is a 1-1 correspondence:*

$$k : \frac{\{\textit{first order deformations of } X\}}{\textit{isomorphism}} \longrightarrow \check{H}^1(X, \Theta_X),$$

called the Kodaira-Spencer correspondence, where $\Theta_X = \operatorname{Hom}(\Omega^1_X, \mathcal{O}_X) = \operatorname{Der}_{\mathbb{K}}(\mathcal{O}_X, \mathcal{O}_X)$, is the hololmorphic tangent bundle of X.

Moreover, $k(\xi) = 0$ if and only if ξ is the trivial deformation class.

Proof. For completeness we repeat this proof from [**31**, Proposition II.1.6] where all details are available.

Let X_ε be a first order deformation of X:

$$\begin{tikzcd} X \arrow[r] \arrow[d] & X_\varepsilon \arrow[d] \\ \operatorname{Spec}(\mathbb{C}) \arrow[r] & \operatorname{Spec}(\mathbb{C}[\varepsilon]), \end{tikzcd}$$

and $\mathcal{U} = \{U_i\}_{i \in I}$ be a Stein open cover of X such that $U_{ij} = U_i \cap U_j$ and $U_{ijk} = U_i \cap U_j \cap U_k$ are Stein for every i, j and $k \in I$.

For any open U_i the deformations $X_{\varepsilon|U_i}$ are trivial, then for each $i \in I$ there exist isomorphisms of deformations

$$\theta_i : U_i \times \operatorname{Spec}(\mathbb{C}[\varepsilon]) \longrightarrow X_{\varepsilon|U_i}.$$

Therefore, for all i and $j \in I$, the composition

$$\theta_{ij} = {\theta_i}^{-1}\theta_j : U_{ij} \times \operatorname{Spec}(\mathbb{C}[\varepsilon]) \longrightarrow U_{ij} \times \operatorname{Spec}(\mathbb{C}[\varepsilon])$$

is an automorphism of the trivial deformation $U_{ij} \times \operatorname{Spec}(\mathbb{C}[\varepsilon])$ of the Stein open subset U_{ij}.

Applying Lemma I.2.6, we conclude that there exists an element $d_{ij} \in \Gamma(U_{ij}, \Theta_X)$ corresponding to θ_{ij}, for all i and $j \in I$.

Moreover, on each U_{ijk} the following equality holds

$$\theta_{jk}{\theta_{ik}}^{-1}\theta_{ij}^{-1} = {\theta_j}^{-1}\theta_k{\theta_k}^{-1}\theta_i{\theta_i}^{-1}\theta_j = id_{|U_{ijk} \times \operatorname{Spec}(\mathbb{K}[\varepsilon])}.$$

Therefore, applying again Lemma I.2.6 yields

$$d_{jk} - d_{ik} + d_{ij} = 0,$$

that is, $\{d_{ij}\}$ is a Čech 1-cocycle and so it defines an element in $\check{H}^1(X, \Theta_X)$.

It can be checked that this element does not depend on the choice of the open cover $\mathcal{U}$.

Let X'_ε be another first order deformation

$$\begin{tikzcd}
X \arrow[r] \arrow[d] & X'_\varepsilon \arrow[d] \\
\mathrm{Spec}(\mathbb{C}) \arrow[r] & \mathrm{Spec}(\mathbb{C}[\varepsilon]),
\end{tikzcd}$$

and ϕ an isomorphism of deformations: $\phi : X_\varepsilon \longrightarrow X'_\varepsilon$.

Then, for each $i \in I$, there exists an induced automorphism

$$\alpha_i = {\theta'_i}^{-1} \circ \phi_{|U_i} \circ \theta_i :$$

$$U_i \times \mathrm{Spec}(\mathbb{C}[\varepsilon]) \xrightarrow{\theta_i} X_{\varepsilon|U_i} \xrightarrow{\phi_{|U_i}} X'_{\varepsilon|U_i} \xrightarrow{{\theta'_i}^{-1}} U_i \times \mathrm{Spec}(\mathbb{C}[\varepsilon])$$

and so a corresponding element $a_i \in \Gamma(U_i, \Theta_X)$.

Therefore, we have $\theta'_i\alpha_i = \phi_{|U_i}\theta_i$ and

$$(\theta'_i\alpha_i)^{-1}(\theta'_j\alpha_j) = {\theta_i}^{-1}\phi^{-1}_{|U_{ij}}\phi_{|U_{ij}}\theta_j = {\theta_i}^{-1}\theta_j.$$

This implies

$${\alpha_i}^{-1}{\theta'_{ij}}^{-1}\alpha_j = \theta_{ij}$$

or equivalently

$$d'_{ij} + a_j - a_i = d_{ij}.$$

In conclusion, the Čech cocycles $\{d_{ij}\}$ and $\{d'_{ij}\}$ are cohomologous and so they represent the same element in $\check{H}^1(X, \Theta_X)$.

Conversely, let $\theta \in \check{H}^1(X, \Theta_X)$ and $\{d_{ij}\} \in Z^1(\mathcal{U}, \Theta_X)$ be a representative of θ with respect to an open Stein cover $\{\mathcal{U}\}$. By Lemma I.2.6, we can associate with each d_{ij} an automorphism θ_{ij} of the trivial deformation $U_{ij} \times \mathrm{Spec}(\mathbb{C}[\varepsilon])$. Since the element $\{d_{ij}\}$ satisfies the cocycle condition, θ_{ij} satisfies this condition too:

$$\theta_{jk}{\theta_{ik}}^{-1}\theta_{ij}^{-1} = id_{|U_{ijk}\times \mathrm{Spec}(\mathbb{C}[\varepsilon])}.$$

Using these automorphisms we can glue together the schemes $U_i \times \mathrm{Spec}(\mathbb{C}[\varepsilon])$ (see [**13**, pag. 69]) to obtain a scheme X_ε that is a first order deformation of X.

At this point the last assertion is clear.

□

THEOREM I.2.9. *$\check{H}^2(X, \Theta_X)$ is a complete obstruction space for* Def_X.

PROOF. For completeness, we take the proof from [**31**, Proposition II.1.8] where details are available.

Let $\mathcal{U} = \{U_i\}_{i\in I}$ be an open Stein cover of X such that U_{ij} and U_{ijk} are Stein for all, i, j and $k \in I$ and

$$0 \longrightarrow J \longrightarrow B \longrightarrow A \longrightarrow 0$$

be a small extension in **Art**.

Let X_A be an infinitesimal deformation of X over $\mathrm{Spec}(A)$. Then we have isomorphisms

$$\theta_i : U_i \times \mathrm{Spec}(A) \longrightarrow X_{A|U_i}$$

such that $\theta_{ij} := {\theta_i}^{-1}\theta_j$ are automorphisms of the trivial deformations $U_{ij} \times \mathrm{Spec}(A)$ and $\theta_{jk}{\theta_{ik}}^{-1}\theta_{ij} = id_{|U_{ijk}\times\mathrm{Spec}(A)}$.

To define a deformation X_B that lifts the deformation X_A is necessary and sufficient to give automorphisms $\{\tilde{\theta}_{ij}\}$ of the trivial deformation $U_{ij} \times \mathrm{Spec}(B)$ such that

i) $\{\tilde{\theta}_{ij}\}$ glues together: $\tilde{\theta}_{jk}\tilde{\theta}_{ik}^{-1}\tilde{\theta}_{ij} = id_{|U_{ijk}\times\mathrm{Spec}(B)}$,
ii) $\{\tilde{\theta}_{ij}\}$ lifts $\{\theta_{ij}\}$: $\tilde{\theta}_{ij}$ restricts to θ_{ij} on $U_{ij} \times \mathrm{Spec}(A)$.

Let us choose automorphisms $\{\tilde{\theta}_{ij}\}$ that satisfy condition *ii*). Then the automorphisms

$$\tilde{\theta}_{ijk} = \tilde{\theta}_{jk}\tilde{\theta}_{ik}^{-1}\tilde{\theta}_{ij}$$

are automorphisms of the trivial deformation that restrict to the identity on $U_{ijk}\times\mathrm{Spec}(A)$. By Lemma I.2.6, there exists $\{\tilde{d}_{ijk}\} \in \Gamma(U_{ijk}, \Theta_X)\otimes J$ that corresponds to $\tilde{\theta}_{ijk}$. An easy calculation shows that $\{\tilde{d}_{ijk}\}$ is a Čech cocylce and so $\{\tilde{d}_{ijk}\} \in Z^2(\mathcal{U}, \Theta_X) \otimes J$.

Next, let $\{\overline{\theta}_{ij}\}$ be different automorphisms of the trivial deformations $U_{ij} \times \mathrm{Spec}(B)$ that satisfy condition *ii*). As above, let $\overline{d}_{ijk}$ be the derivations corresponding to $\overline{\theta}_{ijk}$.

The automorphisms $\overline{\theta}_{ij}\tilde{\theta}_{ij}^{-1}$ of $U_{ij} \times \mathrm{Spec}(B)$ restrict to the identity on $U_{ij} \times \mathrm{Spec}(A)$ for each U_{ij} and so, again by Lemma I.2.6, they correspond to some $\{d_{ij}\} \in \Gamma(U_{ij}, \Theta_X) \otimes J$.

Therefore,

$$\overline{d}_{ijk} = \tilde{d}_{ijk} + d_{jk} - d_{ik} + d_{ij}.$$

This implies that the Čech cocycles $\{\tilde{d}_{ijk}\}$ and $\{\overline{d}_{ijk}\}$ are cohomologous and so their cohomology classes coincide in a well defined element $v_e(X_A)$ in $\check{H}^2(X, \Theta_X) \otimes J$:

$$v_e(X_A) := [\{\tilde{d}_{ijk}\}] = [\{\overline{d}_{ijk}\}] \in \check{H}^2(X, \Theta_X) \otimes J.$$

Moreover, the class $v_e(X_A)$ is zero if and only if the collection of automorphisms also satisfies condition *i*) and it is equivalent to the existence of a lifting X_B of the deformation X_A.

$\square$

I.3. DGLAs and deformation functor

In this section we study the deformation functor associated with a differential graded Lie algebra (DGLA).

In particular, we give the fundamental definition of a DGLA (Definition I.3.5).

We also introduce the Maurer-Cartan functor MC_L (Definition I.3.17) and the deformation functor Def_L associated with a DGLA L (Definition I.3.30).

We start by defining the differential graded vector spaces.

I.3.1. Differential graded vector spaces. Let $\mathbb{K}$ be a fixed field of characteristic 0. Unless otherwise stated, all vector spaces, linear maps, tensor products etc. are intended over $\mathbb{K}$.

Every *graded vector space* is a $\mathbb{Z}$-graded vector space (over $\mathbb{K}$). If $V = \oplus_i V^i$ is a graded vector space and $a \in V$ is a homogeneous element, then we denote by $\deg_V(a)$ the degree of a in V; we will also use the notation $\deg(a) = \bar{a}$, when V is clear from the context.

The morphisms of graded vector space are degree preserving linear maps.

Given two graded vector spaces V and W, we define $\mathrm{Hom}^n_{\mathbb{K}}(V, W)$ as the vector space of $\mathbb{K}$-linear maps $f : V \longrightarrow W$, such that $f(V^i) \subset W^{i+n}$, for each $i \in \mathbb{Z}$.

Let V be a graded vector space, then $V[n]$ is the complex V with degrees shifted by n. More precisely, for $\mathbb{K}[n]$ we have

$$\mathbb{K}[n]^i = \begin{cases} \mathbb{K} & \text{if } i + n = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then $V[n] = \mathbb{K}[n] \otimes V$, which implies $V[n]^i = V^{i+n}$.

Remark I.3.1. We note that there exist isomorphisms

$$\mathrm{Hom}^n_{\mathbb{K}}(V, W) \cong \mathrm{Hom}^0_{\mathbb{K}}(V[-n], W) \cong \mathrm{Hom}^0_{\mathbb{K}}(V, W[n]).$$

A *differential graded vector space* is a pair (V, d) where $V = \oplus V^i$ is a graded vector space and d is a differential of degree 1 ($d : V^i \longrightarrow V^{i+1}$ and $d \circ d = 0$).

For every differential graded vector space (V, d) we use the standard notation $Z^i(V) = \ker(d\colon V^i \to V^{i+1})$, $B^i(V) = \mathrm{Im}(d\colon V^{i-1} \to V^i)$ and $H^i(V) = Z^i(V)/B^i(V)$.

A morphism of differential graded vector spaces is a degree preserving linear map that commutes with the differentials.

A morphism is a *quasi-isomorphism* if it induces isomorphisms in cohomology.

Example I.3.2. Given (V, d), then for each $i \in \mathbb{Z}$, the shifted differential graded vector space $(V[i], d_{[i]})$ is defined as:

$$V[i] = \bigoplus_j V[i]^j = \bigoplus_j V^{i+j} \qquad \text{and} \qquad d_{[i]} = (-1)^i d.$$

Example I.3.3. If (V, d_V) and (W, d_W) are differential graded vector spaces, then we can define a new differential graded vector space

$$\mathrm{Hom}^*(V, W) = \bigoplus_{n\in\mathbb{Z}} \mathrm{Hom}^n_{\mathbb{K}}(V, W)$$

with natural differential d' given by

$$d'(f) := d_W f - (-1)^{\deg(f)} f\, d_V.$$

Moreover, for each i, there exists the following isomorphism

$$H^i(\mathrm{Hom}^*(V, W)) \cong \mathrm{Hom}^i(H^*(V), H^*(W)).$$

Example I.3.4. Given (V, d_V) and (W, d_W), we can also define the following differential graded vector space

$$\mathrm{Htp}(V, W) = \mathrm{Hom}^*(V[1], W) = \bigoplus_i \mathrm{Htp}^i(V, W),$$

with

$$\mathrm{Htp}^i(V, W) = \mathrm{Hom}^i(V[1], W) = \mathrm{Hom}^{i-1}(V, W)$$

and differential δ:

$$\delta(f) = d_W(f) - (-1)^i f d_{V[1]} = d_W f + (-1)^i f d_V \qquad \forall\, f \in \mathrm{Htp}^i(V, W).$$

We will use these differential graded vector spaces in the last chapter (Section V.1.1).

I.3.2. Differential graded Lie algebras (DGLAs).

Definition I.3.5. A *differential graded Lie algebra* (DGLA for short) is a triple $(L, [\,,\,], d)$, where $(L = \bigoplus_{i\in\mathbb{Z}} L^i, d)$ is a differential graded vector space and $[\,,\,] : L \times L \to L$ is a bilinear map, called bracket, satisfying the following conditions:

1. the bracket $[\,,\,]$ is homogeneous and graded skewsymmetric; i.e., $[L^i, L^j] \subset L^{i+j}$ and $[a, b] + (-1)^{\deg(a)\deg(b)}[b, a] = 0$, for every homogeneous a and b.
2. Every triple of homogeneous elements satisfies the graded Jacobi identity
$$[a, [b, c]] = [[a, b], c] + (-1)^{\deg(a)\deg(b)}[b, [a, c]].$$
3. $d(L^i) \subseteq L^{i+1}$, $d \circ d = 0$ and
$$d[a, b] = [da, b] + (-1)^{\deg(a)}[a, db].$$

The last property is called Leibniz's rule and, in particular, it implies that the bracket induces a structure of differential graded Lie algebra (with zero differential) on the cohomology $H^*(L) = \oplus_i H^i(L)$ of a DGLA L.

Example I.3.6. If $L = \oplus L^i$ is a DGLA, then L^0 is a Lie algebra in the usual sense; vice-versa, every Lie algebra is a differential graded Lie algebra concentrated to degree 0.

Remark I.3.7. If the degree of a is even then $[a, a] = 0$ (graded skew-symmetry). If the degree of a is odd, then $[a, [a, b]] = \frac{1}{2}[[a, a], b]$ for all $b \in L$ (graded Jacobi). In particular, $[a, [a, a]] = 0$; moreover, $[a, b] = [b, a]$ for all a and $b \in L$ of odd degree.

Definition I.3.8. A morphism of differential graded Lie algebras $\varphi : L \longrightarrow M$ is a linear map that preserves degrees and commutes with brackets and differentials; written in details we have

- $\varphi(L^i) \subseteq M^i$, for each i;
- $\varphi(d_L a) = d_M(\varphi(a))$, for each $a \in L$;
- $\varphi([a, b]) = [\varphi(a), \varphi(b)]$, for each $a, b \in L$.

A *quasi-isomorphism* of DGLAs is a morphism that induces isomorphisms in cohomology. Two DGLAs L and M are *quasi-isomorphic* if they are equivalent under the equivalence relation $\sim$ defined by: $L \sim M$ if there exists a quasi-isomorphism $\phi : L \longrightarrow M$.

A DGLA L is *formal* if it is quasi-isomorphic to its cohomology graded vector space $H^*(L)$.

Remark I.3.9. The following DGLAs are isomorphic:

$$(L, [\, , \,], d) \cong (L, -[\, , \,], d) \cong (L, [\, , \,], -d) \cong (L, -[\, , \,], -d).$$

Actually, the morphism $\varphi = -id$ gives an isomorphism between $(L, [\, , \,], d)$ and $(L, -[\, , \,], d)$, whereas $\varphi(-) = (-1)^{\deg(-)} id$ is an isomorphism between $(L, [\, , \,], d)$ and $(L, [\, , \,], -d)$.

Definition I.3.10. A linear map $f : L \longrightarrow L$ is called a *derivation* of degree n if $f(L^i) \subset L^{i+n}$ and it satisfies the graded Leibniz rule:

$$f([a, b]) = [f(a), b] + (-1)^{n\bar{a}}[a, f(b)].$$

The graded Leibniz rule implies that the differential d is a derivation of degree 1.

Remark I.3.11. Let $a \in L^i$ and consider the operator

$$[a, \,] : L \longrightarrow L,$$

$$[a, \,](x) = [a, x], \qquad \forall\, x \in L.$$

Then, the graded Jacobi identity implies that $[a, \,]$ is a derivation of degree $i = deg(a)$.

Example I.3.12. Let L be a DGLA and consider the vector space decomposition $L^1 = N^1 \oplus B^1(L)$. Then the graded vector space $N = \oplus N^i$ with

$$\begin{cases} N^i = 0 & \text{for } i \leq 0 \\ N^1 = N^1 & \text{for } i = 1 \\ N^i = L^i & \text{for } i \geq 2 \end{cases}$$

is a sub-DGLA of L.

Example I.3.13. Given a DGLA $(L = \oplus L^i, [\ ,\ ], d)$ we can associate with it a new DGLA $(L' = \oplus L'^i, [\ ,\ ]', d')$ where

$$\begin{cases} L'^i = L^i & \text{for } i \neq 1 \\ L'^1 = L^1 \oplus \mathbb{K}\, d & \text{for } i = 1, \end{cases}$$

$$[v + ad, w + bd]' = [v, w] + a\, d(w) - (-1)^{\deg(v)} b\, d(v)$$

and

$$d'(v + a\, d) = [d, v + ad]' = dv,$$

for any $v, w \in L^i$ and $a, b \in \mathbb{K}$.

Example I.3.14. Let M be a DGLA. Then $M[t, dt] = M \otimes \mathbb{K}[t, dt]$ is a DGLA, where $\mathbb{K}[t, dt]$ is the differential graded algebra of polynomial differential forms over the affine line. More precisely, $\mathbb{K}[t, dt] = \mathbb{K}[t] \oplus \mathbb{K}[t]dt$, where t has degree 0 and dt has degree 1. As vector space $M[t, dt]$ is generated by elements of the form $mp(t) + nq(t)dt$, with $m, n \in M$ and $p(t), q(t) \in \mathbb{K}[t]$. The differential and the bracket on $M[t, dt]$ are defined as follows:

$$d(mp(t) + nq(t)dt) = (dm)p(t) + (-1)^{\deg(m)} mp'(t)dt + (dn)q(t)dt,$$

$$[mp(t), nq(t)] = [m, n]p(t)q(t), \quad [mp(t), nq(t)dt] = [m, n]p(t)q(t)dt.$$

For every $a \in \mathbb{K}$ define the *evaluation morphism* in the following way

$$e_a : M[t, dt] \longrightarrow M,$$

$$e_a(\sum m_i t^i + n_i t^i dt) = \sum m_i a^i.$$

The evaluation morphism is a morphism of DGLAs which is a left inverse of the inclusion and is a surjective quasi-isomorphism for each a.

Example I.3.15. If L is a DGLA and B is a commutative $\mathbb{K}$-algebra then $L \otimes B$ has a natural structure of DGLA, given by

$$[l \otimes a, m \otimes b] = [l, m] \otimes ab;$$

$$d(l \otimes a) = dl \otimes a.$$

If B is also nilpotent (for example $B = m_A$ the maximal ideal of a local artinian $\mathbb{K}$-algebra A) then $L \otimes B$ is a nilpotent DGLA. Therefore, for every $a \in L^0 \otimes B$, we can define an automorphism of the DGLA $L \otimes B$:

$$e^{[a,-]} := \sum_{n \geq 0} \frac{[a, -]^n}{n!} \ : L \otimes B \longrightarrow L \otimes B,$$

where

$$[a, -] : L \otimes B \longrightarrow L \otimes B,$$

$$[a, -](b) := [a, b]$$

is a nilpotent derivation of degree zero (since $[a,-]([b,c]) = [[a,-](b),c] + [b,[a,-](c)]$).

Example I.3.16. Let (L,d) be a DGLA and $\operatorname{Der}^i(L,L)$ the space of derivations of L of degree i. Then $\operatorname{Der}^*(L,L) = \bigoplus_i \operatorname{Der}^i(L,L)$ is a DGLA with bracket

$$[f,g] = fg - (-1)^{\overline{f}\,\overline{g}} gf,$$

and differential d' given by

$$d'(f) = [d,f].$$

I.3.3. Maurer-Cartan functor associated with a DGLA.

Definition I.3.17. The *Maurer-Cartan equation* in a differential graded Lie algebra L is

$$dx + \frac{1}{2}[x,x] = 0, \qquad x \in L^1.$$

The solutions of this equation are called the *Maurer-Cartan elements* of the DGLA L.

Remark I.3.18. Let $x \in L^1$ be an element of degree one in the DGLA L and consider the operator of degree one

$$d(-) + [x,-] : L \longrightarrow L,$$

$$(d(-) + [x,-])(a) = da + [x,a] \qquad \forall\ a \in L.$$

Then, $d(-)+[x,-]$ is a differential ($(d(-)+[x,-])^2 = 0$) if and only if x satisfies the Maurer-Cartan equation. Indeed, using Remark I.3.7,

$$(d(-) + [x,-])^2(a) = (d(-) + [x,-])(da + [x,a]) =$$
$$d^2a + d[x,a] + [x,da] + [x,[x,a]] =$$
$$[dx,a] + (-1)^{deg(x)}[x,da] + [x,da] + \frac{1}{2}[[x,x],a] =$$
$$[dx + \frac{1}{2}[x,x],a].$$

Remark I.3.19. Let L' be the DGLA of Example I.3.13 (with $L'^1 = L^1 \oplus \mathbb{K}\, d$) then

$$dx + \frac{1}{2}[x,x] = 0 \quad \text{if and only if} \quad [x+d, x+d]' = 0.$$

The previous definition led to the following definition of the Maurer-Cartan functor.

Definition I.3.20. Let L be a DGLA; then the *Maurer-Cartan functor associated with L* is

$$MC_L : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$MC_L(A) = \{x \in L^1 \otimes m_A \mid dx + \frac{1}{2}[x,x] = 0\}.$$

Remark I.3.21. A morphism $\phi : L \longrightarrow M$ of DGLA preserves bracket and differential, therefore, it induces a morphism of functors $\phi : \mathrm{MC}_L \longrightarrow \mathrm{MC}_M$.

Remark I.3.22. We note that MC_L is a *homogeneous* functor, since $\mathrm{MC}_L(B \times_A C) \cong \mathrm{MC}_L(B) \times_{\mathrm{MC}_L(A)} \mathrm{MC}_L(C)$ for each pair of morphisms $\gamma : C \longrightarrow A$ and $\beta : B \longrightarrow A$, with β surjective.

Remark I.3.23. By definition, the *tangent space* of MC_L is:

$$t_{\mathrm{MC}_L} := \mathrm{MC}_L(\mathbb{K}[\varepsilon]) = \{x \in L^1 \otimes \mathbb{K}\varepsilon \mid dx + \frac{1}{2}[x, x] = 0\} \cong$$
$$\{x \in L^1 \mid dx = 0\} = Z^1(L).$$

Lemma I.3.24. *$H^2(L)$ is a complete obstruction space for* MC_L.

Proof. Let

$$e : \qquad 0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

be a small extension in **Art** and $x \in \mathrm{MC}_L(A)$.

We want to define a map $v_e : \mathrm{MC}_L(A) \longrightarrow H^2(L) \otimes J$.

Let $\tilde{x} \in L^1 \otimes m_B$ be a lifting of x and define

$$h = d\tilde{x} + \frac{1}{2}[\tilde{x}, \tilde{x}] \in L^2 \otimes m_B.$$

In general, $\tilde{x}$ does not satisfy the Maurer-Cartan equation and so h is in general different from zero.

It turns out that $\alpha(h) = dx + \frac{1}{2}[x, x] = 0$ and so $h \in L^2 \otimes J$. Moreover,

$$dh = d^2\tilde{x} + \frac{1}{2}[d\tilde{x}, \tilde{x}] - \frac{1}{2}[\tilde{x}, d\tilde{x}] =^{\mathrm{a}}$$
$$= [d\tilde{x}, \tilde{x}] = [h, \tilde{x}] - \frac{1}{2}[[\tilde{x}, \tilde{x}], \tilde{x}].$$

By definition, $[h, \tilde{x}] \in [L^2 \otimes J, L^1 \otimes m_B] = 0$ (e is a small extension) and, by Remark I.3.7, $[[\tilde{x}, \tilde{x}], \tilde{x}] = 0$.

In conclusion, $dh = 0$ and so $h \in H^2(L) \otimes J$.

We note that this class does not depend on the choice of the lifting $\tilde{x}$. Indeed, let $y \in L^1 \otimes m_B$ be another lifting of x: $\alpha(y) = \alpha(\tilde{x}) = x$. Then $y = \tilde{x} + t$ for some $t \in L^1 \otimes J$. Using $[L^1 \otimes J, L^1 \otimes m_B] = 0$, we have

$$h' = dy + \frac{1}{2}[y, y] = d\tilde{x} + dt + \frac{1}{2}[\tilde{x} + t, \tilde{x} + t] = d\tilde{x} + \frac{1}{2}[\tilde{x}, \tilde{x}] + dt = h + dt$$

and so h and h' represent the same class in $H^2(L) \otimes J$.

Therefore, the following obstruction map is well defined

$$v_e : \mathrm{MC}_L(A) \longrightarrow H^2(L) \otimes J,$$

---

[a] We have $-\frac{1}{2}[\tilde{x}, d\tilde{x}] = -\frac{1}{2}(-(-1)^2[d\tilde{x}, \tilde{x}]) = \frac{1}{2}[d\tilde{x}, \tilde{x}]$.

$$x \longmapsto v_e(x) = [h].$$

If $[h] = 0$ then $h = dq$ for some $q \in L^1 \otimes J$. This implies that $\overline{x} = \tilde{x} - q$ is a lifting of x that satisfies the Maurer-Cartan equation, i.e.,

$$d\overline{x} + \frac{1}{2}[\overline{x}, \overline{x}] = d\tilde{x} - dq + \frac{1}{2}[\tilde{x} - q, \tilde{x} - q] = d\tilde{x} - dq + \frac{1}{2}[\tilde{x}, \tilde{x}] = h - dq = 0.$$

Thus, v_e satisfies condition 1 of Definition I.1.22 of obstruction theory. The other property (change of basis) is an easy calculation.

If $x \in \mathrm{MC}_L(A)$ can be lifted to $x' \in MC_L(B)$ then $[h] = 0$.

In conclusion, $(H^2(L), v_e)$ is a complete obstruction theory for MC_L.

□

Remark I.3.25. (**About smoothness**)

If $H^2(L) = 0$ then MC_L is smooth.

If L is abelian then MC_L is *smooth*. Actually, in this case, $\mathrm{MC}_L(A) = Z^1(L) \otimes m_A$. Moreover, if $B \twoheadrightarrow A$ then $Z^1(L) \otimes m_A \twoheadrightarrow Z^1(L) \otimes m_B$.

I.3.4. Gauge action.

Definition I.3.26. Two elements x and $y \in L^1 \otimes m_A$ are said to be *gauge equivalent* if there exists $a \in L^0 \otimes m_A$ such that

$$y = e^a * x := x + \sum_{n \geq 0} \frac{[a, -]^n}{(n+1)!}([a, x] - da).$$

The operator $*$ is called the gauge action of the group $\exp(L^0 \otimes m_A)$ on $L \otimes m_A$; indeed $e^a * e^b * x = e^{a \bullet b} * x$, where $\bullet$[b] is the Baker-Campbell-Hausdorff product in the nilpotent DGLA $L \otimes m_A$.

Remark I.3.27. For a better understanding of the gauge action, it is convenient to consider the DGLA L' of Example I.3.13 (with $L'^1 = L^1 \oplus \mathbb{K}\, d$) and the affine embedding

$$\phi : L^1 \longrightarrow L'^1, \qquad \phi(x) = x + d \qquad \forall\, x \in L^1.$$

As already observed, $dx + \frac{1}{2}[x, x] = 0$ if and only if $[\phi(x), \phi(x)]' = 0$.

As in Example I.3.15, for each $A \in \mathbf{Art}$ and $a \in L'^0 \otimes m_A$, we can consider the exponential of the adjoint action $e^{[a,]'} : L'^1 \otimes m_A \longrightarrow L'^1 \otimes m_A$. Using the embedding ϕ this action induces the gauge action of $L^0 \otimes m_A$ on $L^1 \otimes m_A$. Actually, for any $a \in L^0 \otimes m_A$ and $x \in L^1 \otimes m_A$

$$\phi^{-1}(e^{[a,]'}\phi(x)) = e^{[a,]'}(x + d) - d =$$

$$x + \sum_{n \geq 1} \frac{[a, -]'^n}{(n)!}(x + d) = x + \sum_{n \geq 0} \frac{[a, -]'^{n+1}}{(n+1)!}(x + d) =$$

$$x + \sum_{n \geq 0} \frac{[a, -]'^n}{(n+1)!}([a, x] - da) = e^a * x.$$

---

[b] $a \bullet b = a + b + \frac{1}{2}[a, b] + \frac{1}{12}[a, [a, b]] - \frac{1}{12}[b, [b, a]] + \cdots$

Example I.3.28. Let J be an ideal of $A \in \mathbf{Art}$ ($J \subset m_A$) such that $J \cdot m_A = 0$ (for example, J may be the kernel of a small extension).

If $x \in L^1 \otimes J$ then, for each $a \in L^0 \otimes m_A$ we have

$$e^a * x = x + \sum_{n=0}^{\infty} \frac{[a,-]^n}{(n+1)!}([a,x] - da) = x + \sum_{n=0}^{\infty} \frac{[a,-]^n}{(n+1)!}(-da) = x + e^a * 0,$$

or, in general, if $y \in L^1 \otimes m_A$ then $e^a * (x+y) = x + e^a * y$.

If $a \in L^0 \otimes J$ then, for each $x \in L^1 \otimes m_A$:

$$e^a * x = x + \sum_{n=0}^{\infty} \frac{[a,-]^n}{(n+1)!}([a,x] - da) = x - da.$$

In general, if $b \in L^0 \otimes m_A$, then $e^{a+b} * x = e^b * x - da$.

We note that

$$e^a * x = x \qquad \text{if and only if} \qquad [a,x] = da. \tag{3}$$

Actually, $e^a * x = x$ if and only if $0 = \dfrac{e^{[a,-]} - id}{[a,-]}([a,x] - da)$. Applying the inverse of the operator $\dfrac{e^{[a,-]} - id}{[a,-]}$, yields $e^a * x = x$ if and only if $[a,x] - da = 0$.

Remark I.3.29. The solutions of the Maurer-Cartan equation are preserved under the gauge action.

Actually, we have

$$d(e^a * x) = d'(e^{[a,]'}(d+x) - d) = [d, e^{[a,]'}(d+x) - d]' = [d, e^{[a,]'}(d+x)]'$$

and using Remark I.3.19

$$[e^a * x, e^a * x] = [e^{[a,]'}(d+x) - d, e^{[a,]'}(d+x) - d]' =$$
$$[e^{[a,]'}(d+x), e^{[a,]'}(d+x)]' - 2[d, e^{[a,]'}(d+x)]' =$$
$$e^{[a,]'}[d+x, d+x]' - 2[d, e^{[a,]'}(d+x)]' = -2[d, e^{[a,]'}(d+x)]'.$$

Therefore,

$$d(e^a * x) + \frac{1}{2}[e^a * x, e^a * x] = 0.$$

Finally, for each $x \in \mathrm{MC}_L(A)$, we define the *irrelevant stabilizer* of x:

$$Stab_A(x) = \{e^{dh + [x,h]} \mid h \in L^{-1} \otimes m_A\} \subset \exp(L^0 \otimes A).$$

The name irrelevant stabilizer is due to the fact that $e^{dh+[x,h]} * x = x$. Actually, $dh + [x,h]$ satisfies condition (3), i.e.,

$$[dh + [x,h], x] = [dh, x] + [[x,h], x] = d[h,x] + [h, dx] + \frac{1}{2}[h, [x,x]] =$$
$$d[h,x] + [h, dx + \frac{1}{2}[x,x]] = d[h,x] = d(dh + [x,h]).$$

Moreover, we observe that $Stab_A(x)$ is a subgroup of $\exp(L^0 \otimes A)$ and that for any $a \in L^0 \otimes A$

$$e^a Stab_A(x) e^{-a} = Stab_A(y), \qquad \text{with} \qquad y = e^a * x.$$

I.3.5. Deformation functor associated with a DGLA.

Definition I.3.30. The *deformation functor* associated with a differential graded Lie algebra L is:

$$\mathrm{Def}_L : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{Def}_L(A) = \frac{MC_L}{\exp\left(L^0 \otimes m_A\right)}.$$

In this case too a morphism of DGLAs $\phi : L \longrightarrow M$ induces a morphism of the associated functors $\phi : \mathrm{Def}_L \longrightarrow \mathrm{Def}_M$.

The name deformation functor is justified by the following proposition.

Proposition I.3.31. Def_L *is a deformation funcotr, i.e., it satisfies the conditions of Definition I.1.10.*

Proof. If $A = \mathbb{K}$, then it is clear that $\mathrm{Def}_L(B \times C) = \mathrm{Def}_L(B) \times \mathrm{Def}_L(C)$ and so condition *ii*) of Definition I.1.10 is satisfied.

Next, let $\beta : B \longrightarrow A$ and $\gamma : C \longrightarrow A$ be morphisms in **Art** with β surjective. Let $(l, m) \in \mathrm{Def}_L(B) \times_{\mathrm{Def}_L(A)} \mathrm{Def}_L(C)$ and $\tilde{l} \in MC_L(B)$ and $\tilde{m} \in \mathrm{MC}_L(C)$ be liftings of l and m, respectively, such that $\beta(\tilde{l}) = \gamma(\tilde{m}) \in \mathrm{Def}_L(A)$. Therefore, there exists $a \in L^0 \otimes m_A$ such that $e^a * \beta(\tilde{l}) = \gamma(\tilde{m})$. Let $b \in L^0 \otimes m_B$ be a lifting of a. By replacing $\tilde{l}$ with its gauge equivalent element $l' = e^b * \tilde{l}$ we can suppose $\beta(l') = \gamma(\tilde{m})$ in $\mathrm{MC}_L(A)$. By Remark I.3.22, MC_L is homogeneous and so there exists $n \in \mathrm{MC}_L(B \times_A C)$ that lifts $(l', \tilde{m})$. This implies that

$$\mathrm{Def}_L(B \times_A C) \longrightarrow \mathrm{Def}_L(B) \times_{\mathrm{Def}_L(A)} \mathrm{Def}_L(C)$$

is surjective. Hence condition *i*) of Definition I.1.10 also holds.

□

Remark I.3.32. By definition, the *tangent space* of Def_L is:

$$t_{\mathrm{Def}_L} := \mathrm{Def}_L(\mathbb{K}[\varepsilon]) = \frac{\{x \in L^1 \otimes \mathbb{K}\varepsilon \mid dx = 0\}}{\{da \mid a \in L^0 \otimes \mathbb{K}\varepsilon\}} \cong$$

$$H^1(L).$$

In general, if $L \otimes m_A$ is abelian then $\mathrm{Def}_L(A) = H^1(L) \otimes m_A$.

Lemma I.3.33. *The projection* $\pi : \mathrm{MC}_L \longrightarrow \mathrm{Def}_L$ *is a smooth morphism of functors.*

Proof. Let $\beta : B \longrightarrow A$ be a surjection in **Art**; we prove that

$$\mathrm{MC}_L(B) \longrightarrow \mathrm{Def}_L(B) \times_{\mathrm{Def}_L(A)} \mathrm{MC}_L(A),$$

induced by

$$\begin{tikzcd}
\mathrm{MC}_L(B) \arrow[r, "\beta"] \arrow[d, "\pi"'] & \mathrm{MC}_L(A) \arrow[d, "\pi"] \\
\mathrm{Def}_L(B) \arrow[r, "\beta"'] & \mathrm{Def}_L(A),
\end{tikzcd}$$

is surjective.

Let $(b, a) \in \mathrm{Def}_L(B) \times_{\mathrm{Def}_L(A)} \mathrm{MC}_L(A)$ and $\tilde{b} \in \mathrm{MC}_L(B)$ be a lifting of b. Then $\beta(\tilde{b})$ and a have a common image in $\mathrm{Def}_L(A)$ and so $\beta(\tilde{b}) = e^t * a$, for some $t \in L^0 \otimes m_A$.

Let $s \in L^0 \otimes m_B$ be a lifting of t and define $b' = e^{-s} * \tilde{b} \in \mathrm{MC}_L(B)$. Then $\beta(b') = e^{-t} * \beta(\tilde{b}) = a$ and b' lifts b. □

Therefore, by Corollary I.1.35, π induces an isomorphism between universal obstruction theories.

In conclusion, Lemma I.3.24 implies that $H^2(L)$ *is a complete obstruction space of* Def_L.

Theorem I.3.34. *Let $\phi : L \longrightarrow M$ be a morphism of DGLAs and denote by*

$$H^i(\phi) : H^i(L) \longrightarrow H^i(M)$$

the induced maps in cohomology.

- i) *If $H^1(\phi)$ is surjective (resp. bijective) and $H^2(\phi)$ injective, then the morphism $\mathrm{Def}_L \longrightarrow \mathrm{Def}_M$ is smooth (resp. étale).*
- ii) *If in addition to i) $H^0(\phi)$ is surjective, then the morphism $\mathrm{Def}_L \longrightarrow \mathrm{Def}_M$ is an isomorphism.*

Proof. *i)* follows from Proposition I.1.31 (resp. Corollary I.1.32). For a proof of *ii)* see [**20**, Theorem 3.1] (it also follows from the inverse function Theorem III.2.14 on the extended case). □

Corollary I.3.35. *Let $L \longrightarrow M$ be a quasi-isomorphism of DGLAs. Then the induced morphism $\mathrm{Def}_L \longrightarrow \mathrm{Def}_M$ is an isomorphism.*

Corollary I.3.36. *If $H^0(L) = 0$, then Def_L is homogenous.*

Proof. Let N be the DGLA introduced in Example I.3.12. Then the natural inclusion $N \longrightarrow L$ gives isomorphisms $H^i(N) \longrightarrow H^i(L)$ for each $i \geq 1$. Since $H^0(L) = 0$, $H^0(N) \longrightarrow H^0(L)$ is surjective. Therefore, Theorem I.3.34 *ii)* implies that $\mathrm{Def}_N \longrightarrow \mathrm{Def}_L$ is an isomorphism, with $\mathrm{Def}_N \cong \mathrm{MC}_N$ homogeneous.

□

Remark I.3.37. Let $F : \mathbf{Art} \longrightarrow \mathbf{Set}$ be the functor of the infinitesimal deformations of some algebro-geometric object defined over $\mathbb{K}$.

Then the guiding principle of Kontsevich (see [**18**]) affirms the existence of a DGLA L such that $F \cong \mathrm{Def}_L$. In spite of Corollary I.3.35, it is clear that this DGLA is defined only up to quasi-isomorphism. In this case we say that L *controls* the deformation functor F.

In Section II.7 we will prove the existence of a DGLA that controls the infinitesimal deformations of X (Theorem II.7.3) and in Section IV.2 the existence of a DGLA that controls the infinitesimal deformations of a holomorphic map f (Theorem IV.2.6).

CHAPTER II

Deformation of complex manifolds

In the first part of this chapter we fix notation and recall some known facts about complex manifolds that will be useful in the sequel.

Therefore, any book on complex varieties is a good reference for this chapter (for example [**11**], [**23**], [**33**], etc.).

In particular, we wish to recall the Čech cohomology and Leray's theorem (Section II.3.1) and some properties of Kähler manifolds (Section II.2). We also study the maps f_* and f^* induced by a holomorphic map f (Section II.6).

Moreover, we give the fundamental definition of the *Kodaira-Spencer differential graded Lie algebra* KS_X associated with a compact complex manifold X (Definition II.4.1), of the *contraction map* $\boldsymbol{i}$ and of the *holomorphic Lie derivative* $\boldsymbol{l}$ (Section II.5).

In the second part (Section II.7) we prove that the functor Def_X of the infinitesimal deformations of a compact complex manifold X (see Definition I.2.4) is isomorphic to the deformation functor Def_{KS_X} associated with the Kodaira-Spencer algebra KS_X of X (Theorem II.7.3).

Theorem. *Let X be a complex compact manifold and KS_X its Kodaira-Spencer algebra. Then there exists an isomorphism of functors*

$$\mathrm{Def}_{KS_X} \longrightarrow \mathrm{Def}_X .$$

Therefore, in spite of Remark I.3.37 we can say that the differential graded Lie algebra of Kodaira-Spencer KS_X controls the infinitesimal deformations of a complex compact manifold X.

This theorem is well known and a proof based on the theorem of Newlander-Nirenberg can be found in [**4**], [**10**] or more recently in [**23**]. Here we are interested in a simpler proof that avoids the use of this theorem.

Note that in this chapter we will work over the complex numbers and so $\mathbb{K} = \mathbb{C}$.

We also assume that every variety X is smooth (complex), compact, and connected.

II.1. Differential forms

Let X be such a manifold of dimension n and $T_{X,\mathbb{C}} = T_X^{1,0} \oplus T_X^{0,1}$ its complex tangent bundle, with $T_X^{1,0} =: \Theta_X$ the holomorphic tangent bundle and $T_X^{0,1} = \overline{T_X^{1,0}}$.

This decomposition induces a dual decomposition on the sheaf of differentiable forms

$$\mathcal{A}_X^1 = \mathcal{A}_X^{1,0} \oplus \mathcal{A}_X^{0,1},$$

with $\mathcal{A}_X^{1,0}$ the sheaf of complex differentiable forms of type $(1,0)$. If $z_1, \dots, z_n$ are local holomorphic coordinates on X, then $\mathcal{A}_X^{1,0}$ is generated by the dz_i: each $\alpha \in \mathcal{A}_X^{1,0}$ has the form $\alpha = \sum_i \alpha_i dz_i$, with $\alpha_i \in \mathcal{A}_X^{0,0}$ for any i.
In general, a (p,q)-form α can be locally written as $\alpha = \sum_{K,J} \alpha_{K,J} dz_K \wedge d\overline{z}_J$ with $\alpha_{K,J} \in \mathcal{A}_X^{0,0}$, $K = (1 \leq k_1 < k_2 < \dots < k_p \leq n)$ a multi-index of length p and $J = (1 \leq j_1 < j_2 < \dots < j_q \leq n)$ a multi-index of length q, such that $dz_K = dz_{k_1} \wedge dz_{k_2} \wedge \dots \wedge dz_{k_p}$ and $d\overline{z}_J = d\overline{z}_{j_1} \wedge d\overline{z}_{j_2} \wedge \dots \wedge d\overline{z}_{j_q}$.

If $\alpha = f \in \mathcal{A}_X^{0,0}$, then

$$df = \sum_{h=1}^n \frac{\partial f}{\partial z_h} dz_h + \sum_{h=1}^n \frac{\partial f}{\partial \overline{z}_h} d\overline{z}_h = \partial f + \overline{\partial} f,$$

with $\partial f \in \mathcal{A}_X^{1,0}$ and $\overline{\partial} f \in \mathcal{A}_X^{0,1}$.

In general, for $\alpha = \sum_{K,J} \alpha_{K,J} dz_K \wedge d\overline{z}_J \in \mathcal{A}_X^{p,q}$, we have

$$d\alpha = \sum_{K,J} d\alpha_{K,J} \wedge dz_K \wedge d\overline{z}_J = \partial\alpha + \overline{\partial}\alpha,$$

with

$$\partial\alpha = \sum_{I,J} \partial\alpha_{I,J} \wedge dz_K \wedge d\overline{z}_J \in \mathcal{A}^{p+1,q}$$

and

$$\overline{\partial}\alpha = \sum_{I,J} \overline{\partial}\alpha_{I,J} \wedge dz_K \wedge d\overline{z}_J \in \mathcal{A}^{p,q+1}.$$

Obviously, since $d^2 = 0$ we have $\partial^{\,2} = \overline{\partial}^{\,2} = \partial\overline{\partial} + \overline{\partial}\partial = 0$.

Proposition II.1.1. *Let α be a form of type (p,q), with $q > 0$, such that $\overline{\partial}\alpha = 0$. Then there exists, locally on X, a form β of type $(p, q-1)$ such that $\overline{\partial}\beta = \alpha$.*

Proof. See [**33**, Proposition 2.31]. □

Definition II.1.2. $(\mathcal{A}_X^{*,*}, \wedge)$ is the sheaf of graded algebras of differential forms of X, i.e., if $\mathcal{A}_X^{(p,q)}$ is the sheaf of differentiable (p,q)-forms then

$$\mathcal{A}_X^{*,*} := \bigoplus_i \mathcal{A}_X^i, \qquad \text{with} \qquad \mathcal{A}_X^i = \bigoplus_{p+q=i} \mathcal{A}_X^{p,q}.$$

We use the notation $A_X^{p,q} = \Gamma(X, \mathcal{A}_X^{p,q})$ for the vector space of global sections of $\mathcal{A}_X^{p,q}$.

Definition II.1.3. $Der^*(\mathcal{A}_X^{*,*})$ is the sheaf of $\mathbb{C}$-linear derivations on $\mathcal{A}_X^{*,*}$; more precisely, if $Der^{a,b}(\mathcal{A}_X^{*,*})$ are the derivations of bi-degree (a,b) then

$$Der^*(\mathcal{A}_X^{*,*}) := \bigoplus_k \bigoplus_{a+b=k} Der^{a,b}(\mathcal{A}_X^{*,*}).$$

We note that ∂ and $\overline{\partial}$ are global sections of $Der^{1,0}(\mathcal{A}_X^{*,*})$ and $Der^{0,1}(\mathcal{A}_X^{*,*})$, respectively.

Remark II.1.4. $Der^*(\mathcal{A}_X^{*,*})$ is a *sheaf of differential graded Lie algebras* with bracket and differential given by the following formulas:

$$[f,g] := f \circ g - (-1)^{\deg(f)\,\deg(g)} g \circ f$$

and

$$d(f) := [\partial + \overline{\partial}, f] = \partial f + \overline{\partial} f - (-1)^{\deg(f)}(f\partial + f\overline{\partial}).$$

In particular, fixing $p = 0$, $(\mathcal{A}_X^{0,*}, \wedge)$ is a sheaf of graded algebras and $Der^*(\mathcal{A}_X^{0,*}, \mathcal{A}_X^{0,*}) := \bigoplus_p Der^p(\mathcal{A}_X^{0,*}, \mathcal{A}_X^{0,*})$ is a sheaf of DGLAs (in this case the differential reduces to $d(f) = [\overline{\partial}, f] = \overline{\partial} f - (-1)^{\deg(f)} f\overline{\partial}$).

II.2. Kähler manifolds

This section is devoted to the compact Kähler manifolds. For definitions and properties of Kähler manifolds see for example [**11**], [**23**] or [**33**].

We include this section just to prove an important application (Lemma II.2.2) of the $\partial\overline{\partial}$-Lemma (Lemma II.2.1) that will be fundamental in the obstruction calculus of the last chapter of this thesis (Theorem V.1.4).

Lemma II.2.1 ($\partial\overline{\partial}$-Lemma). *Let X be a compact Kälher manifold and consider the operators ∂ and $\overline{\partial}$ on A_X. Then*

$$\operatorname{Im} \overline{\partial}\partial = \ker \partial \cap \operatorname{Im} \overline{\partial} = \ker \overline{\partial} \cap \operatorname{Im} \partial.$$

Proof. See for example [**23**, Theorem 6.37] and [**33**, Proposition 6.17].

□

Let $f : X \longrightarrow Y$ be a holomorphic map of compact complex manifolds. Let $\Gamma \subset X \times Y$ be the graph of f and $p : X \times Y \longrightarrow X$ and $q : X \times Y \longrightarrow Y$ be the natural projections.

Lemma II.2.2. *If X and Y are compact Kähler, then the subcomplexes $Im(\partial) = \partial A_{X\times Y}$, ∂A_Γ, $\partial A_{X\times Y} \cap q^* A_Y$ and $\partial A_{X\times Y} \cap p^* A_X$ are acyclic.*

Proof. By hypothesis, $X \times Y$ is Kähler. Then applying the $\partial\overline{\partial}$-Lemma II.2.1 to $A_{X\times Y}$ we get

$$\ker(\overline{\partial}) \cap \operatorname{Im}(\partial) = \operatorname{Im}(\overline{\partial}\partial)$$

and so $H^*_{\overline{\partial}}(\partial(A_{X\times Y})) = 0$. $\Gamma \subset X \times Y$ is also Kähler and so the same conclusion holds for ∂A_Γ: ∂A_Γ is acyclic.

Analogously, since Y is Kähler ∂A_Y and $q^*\partial A_Y$ are acyclic. Therefore, to prove that $\partial A_{X\times Y} \cap q^* A_Y$ is acyclic it suffices to prove that $\partial A_{X\times Y} \cap q^* A_Y = q^*\partial A_Y$.

The inclusion $\supseteq$ is obvious. Let $p \in q^*A_Y \cap \partial A_{X\times Y}$, then $p = q^*\phi = \partial z$ with $\phi \in A_Y$ and $z \in A_{X\times Y}$. The map $q^* : H_\partial(A_Y) \longrightarrow H_\partial(A_{X\times Y})$ is injective. Therefore, $\partial p = q^*\partial\phi = \partial\partial z = 0$ and so ϕ is ∂-closed ($\phi \in H_\partial(A_Y)$); moreover, $q^*[\phi] = [\partial z] = 0$. Thus ϕ is ∂-exact, that is, $\phi = \partial t$ with $t \in A_Y$. This implies $p = q^*\partial t \in q^*\partial A_Y$.

The case $\partial A_{X\times Y} \cap p^* A_X$ can be proved in the same way. $\square$

Remark II.2.3. In the previous lemma the Kähler hypothesis on X and Y can be substituted by the validity of the $\partial\overline{\partial}$-lemma in A_X,A_Y, $A_{X\times Y}$ and A_Γ.

II.3. Holomorphic fiber bundle and Dolbeault's cohomology

Let E be a holomorphic fiber bundle on X. Then the $\overline{\partial}$ operator can be extended to the *Dolbeault* operator

$$\overline{\partial}_E : \mathcal{A}^{p,q}_X(E) \longrightarrow \mathcal{A}^{p,q+1}_X(E).$$

If $e_1, \ldots, e_n$ is a local frame for E then

$$\overline{\partial}_E(\sum_i \phi_i e_i) = \sum_i \overline{\partial}(\phi) e_i.$$

Since E is a holomorphic fiber bundle, this definition does not depend on the choice of the local frame. By definition, $\overline{\partial}_E$ satisfies the property $\overline{\partial}_E{}^2 = 0$.

Let $A^{p,q}_X(E) = \Gamma(X, \mathcal{A}^{p,q}_X(E))$ be the vector space of global sections of the sheaf $\mathcal{A}^{p,q}_X(E)$. Then we can consider, for each $p \geq 0$, the following complex:

$$0 \longrightarrow A^{p,0}_X(E) \xrightarrow{\overline{\partial}_E} A^{p,1}_X(E) \xrightarrow{\overline{\partial}_E} \cdots \xrightarrow{\overline{\partial}_E} A^{p,q}_X(E) \xrightarrow{\overline{\partial}_E} \cdots .$$

The cohomology of this complex is *Dolbeault's cohomology* $H^{p,*}_{\overline{\partial}_E}(X, E)$ of E. We note that, for $p = 0$, $\ker(\overline{\partial}_E : A^{0,0}_X(E) \longrightarrow A^{0,1}_X(E))$ coincides with the holomorphic sections of E and

$$H^q_{\overline{\partial}_E}(X, E) := H^{0,q}_{\overline{\partial}_E}(X, E) = \frac{\ker(\overline{\partial}_E : A^{0,q}_X(E) \longrightarrow A^{0,q+1}_X(E))}{\operatorname{Im}(\overline{\partial}_E : A^{0,q-1}_X(E) \longrightarrow A^{0,q}_X(E))}.$$

PROPOSITION II.3.1. *Let α be a differential form with coefficients in E of type $(0,q)$ with $q>0$. If $\overline{\partial}_E\alpha=0$, then there exists, locally on X, a differential form β of type $(0,q-1)$, with coefficients in E, such that $\overline{\partial}_E\beta=\alpha$.*

PROOF. See [**33**, Proposition 2.36]. □

II.3.1. Čech cohomology and Leray's theorem. We follow [**23**, Section 1.3].

Let E be a holomorphic bundle on the complex manifold X. Let $\mathcal{U}=\{U_i\}_{i\in I}$ be a locally finite open covering of X and denote $U_{i_0\cdots i_k}=U_{i_0}\cap\cdots\cap U_{i_k}$.

Define the Čech q-chains of E:

$$\check{C}^k(\mathcal{U},E)=\{f_{i_0\cdots i_k}\mid f_{i_0\cdots i_k}:U_{i_0\cdots i_k}\longrightarrow E \text{ is a holomorphic section}\}$$

and the Čech differential

$$\check{\delta}:\check{C}^k(\mathcal{U},E)\longrightarrow\check{C}^{k+1}(\mathcal{U},E),$$

$$(\check{\delta}f)_{i_0\cdots i_{k+1}}=\sum_{j=0}^{k+1}(-1)^j f_{i_0\cdots\widehat{i_j}\cdots i_{k+1}}.$$

A simple calculation shows that $\check{\delta}^2=0$ and so we can define the $\mathbb{C}$-vector space of Čech cohomology

$$\check{H}^k(\mathcal{U},E)=\frac{\ker(\check{\delta}:\check{C}^k(\mathcal{U},E)\longrightarrow\check{C}^{k+1}(\mathcal{U},E)}{\operatorname{Im}(\check{\delta}:\check{C}^{k-1}(\mathcal{U},E)\longrightarrow\check{C}^{k}(\mathcal{U},E)}.$$

Next, define a morphism $\theta:\check{H}^k(\mathcal{U},E)\longrightarrow H^{0,k}_{\overline{\partial}_E}(X,E)$.

Let $t_i:X\longrightarrow\mathbb{C}$, with $i\in I$, be a partition of unity subordinate to the cover $\mathcal{U}$, that is, *supp*$(t_i)\subset U_i$, $\sum_i t_i=1$ and $\sum_i\overline{\partial}t_i=0$.

For each $f\in\check{C}^k(\mathcal{U},E)$ and $i\in I$ we define

$$\phi_i(f)=\sum_{j_1\cdots j_k}f_{ij_1\cdots j_k}\overline{\partial}t_{j_1}\wedge\cdots\wedge\overline{\partial}t_{j_k}\in\Gamma(U_i,\mathcal{A}^{0,k}(E))$$

and then

$$\phi(f)=\sum_i t_i\phi_i(f)\in\Gamma(X,\mathcal{A}^{0,k}(E)).$$

It can be proved that ϕ is a well defined morphism of complexes that induces a morphism θ in cohomology (for details see [**23**, Proposition 1.22]).

THEOREM II.3.2. *Let $\mathcal{U}=\{U_i\}_{i\in I}$ be a locally finite countable open covering of a complex manifold X and E a holomorphic vector bundle. If $H^{k-q}_{\overline{\partial}_E}(U_{i_0\cdots i_q},E)=0$ for every $q<k$ and $i_0\cdots i_k$, then the morphism θ is an isomorphism*

$$\theta:\check{H}^k(\mathcal{U},E)\longrightarrow H^k_{\overline{\partial}_E}(X,E).$$

PROOF. See [**23**, Theorem 1.24] or [**33**, Theorem 4.41]. $\square$

REMARK II.3.3. If the open sets U_i of the cover $\mathcal{U}$ are biholomorphic to open convex subsets of $\mathbb{C}^n$ then $\mathcal{U}$ satisfies the hypothesis of Theorem II.3.2.

REMARK II.3.4. It is convenient to give an explicit description of the inverse map of θ, at least for $k = 2$:

$$\theta^{-1} : H^2_{\overline{\partial}_E}(X, E) \longrightarrow \check{H}^2(\mathcal{U}, E).$$

Let $h \in H^2_{\overline{\partial}_E}(X, E)$. By applying Proposition II.3.1 for each $i \in I$, there exists $\tau_i \in \Gamma(U_i, \mathcal{A}^{0,1}(E))$ such that $h_{|U_i} = \overline{\partial}\tau_i$.

Define $\sigma_{ij} = (\tau_i - \tau_j)_{|U_{ij}} \in \Gamma(U_{ij}, \mathcal{A}^{0,1}(E))$. σ_{ij} is $\overline{\partial}$-closed; actually,

$$\overline{\partial}\sigma_{ij} = (\overline{\partial}\tau_i - \overline{\partial}\tau_j)_{|U_{ij}} = h_{|U_{ij}} - h_{|U_{ji}} = 0.$$

Therefore, for each U_{ij} there exists $\rho_{ij} \in \Gamma(U_{ij}, \mathcal{A}^{0,0}(E))$, such that $\overline{\partial}\rho_{ij} = \sigma_{ij}$.

We observe that $(\sigma_{jk} - \sigma_{ik} + \sigma_{ij})_{|U_{ijk}} = 0$; indeed,

$$(\sigma_{jk} - \sigma_{ik} + \sigma_{ij})_{|U_{ijk}} =$$

$$((\tau_j - \tau_k) - (\tau_i - \tau_k) + (\tau_i - \tau_j))_{|U_{ijk}} = 0.$$

Define $\alpha_{ijk} = (\rho_{jk} - \rho_{ik} + \rho_{ij})_{|U_{ijk}} \in \Gamma(U_{ijk}, \mathcal{A}^{0,0}(E))$. First of all, we have that $\overline{\partial}\alpha_{ijk} = 0$; actually,

$$\overline{\partial}\alpha_{ijk} = (\overline{\partial}\rho_{jk} - \overline{\partial}\rho_{ik} + \overline{\partial}\rho_{ij})_{|U_{ijk}} = (\sigma_{jk} - \sigma_{ik} + \sigma_{ij})_{|U_{ijk}} = 0$$

This implies that $\alpha_{ijk} \in \Gamma(U_{ijk}, E)$.

Moreover, $(\check{\delta}\alpha)_{ijkl} = 0$; indeed,

$$(\check{\delta}\alpha)_{ijkl} = (\alpha_{jkl} - \alpha_{ikl} + \alpha_{ijl} - \alpha_{ijk})_{|U_{ijkl}} =$$

$$((\rho_{kl} - \rho_{jl} + \rho_{jk}) - (\rho_{kl} - \rho_{il} + \rho_{ik}) + (\rho_{jl} - \rho_{il} + \rho_{ij}) - (\rho_{jk} - \rho_{ik} + \rho_{ij}))_{|U_{ijkl}} = 0.$$

This implies that $\alpha \in \check{H}^2(X, E)$.

α is independent of the choices of the lifitngs τ_i. Actually, if we choose $\overline{\tau}_i$, such that $h_{|U_i} = \overline{\partial}\overline{\tau}_i$, then $\overline{\tau}_i = \tau_i + \overline{\partial}t_i$ and this change does not affect the choice of α_{ijk}.

If we choose $\overline{\rho}_{ij} \in \Gamma(U_{ij}, \mathcal{A}^{0,0}(E))$ such that $\overline{\partial}\overline{\rho}_{ij} = \sigma_{ij}$, then $\overline{\rho}_{ij} = \rho_{ij} + s_{ij}$, with $s_{ij} \in \Gamma(U_{ij}, \mathcal{A}^{0,0}(E))$ such that $\overline{\partial}s_{ij} = 0$. This implies that $s_{ij} \in \Gamma(U_{ij}, E)$. Therefore, $\{\overline{\alpha}_{ijk}\} = \{\alpha_{ijk}\} + \{\check{\delta}(s_{ij})\}$ and so $\overline{\alpha}_{ijk}$ and α_{ijk} represent the same class in cohomology.

In conclusion, we have defined a map

$$\vartheta : H^2_{\overline{\partial}_E}(X, E) \longrightarrow \check{H}^2(\mathcal{U}, E),$$

$$[h] \longmapsto [\alpha].$$

Finally, it can be proved that this map ϑ is the inverse of θ (for details see [**23**, Theorem 1.24] or [**33**, Theorem 4.41]).

II.4. The Kodaira-Spencer algebra KS_X

Definition II.4.1. Let Θ_X be the holomorphic tangent bundle of a complex manifold X. The *Kodaira-Spencer* (differential graded Lie) algebra of X is

$$KS_X = \bigoplus_i \Gamma(X, \mathcal{A}_X^{0,i}(\Theta_X)) = \bigoplus_i A_X^{0,i}(\Theta_X).$$

In particular, KS_X^i is the vector space of the global sections of the sheaf of germs of the differential $(0,i)$-forms with coefficients in Θ_X.

The differential $\tilde{d}$ on KS_X is the opposite of Dolbeault's differential, whereas the bracket is defined in local coordinates as the $\overline{\Omega}^*$-bilinear extension of the standard bracket on $\mathcal{A}_X^{0,0}(\Theta_X)$ ($\overline{\Omega}^* = \ker(\partial : \mathcal{A}_X^{0,*} \longrightarrow \mathcal{A}_X^{1,*})$ is the sheaf of antiholomorphic differential forms).

Explicitly, if $z_1, \ldots, z_n$ are local holomorphic coordinates on X, we have

$$\tilde{d}(f d\overline{z}_I \frac{\partial}{\partial z_i}) = -\overline{\partial}(f) \wedge d\overline{z}_I \frac{\partial}{\partial z_i}.$$

$$[f \frac{\partial}{\partial z_i}\, d\overline{z}_I, g \frac{\partial}{\partial z_j}\, d\overline{z}_J] = (f \frac{\partial g}{\partial z_i} \frac{\partial}{\partial z_j} - g \frac{\partial f}{\partial z_j} \frac{\partial}{\partial z_i})\, d\overline{z}_I \wedge d\overline{z}_J, \qquad \forall\, f, g \in \mathcal{A}_X^{0,0},$$

($\mathcal{A}_X^{0,*}(T_X)$ is a sheaf of DGLAs).
We note that by Dolbeault theorem we have $H^i(A_X^{0,*}(\Theta_X)) \cong H^i(X, \Theta_X)$ for all i, then

$$H^q(KS_X) = \frac{\ker(\overline{\partial} : A_X^{0,q}(\Theta_X) \longrightarrow A_X^{0,q+1}(\Theta_X))}{\operatorname{Im}(\overline{\partial} : A_X^{0,q-1}(\Theta_X) \longrightarrow A_X^{0,q}(\Theta_X))} \cong H^q_{\overline{\partial}}(X, \Theta_X).$$

In Theorem II.7.3, we will prove that the DGLA KS_X controls the infinitesimal deformations of X.

II.5. Contraction map and holomorphic Lie derivative

In general, for any vector space V and linear functional $\alpha : V \longrightarrow \mathbb{C}$, we can define the *contraction operator*

$$\alpha \lrcorner : \bigwedge^k V \longrightarrow \bigwedge^{k-1} V,$$

$$\alpha \lrcorner (v_1 \wedge \ldots \wedge v_k) = \sum_{i=1}^{k} (-1)^{i-1} \alpha(v_i)(v_1 \wedge \ldots \wedge \hat{v}_i \wedge \ldots \wedge v_k),$$

that is a derivation of degree -1 of the graded algebra $(\bigwedge^k V, \wedge)$.

Then considering the contraction $\lrcorner$ of the differential forms with vector fields we can define two injective morphisms of sheaves:

- the *contraction map*

$$\boldsymbol{i} : \mathcal{A}_X^{0,*}(\Theta_X) \longrightarrow Der^*(\mathcal{A}_X^{*,*})[-1],$$
$$a \longmapsto \boldsymbol{i}_a \quad \text{with} \quad \boldsymbol{i}_a(\omega) = a\lrcorner\omega$$

- the *holomorphic Lie derivative*

$$\boldsymbol{l} : \mathcal{A}_X^{0,*}(\Theta_X) \longrightarrow Der^*(\mathcal{A}_X^{*,*}),$$
$$a \longmapsto \boldsymbol{l}_a = [\partial, \boldsymbol{i}_a] \quad \text{with} \quad \boldsymbol{l}_a(\omega) = \partial(a\lrcorner\omega) + (-1)^{\deg(a)} a\lrcorner\partial\omega$$

for each $a \in \mathcal{A}_X^{0,*}(\Theta_X)$ and $\omega \in \mathcal{A}_X^{*,*}$.

Lemma II.5.1. *With the notation above, for every* $a, b \in \mathcal{A}_X^{0,*}(\Theta_X)$ *we have*

$$\boldsymbol{i}_{\tilde{d}a} = -[\overline{\partial}, \boldsymbol{i}_a], \qquad \boldsymbol{i}_{[a,b]} = [\boldsymbol{i}_a, [\partial, \boldsymbol{i}_b]] = [[\boldsymbol{i}_a, \partial], \boldsymbol{i}_b], \qquad [\boldsymbol{i}_a, \boldsymbol{i}_b] = 0.$$

Proof. See [**22**, Lemma 2.1]. Let $z_1, z_2, \dots, z_n$ be local holomorphic coordinates on X. By linearity, we can assume that $a = f d\overline{z}_I \dfrac{\partial}{\partial z_i}$ and $b = g d\overline{z}_J \dfrac{\partial}{\partial z_j}$ $(i \neq j)$, with $f, g \in \mathcal{A}_X^{0,0}$.

All the expressions vanish on $\mathcal{A}_X^{0,*}$ and $\mathcal{A}_X^{*,*}$ is generated as $\mathbb{C}$-algebra by $\mathcal{A}_X^{0,0} \oplus \mathcal{A}_X^{0,1} \oplus \mathcal{A}_X^{1,0}$. Therefore, it is sufficient to verify the equalities on the dz_h (that generate $\mathcal{A}_X^{1,0}$).

Moreover, we note that $\overline{\partial} dz_h = \partial dz_h = \boldsymbol{i}_a\boldsymbol{i}_b dz_h = \boldsymbol{i}_b\boldsymbol{i}_a dz_h = 0$. Therefore, $[\boldsymbol{i}_a, \boldsymbol{i}_b] = 0$ and the other equalities follow from the easy calculations below.

Let $\omega = dz_h$ and $\tilde{d}(a) = -\overline{\partial}(f) \wedge d\overline{z}_I \dfrac{\partial}{\partial z_i}$. Then

$$\boldsymbol{i}_{\tilde{d}a}(dz_h) = \tilde{d}a\lrcorner dz_h = \begin{cases} 0 & h \neq i, \\ -\overline{\partial}(f) \wedge d\overline{z}_I & h = i. \end{cases}$$

On the other hand,

$$-[\overline{\partial}, \boldsymbol{i}_a](\omega) = (-\overline{\partial}\boldsymbol{i}_a + (-1)^{\overline{a}-1}\boldsymbol{i}_a\overline{\partial})(dz_h) = -\overline{\partial}\boldsymbol{i}_a(dz_h) =$$
$$-\overline{\partial}(a\lrcorner dz_h) = \begin{cases} 0 & h \neq i, \\ -\overline{\partial}(f d\overline{z}_I) & h = i. \end{cases}$$

Then the first equality holds.

As to $\boldsymbol{i}_{[a,b]}$, we have

$$[a, b] = (f \frac{\partial g}{\partial z_i}\frac{\partial}{\partial z_j} - g\frac{\partial f}{\partial z_j}\frac{\partial}{\partial z_i})\, d\overline{z}_I \wedge d\overline{z}_J$$

and then

$$\boldsymbol{i}_{[a,b]}(dz_h) = \begin{cases} 0 & h \neq i, j, \\ f\dfrac{\partial g}{\partial z_i}\, d\overline{z}_I \wedge d\overline{z}_J & h = j, \\ -\, g\dfrac{\partial f}{\partial z_j}\, d\overline{z}_I \wedge d\overline{z}_J & h = i. \end{cases}$$

On the other hand,

$$[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]]=[\boldsymbol{i}_a,\partial\boldsymbol{i}_b-(-1)^{\overline{b}-1}\boldsymbol{i}_b\partial]=$$

$$\boldsymbol{i}_a\partial\boldsymbol{i}_b-(-1)^{\overline{b}-1}\boldsymbol{i}_a\boldsymbol{i}_b\partial-(-1)^{(\overline{a}-1)\overline{b}}(\partial\boldsymbol{i}_b\boldsymbol{i}_a-(-1)^{\overline{b}-1}\boldsymbol{i}_b\partial\boldsymbol{i}_a).$$

Then

$$[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]](dz_h)=\boldsymbol{i}_a\partial\boldsymbol{i}_b(dz_h)-(-1)^{\overline{a}\overline{b}}\boldsymbol{i}_b\partial\boldsymbol{i}_a(dz_h)=$$

$$\begin{cases}0 & h\neq i,j,\\ \boldsymbol{i}_a\partial\boldsymbol{i}_b(dz_j)=f\dfrac{\partial g}{\partial z_i}d\overline{z}_I\wedge d\overline{z}_J & h=j,\\ -(-1)^{\overline{a}\overline{b}}\boldsymbol{i}_b\partial\boldsymbol{i}_a(dz_i)=-(-1)^{\overline{a}\overline{b}}g\dfrac{\partial f}{\partial z_j}d\overline{z}_J\wedge d\overline{z}_I & h=i.\end{cases}$$

□

The previous set of equalities is referred to as *Cartan formulas.*

Definition II.5.2. Let L and M be two differential graded Lie algebras and let d' be the differential on the graded vector space $\operatorname{Hom}^*(L,M)$. A linear map $i\in\operatorname{Hom}^{-1}(L,M)$ is called a *Cartan homotopy* if

$$i([a,b])=[i(a),d'i(b)]\qquad\text{and}\qquad[i(a),i(b)]=0\qquad\forall\,a,b\in L.$$

We recall that, by definition (see Example I.3.3), we have

$$d'i(a)=d_M(i(a))+i(d_L(a)).$$

Corollary II.5.3. *$\boldsymbol{i}$ is a Cartan homotopy and the Lie derivative $\boldsymbol{l}$ is a morphism of sheaves of DGLAs.*

Proof. Using Cartan formulas we get $d'(\boldsymbol{i}_b)=[d,\boldsymbol{i}_b]+\boldsymbol{i}_{\tilde{d}b}=[\partial+\overline{\partial},\boldsymbol{i}_b]-[\overline{\partial},\boldsymbol{i}_b]=[\partial,\boldsymbol{i}_b]$. Then $\boldsymbol{i}_{[a,b]}=[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]]=[\boldsymbol{i}_a,d'(\boldsymbol{i}_b)]$. Moreover, by Lemma II.5.1, $[\boldsymbol{i}_a,\boldsymbol{i}_b]=0$ and so $\boldsymbol{i}$ is a Cartan homotopy.

As to $\boldsymbol{l}$, we have

$$\boldsymbol{l}_{\tilde{d}a}=[\partial,\boldsymbol{i}_{\tilde{d}a}]=-[\partial,[\overline{\partial},\boldsymbol{i}_a]].$$

Moreover,

$$-[\partial,[\overline{\partial},\boldsymbol{i}_a]]=-[\partial,\overline{\partial}\boldsymbol{i}_a-(-1)^{\deg(\boldsymbol{i}_a)}\boldsymbol{i}_a\overline{\partial}]=$$

$$-\partial\overline{\partial}\boldsymbol{i}_a+(-1)^{\deg(\boldsymbol{i}_a)}\partial\boldsymbol{i}_a\overline{\partial}-(-1)^{\deg(\boldsymbol{i}_a)}\overline{\partial}\boldsymbol{i}_a\partial+\boldsymbol{i}_a\overline{\partial}\partial.$$

Therefore,

$$[d,\boldsymbol{l}_a]=[\partial+\overline{\partial},\boldsymbol{l}_a]=[\partial,\boldsymbol{l}_a]+[\overline{\partial},\boldsymbol{l}_a]=$$

$$=[\partial,[\partial,\boldsymbol{i}_a]]+[\overline{\partial},[\partial,\boldsymbol{i}_a]]=-[\partial,[\overline{\partial},\boldsymbol{i}_a]]=\boldsymbol{l}_{\tilde{d}a}.$$

As to the equality $\boldsymbol{l}_{[a,b]}=[\boldsymbol{l}_a,\boldsymbol{l}_b]$, we can prove it as follows

$$\boldsymbol{l}_{[a,b]}=[\partial,\boldsymbol{i}_{[a,b]}]=[\partial,[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]]]=$$

$$\partial[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]]-(-1)^{\overline{a}-1+\overline{b}}[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]]\partial=$$

$$\partial(\boldsymbol{i}_a[\partial,\boldsymbol{i}_b]-(-1)^{(\overline{a}-1)\overline{b}}[\partial,\boldsymbol{i}_b]\boldsymbol{i}_a)-(-1)^{\overline{a}-1+\overline{b}}(\boldsymbol{i}_a[\partial,\boldsymbol{i}_b]-(-1)^{(\overline{a}-1)\overline{b}}[\partial,\boldsymbol{i}_b]\boldsymbol{i}_a)\partial=$$

$$\partial\boldsymbol{i}_a[\partial,\boldsymbol{i}_b]-(-1)^{\overline{a}\overline{b}}[\partial,\boldsymbol{i}_b]\partial\boldsymbol{i}_a-(-1)^{\overline{a}-1}\boldsymbol{i}_a\partial[\partial,\boldsymbol{i}_b]-(-1)^{\overline{a}+\overline{a}\overline{b}}[\partial,\boldsymbol{i}_b]\boldsymbol{i}_a\partial=$$

$$(\partial\boldsymbol{i}_a - (-1)^{\overline{a}-1}\boldsymbol{i}_a\partial)[\partial,\boldsymbol{i}_b] - (-1)^{\overline{a}\overline{b}}[\partial,\boldsymbol{i}_b](\partial\boldsymbol{i}_a + (-1)^{\overline{a}}\boldsymbol{i}_a\partial) =$$
$$[\partial,\boldsymbol{i}_a][\partial,\boldsymbol{i}_b] - (-1)^{\overline{a}\overline{b}}[\partial,\boldsymbol{i}_b][\partial,\boldsymbol{i}_a] = [[\partial,\boldsymbol{i}_a],[\partial,\boldsymbol{i}_b]] = [\boldsymbol{l}_a,\boldsymbol{l}_b].$$

□

In particular, we have an injective morphism of sheaves

$$\boldsymbol{l} : \mathcal{A}_X^{0,*}(\Theta_X) \longrightarrow Der^*(\mathcal{A}_X^{0,*},\mathcal{A}_X^{0,*})$$

$$a \longmapsto \boldsymbol{l}_a \ \text{ with } \ \boldsymbol{l}_a(\omega) = (-1)^{\deg(a)} a \lrcorner \partial\omega. \tag{4}$$

Explicitly, in local holomorphic coordinates $z_1, z_2, \dots, z_n$, if $a = g\ d\overline{z}_I \dfrac{\partial}{\partial z_i}$ and $\omega = f d\overline{z}_J$, then

$$\boldsymbol{l}_a(\omega) = (-1)^{\deg(a)} \frac{\partial f}{\partial z_i}\, g\, d\overline{z}_I \wedge d\overline{z}_J.$$

Using $\boldsymbol{l}$, for each $(A, m_A) \in \mathbf{Art}$, we can define the following morphism:

$$\boldsymbol{l} : \mathcal{A}_X^{0,*}(\Theta_X) \otimes A \longrightarrow Der^*(\mathcal{A}_X^{0,*} \otimes A, \mathcal{A}_X^{0,*} \otimes A).$$

In particular, for each solution of the Maurer-Cartan equation in KS_X we have the fundamental lemma below.

Lemma II.5.4. *$x \in \mathrm{MC}_{KS_X}(A)$ if and only if*

$$\overline{\partial} + \boldsymbol{l}_x : \mathcal{A}_X^{0,*} \otimes A \longrightarrow \mathcal{A}_X^{0,*+1} \otimes A$$

is a differential of degree 1 on $\mathcal{A}_X^{0,*} \otimes A$.

Proof. Since $\boldsymbol{l}$ is a morphism of DGLAs, we have

$$(\overline{\partial} + \boldsymbol{l}_x)^2 = \overline{\partial}\boldsymbol{l}_x + \boldsymbol{l}_x\overline{\partial} + \boldsymbol{l}_x^2 = [\overline{\partial}, \boldsymbol{l}_x] + \frac{1}{2}[\boldsymbol{l}_x, \boldsymbol{l}_x] = \boldsymbol{l}(\tilde{d}x + \frac{1}{2}[x,x]).$$

□

Moreover, using $\boldsymbol{l}$, we can also define, for any $(A, m_A) \in \mathbf{Art}$ and $a \in \mathcal{A}_X^{0,0}(\Theta_X) \otimes m_A$, an automorphism e^a of $\mathcal{A}_X^{0,*} \otimes A$:

$$e^a : \mathcal{A}_X^{0,*} \otimes A \longrightarrow \mathcal{A}_X^{0,*} \otimes A, \qquad e^a(f) = \sum_{n=0}^{\infty} \frac{\boldsymbol{l}_a^n}{n!}(f). \tag{5}$$

Lemma II.5.5. *For every local Artinian $\mathbb{C}$-algebra (A, m_A), $a \in \mathcal{A}_X^{0,0}(\Theta_X) \otimes m_A$ and $x \in \mathrm{MC}_{KS_X}(A)$ we have*

$$e^a \circ (\overline{\partial} + \boldsymbol{l}_x) \circ e^{-a} = \overline{\partial} + e^a * \boldsymbol{l}_x \ : \ \mathcal{A}_X^{0,0} \otimes A \longrightarrow \mathcal{A}_X^{0,1} \otimes A, \tag{6}$$

where $*$ is the gauge action (and $e^a * \boldsymbol{l}_x \in \mathcal{A}_X^{0,1}(\Theta_X) \otimes m_A$ acts on $\mathcal{A}_X^{0,0} \otimes A$ as defined in (4)). In particular,

$$\ker(\overline{\partial} + e^a * \boldsymbol{l}_x : \mathcal{A}_X^{0,0} \otimes A \longrightarrow \mathcal{A}_X^{0,1} \otimes A) = e^a(\ker(\overline{\partial} + \boldsymbol{l}_x : \mathcal{A}_X^{0,0} \otimes A \longrightarrow \mathcal{A}_X^{0,1} \otimes A)).$$

PROOF. It follows from the definition of gauge action. More precisely, since $e^a \circ e^b \circ e^{-a} = e^{[a,]}(b)$, we have

$$e^a \circ (\overline{\partial} + l_x) \circ e^{-a} = e^{[a,]'}(\overline{\partial} + l_x) =$$

$$\sum_{n=0}^{\infty} \frac{([a,]')^n}{n!}(\overline{\partial} + l_x) = \overline{\partial} + l_x + \sum_{n=1}^{\infty} \frac{([a,]')^n}{n!}(\overline{\partial} + l_x) =$$

$$= \overline{\partial} + l_x + \sum_{n=0}^{\infty} \frac{([a,]')^{n+1}}{(n+1)!}(\overline{\partial} + l_x) = \overline{\partial} + l_x + \sum_{n=0}^{\infty} \frac{([a,]')^n}{(n+1)!}([a, \overline{\partial}]' + [a, l_x]) =$$

$$\overline{\partial} + l_x + \sum_{n=0}^{\infty} \frac{([a,]')^n}{(n+1)!}([a, l_x] - \overline{\partial}a) = \overline{\partial} + e^a * l_x.$$

□

REMARK II.5.6. Let ϕ_i be an automorphism of the A-module $\mathcal{A}_X^{0,*} \otimes A$ whose specialization to the residue field $\mathbb{C}$ is the identity. Let $\phi = \sum_i \phi_i = id + \eta$ with $\eta \in \mathrm{Hom}^0(\mathcal{A}_X^{0,*}, \mathcal{A}_X^{0,*}) \otimes m_A$. Since we are in characteristic zero, we can take the logarithm so that $\phi = e^a$ with $a \in \mathrm{Hom}^0(\mathcal{A}_X^{0,*}, \mathcal{A}_X^{0,*}) \otimes m_A$

II.5.1. The DGLA of a submanifold. Let X be a complex manifold and $i : X \hookrightarrow Y$ be the inclusion of a submanifold X. Let $i^* : \mathcal{A}_X^{0,*} \longrightarrow \mathcal{A}_Y^{0,*}$ be the restriction morphism (of sheaves of DGLAs). Finally, denote by Θ_Y the holomorphic tangent bundle of Y and by $N_{X|Y}$ the normal bundle of X in Y. Define the sheaf $\mathcal{L}' = \oplus_i \mathcal{L}'^i$ such that

$$0 \longrightarrow \mathcal{L}' \longrightarrow \mathcal{A}_Y^{0,*}(\Theta_Y) \longrightarrow \mathcal{A}_X^{0,*}(N_{X|Y}) \longrightarrow 0.$$

Let $z_1, \ldots, z_n$ be holomorphic coordinates on Y such that $Y \supset X = \{z_{t+1} = \cdots = z_n = 0\}$. Then $\eta \in \mathcal{L}'^i$ if and only if $\eta = \sum_{j=1}^{n} \omega_j \frac{\partial}{\partial z_j}$, with $\omega_j \in \mathcal{A}_Y^{0,i}$ such that $\omega_j \in \ker i^*$ for $j \geq t$. In particular, $\mathcal{L}'^0$ is the sheaf of differentiable vector fields on Y that are tangent to X.

LEMMA II.5.7. *$\mathcal{L}'$ is a sheaf of differential graded Lie subalgebras of $\mathcal{A}_Y^{0,*}(\Theta_Y)$ such that $\boldsymbol{l}_a(\ker i^*) \subset \ker i^*$ if and only if $a \in \mathcal{L}' \subset \mathcal{A}_Y^{0,*}(\Theta_Y)$.*

PROOF. See [**24**, Section 5]. It is an easy calculation in local holomorphic coordinates.

□

Moreover, consider the automorphism e^a of $\mathcal{A}_X^{0,*} \otimes A$ defined in (5): if $a \in \mathcal{L}'^0 \otimes m_A$ then $e^a(\ker(i^*) \otimes A) = \ker(i^*) \otimes A$.

Let L' be the DGLA of the global sections of $\mathcal{L}'$:

$$0 \longrightarrow L' \longrightarrow A_Y^{0,*}(\Theta_Y) \xrightarrow{\pi'} A_X^{0,*}(N_{X|Y}) \longrightarrow 0.$$

In the literature, the notation $L' = A_Y^{0,*}(\Theta_Y(-log\ X))$ can also be found.

In Section V.4 we will prove that L' controls the embedded deformations of the inclusion $i : X \hookrightarrow Y$ (Corollary V.4.1).

II.6. Induced maps f_* and f^* by a holomorphic map f

This section is devoted to the study of the maps f_* and f^* induced by a holomorphic map f. In particular, we prove a property of these maps (Lemma II.6.1) that will be used in the last chapter (Section V.1.2).

Let $f : X \longrightarrow Y$ be a holomorphic map of compact complex manifolds.

Let $\mathcal{U} = \{U_i\}_{i\in I}$ and $\mathcal{V} = \{V_i\}_{i\in I}$ be finite Stein open covers of X and Y, respectively, such that $f(U_i) \subset V_i$ (U_i is allowed to be empty). Then f induces morphisms

$$f^* : \check{C}^p(\mathcal{V}, \Theta_Y) \longrightarrow \check{C}^p(\mathcal{U}, f^*\Theta_Y),$$

and

$$f_* : \check{C}^p(\mathcal{U}, \Theta_X) \longrightarrow \check{C}^p(\mathcal{U}, f^*\Theta_Y).$$

Explicitly, for each $i \in I$ and local holomorphic coordinate systems $z = (z_1, z_2, \dots, z_n)$ on U_i and $w = (w_1, w_2, \dots w_m)$ on V_i such that $f(z_1, z_2, \dots, z_n) = (f_1(z), \dots, f_m(z))$, we have

$$f^* : \Gamma(V_i, \Theta_Y) \longrightarrow \Gamma(U_i, f^*\Theta_Y),$$

$$f^*(\sum_j g_j(w)\frac{\partial}{\partial w_j}) = \sum_j g_j(f(z))\frac{\partial}{\partial w_j}$$

and

$$f_* : \Gamma(U_i, \Theta_X) \longrightarrow \Gamma(U_i, f^*\Theta_Y),$$

$$f_*(\sum_k h_k(z)\frac{\partial}{\partial z_k}) = \sum_{k,j} h_k(z)\frac{\partial f_j(z)}{\partial z_k}\frac{\partial}{\partial w_j}.$$

Moreover, f_* and f^* commute with the Čech differential and they do not depend on the choice of the cover. Therefore, we get linear maps in cohomology :

$$f^* : \check{H}^p(Y, \Theta_Y) \longrightarrow \check{H}^p(X, f^*\Theta_Y)$$

and

$$f_* : \check{H}^p(X, \Theta_X) \longrightarrow \check{H}^p(X, f^*\Theta_Y).$$

Analogously, f induces morphisms

$$f^* : A^{p,q}_Y(\Theta_Y) \longrightarrow A^{p,q}_X(f^*\Theta_Y)$$

and

$$f_* : A^{p,q}_X(\Theta_X) \longrightarrow A^{p,q}_X(f^*\Theta_Y).$$

Let $\mathcal{U}$ and $\mathcal{V}$ be Stein covers and $z = (z_1, z_2, \dots, z_n)$ on U_i and $w = (w_1, w_2, \dots w_m)$ on V_i local holomorphic coordinate systems as above.

Let $K = (1 \leq k_1 < k_2 < \cdots < k_p \leq n)$ be a multi-index of length p and $J = (1 \leq j_1 < j_2 < \cdots < j_q \leq n)$ a multi-index of length q. Then

$$f_* : A^{p,q}(U_i, \Theta_X) \longrightarrow A_X^{p,q}(U_i, f^*\Theta_Y)$$

$$f_*\left(h(z)dz_K \wedge d\overline{z}_J \frac{\partial}{\partial z_i}\right) = h(z)dz_K \wedge d\overline{z}_J \sum_j \frac{\partial f_j(z)}{\partial z_i}\frac{\partial}{\partial w_j},$$

and

$$f^* : A^{p,q}(V_i, \Theta_Y) \longrightarrow A_X^{p,q}(U_i, f^*\Theta_Y),$$

$$f^*\left(g(w)dw_K \wedge d\overline{w}_J \frac{\partial}{\partial w_i}\right) = g(f(z))\partial f_K \wedge \overline{\partial} f_J \frac{\partial}{\partial w_i},$$

where

$$\partial f_K = \sum_{h=1}^{n} \frac{\partial f_{k_1}}{\partial z_h} dz_h \wedge \cdots \wedge \sum_{h=1}^{n} \frac{\partial f_{k_p}}{\partial z_h} dz_h$$

and

$$\overline{\partial} f_J = \sum_{h=1}^{n} \frac{\partial f_{j_1}}{\partial \overline{z}_h} d\overline{z}_h \wedge \cdots \wedge \sum_{h=1}^{n} \frac{\partial f_{j_q}}{\partial \overline{z}_h} d\overline{z}_h.$$

We note that f^* and f_* commute with ∂ and $\overline{\partial}$.

Moreover, for each k, there exist the following commutative diagrams

$$\begin{tikzcd}
\check{C}^q(\mathcal{V}, \Theta_Y) \arrow[r, "\phi"] \arrow[d, "f^*"] & A_Y^{0,q}(\Theta_Y) \arrow[d, "f^*"] \\
\check{C}^q(\mathcal{U}, f^*\Theta_Y) \arrow[r, "\phi"] & A_X^{0,q}(f^*\Theta_Y)
\end{tikzcd}$$

and

$$\begin{tikzcd}
\check{C}^q(\mathcal{U}, \Theta_X) \arrow[r, "\phi"] \arrow[d, "f_*"] & A_X^{0,q}(\Theta_X) \arrow[d, "f_*"] \\
\check{C}^q(\mathcal{U}, f^*\Theta_Y) \arrow[r, "\phi"] & A_X^{0,q}(f^*\Theta_Y),
\end{tikzcd}$$

where ϕ is the map defined in Section II.3.1. Therefore, $f_*\phi = \phi f_*$ and $f^*\phi = \phi f^*$.

Lemma II.6.1. *Let $f : X \longrightarrow Y$ be a holomorphic map of complex manifolds. Let $\chi \in \mathcal{A}_Y^{0,*}(\Theta_Y)$ and $\eta \in \mathcal{A}_X^{0,*}(\Theta_X)$ such that $f^*\chi = f_*\eta \in \mathcal{A}_X^{0,*}(f^*\Theta_Y)$. Then for any $\omega \in \mathcal{A}_Y^{*,*}$*

$$f^*(\chi \lrcorner \omega) = \eta \lrcorner f^*\omega.$$

Proof. Let $\mathcal{U} = \{U_i\}_{i\in I}$ and $\mathcal{V} = \{V_i\}_{i\in I}$ be finite open Stein covers of X and Y, respectively, as above. For each $i \in I$, let z be local holomorphic coordinate systems on U_i and w on V_i such that $f(z) = (f_1(z), \ldots, f_m(z))$.

Let

$$\mathcal{A}^{0,r}_X(\Theta_X) \ni \eta = \sum_{i=1}^n h_i(z)d\overline{z}_I\frac{\partial}{\partial z_i}$$

and

$$\mathcal{A}^{0,r}_Y(\Theta_Y) \ni \chi = \sum_{h=1}^m \varphi_h(w)d\overline{w}_H\frac{\partial}{\partial w_h},$$

with $I = (1 \leq i_1 < i_2 < \cdots < i_r \leq n)$ and $H = (1 \leq h_1 < h_2 < \cdots < h_r \leq n)$ multi-indexes of length r.

Therefore,

$$f_*\eta = \sum_{i=1}^n h_i(z)d\overline{z}_I\sum_{h=1}^m\frac{\partial f_h}{\partial z_i}\frac{\partial}{\partial w_h} = \sum_{h=1}^m\left(\sum_{i=1}^n h_i(z)\frac{\partial f_h}{\partial z_i}d\overline{z}_I\right)\frac{\partial}{\partial w_h}$$

and

$$f^*\chi = \sum_{h=1}^m \varphi_h(f(z))\overline{\partial}f_H\frac{\partial}{\partial w_h}.$$

By hypothesis, $f^*\chi = f_*\eta \in \mathcal{A}^{0,*}_X(f^*\Theta_Y)$, then

$$\varphi_h(f(z))\overline{\partial}f_H = \sum_{i=1}^n h_i(z)\frac{\partial f_h}{\partial z_i}d\overline{z}_I, \qquad \forall\ h = 1,\dots,m. \tag{7}$$

Next, let

$$\mathcal{A}^{p,q}(V_i) \ni \omega = g(w)\,dw_K \wedge d\overline{w}_J,$$

with $K = (1 \leq k_1 < k_2 < \cdots < k_p \leq n)$ a multi-index of length p and $J = (1 \leq j_1 < j_2 < \cdots < j_q \leq n)$ a multi-index of length q. Then

$$f^*\omega = g(f(z))\,\partial f_K \wedge \overline{\partial}f_J$$

and

$$\chi\lrcorner\omega = \sum_{h=1}^m \varphi_h(w)g(w)\,d\overline{w}_H \wedge \left(\frac{\partial}{\partial w_h}\lrcorner dw_K\right)\wedge d\overline{w}_J =$$

$$\sum_{h=1}^p(-1)^{h-1}\varphi_{k_h}(w)g(w)\,d\overline{w}_H \wedge dw_{K-\{k_h\}} \wedge d\overline{w}_J,$$

with $dw_{K-\{k_h\}} = dw_{k_1}\wedge\ldots\wedge\widehat{dw_{k_h}}\wedge\ldots\wedge dw_{k_p}$.

Therefore,

$$f^*(\chi\lrcorner\omega) = \sum_{h=1}^p(-1)^{h-1}\varphi_{k_h}(f(z))g(f(z))\,\overline{\partial}f_H \wedge \partial f_{K-\{k_h\}} \wedge \overline{\partial}f_J$$

and using (7) we get

$$f^*(\chi\lrcorner\omega) = \sum_{h=1}^p(-1)^{h-1}g(f(z))\left(\sum_{i=1}^n h_i(z)\frac{\partial f_{k_h}}{\partial z_i}d\overline{z}_I\right)\wedge \partial f_{K-\{k_h\}} \wedge \overline{\partial}f_J.$$

On the other hand,

$$\eta\lrcorner f^*\omega =$$

$$\sum_{i=1}^{n} h_i(z)g(f(z))\, d\overline{z}_I \left(\frac{\partial}{\partial z_i} \lrcorner \partial f_K\right) \wedge \overline{\partial} f_J =$$

$$\sum_{i=1}^{n} h_i(z)g(f(z))\, d\overline{z}_I \wedge \left(\sum_{h=1}^{p} (-1)^{h-1} \frac{\partial f_{k_h}}{\partial z_i} \partial f_{K-\{k_h\}}\right) \wedge \overline{\partial} f_J =$$

$$f^*(\chi \lrcorner \omega).$$

□

II.7. Deformations of complex manifolds

In this section we prove that the infinitesimal deformations of a compact complex manifold X are controlled by the differential graded Lie algebra of Kodaira-Spencer KS_X, i.e., $\mathrm{Def}_{KS_X} \cong \mathrm{Def}_X$.

We start with some lemmas and we postpone the proof to Section II.7.1, where we also give an explicit description of the isomorphism (see Theorem II.7.3).

Lemma II.7.1. *Let $A \in$ **Art** and $x \in MC_{KS_X}(A)$, then there exists a cover $\mathcal{U} = \{U_i\}$ of X, such that $x_{|U_i} \sim 0$ for each i.*

Proof. By Proposition II.3.1, there exists a cover $\mathcal{U} = \{U_i\}$ such that $H^1(U_i, \Theta_X) = 0$ for all i. Moreover, by Remark I.3.32, $H^1(X, \Theta_X)$ is the tangent space of the deformation functor Def_{KS_X}. Therefore, by Corollary I.1.21, Def_{KS_X} is locally trivial and so any $x \in MC_{KS_X}(A)$ is locally gauge equivalent to zero.

□

In Section II.4 we defined a morphism of sheaves

$$\boldsymbol{l} : \mathcal{A}_X^{0,*}(\Theta_X) \otimes A \longrightarrow Der^*(\mathcal{A}_X^{0,*} \otimes A, \mathcal{A}_X^{0,*} \otimes A),$$

$$a \longmapsto \boldsymbol{l}_a \quad \text{with} \quad \boldsymbol{l}_a(\omega) = (-1)^{\deg(a)} a \lrcorner \partial\omega.$$

Let $x \in MC_{KS_X}(A)$. Explicitly, in local holomorphic coordinates $z_1, z_2, \ldots, z_n$, if $x = \sum_{i,j} x_{ij} d\bar{z}_i \frac{\partial}{\partial z_j}$ and $\omega = f d\overline{z}_J$, then

$$\boldsymbol{l}_x(f) = -\sum_{i,j} x_{ij} \frac{\partial f}{\partial z_j} d\bar{z}_i \wedge d\overline{z}_J.$$

We also proved that for each $x \in MC_{KS_X}(A)$

$$\overline{\partial} + \boldsymbol{l}_x : \mathcal{A}_X^{0,*} \otimes A \longrightarrow \mathcal{A}_X^{0,*+1} \otimes A$$

is a differential (Lemma II.5.4).

Define $\mathcal{O}_A(x)$ as the kernel of $\overline{\partial} + \boldsymbol{l}_x : \mathcal{A}_X^{0,0} \otimes A \longrightarrow \mathcal{A}_X^{0,1} \otimes A$. Then we have the complex

$$0 \longrightarrow \mathcal{O}_A(x) \longrightarrow \mathcal{A}_X^{0,0} \otimes A \xrightarrow{\overline{\partial} + \boldsymbol{l}_x} \mathcal{A}_X^{0,1} \otimes A \xrightarrow{\overline{\partial} + \boldsymbol{l}_x} \cdots \xrightarrow{\overline{\partial} + \boldsymbol{l}_x} \mathcal{A}_X^{0,n} \otimes A \longrightarrow 0.$$

In Section II.4 we also defined, for each $s \in \mathcal{A}^{0,0}_X(\Theta_X) \otimes m_A$, an automorphism e^s of $\mathcal{A}^{0,*}_X \otimes A$.

Lemma II.7.2. *Let $F, G : \mathbf{Art} \longrightarrow \mathbf{Set}$ be the following functors*

$$F(A) := \{\textit{isomorphisms of complexes } e^s : (\mathcal{A}^{0,*}_X \otimes A, \overline{\partial} + \boldsymbol{l}_x) \longrightarrow (\mathcal{A}^{0,*}_X \otimes A, \overline{\partial} + \boldsymbol{l}_y)$$

$$\textit{with } s \in \mathcal{A}^{0,0}_X(\Theta_X) \otimes m_A \textit{ that specialize to identity}\}$$

$$G(A) := \{\textit{isomorphisms of sheaves of } A\textit{-module } \psi : \mathcal{O}_A(x) \longrightarrow \mathcal{O}_A(y)$$

$$\textit{that specialize to identity}\}.$$

Then the restriction morphism $\phi : F \longrightarrow G$ is surjective.

Proof. We proceed by induction on $d = dim_{\mathbb{C}} A$.
If $A = \mathbb{C}$, then $G(\mathbb{C}) = \{\text{identity}\}$ and so it can be lifted.
Assume that $d \geq 2$ and let

$$0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

be a small extension; by induction, each element in $G(A)$ can be lifted to $F(A)$.
Let ψ be an isomorphism between $\mathcal{O}_B(x)$ and $\mathcal{O}_B(y)$ ($\psi \in G(B)$); we want to lift it to an isomorphism e^s.

$\alpha(x)$ and $\alpha(y)$ are in $MC_{KS_X}(A)$ and ψ induces an isomorphism of sheaves of A-modules $\overline{\psi} : \mathcal{O}_A(\alpha(x)) \longrightarrow \mathcal{O}_A(\alpha(y))$. Therefore, by the induction hypothesis we can lift $\overline{\psi}$ to an isomorphism of complexes $e^{\overline{s}}$, i.e., $e^{\overline{s}^{-1}} \circ (\overline{\partial} + \boldsymbol{l}_{\alpha(x)}) \circ e^{\overline{s}} = \overline{\partial} + \boldsymbol{l}_{\alpha(y)}$ with $\overline{s} \in \mathcal{A}^{0,0}_X(\Theta_X) \otimes m_A$.

Then we can suppose that $\alpha(x) = \alpha(y) \in A^{0,1}(\Theta_X) \otimes m_A$ and that $e^{\overline{s}}$ is the identity.

This implies the existence of an element $p \in \mathcal{A}^{0,1}_X(\Theta_X) \otimes J$ such that $x = y + p$. Since x and y satisfy the Maurer-Cartan equation, $\overline{\partial} p = 0$. Indeed

$$0 = dx + \frac{1}{2}[x,x] = d(y+p) + \frac{1}{2}[y+p, y+p] = dy + dp + \frac{1}{2}[y,y] = dp.$$

Therefore, by Proposition II.3.1, there exists a Stein cover $\mathcal{U} = \{U_i\}_{i \in I}$ of X such that p is locally $\overline{\partial}$-exact, i.e., for each $i \in I$ there exists $t_i \in \mathcal{A}^{0,0}_X(\Theta_X) \otimes J$ such that $\overline{\partial} t_i = p_{|U_i}$. Then

$$y_{|U_i} = (x-p)_{|U_i} = x_{|U_i} - \overline{\partial} t_i = e^{t_i} * x_{|U_i},$$

where we use the fact that $J \cdot m_B = 0$ as in Example I.3.28.

In particular, by Lemma II.5.5, $e^{t_i} : (\mathcal{A}^{0,*}(U_i) \otimes B, \overline{\partial} + \boldsymbol{l}_x) \longrightarrow (\mathcal{A}^{0,*}(U_i) \otimes B, \overline{\partial} + \boldsymbol{l}_y)$ is an isomorphism of complexes, that lifts the isomorphism $e^{t_i} : \mathcal{O}_B(x)(U_i) \longrightarrow \mathcal{O}_B(y)(U_i)$. We note that e^{t_i} restricts to the identity on $\mathcal{O}_A(x)(U_i)$.

On the other hand, by Lemma II.7.1, the Maurer-Cartan element x is locally gauge equivalent to zero. Then for each $i \in I$ there exists $a_i \in \mathcal{A}^{0,0}(U_i, \Theta_X) \otimes m_B$ such that $e^{a_i} * x_{|U_i} = 0$. As before, Lemma II.5.5

implies that $e^{a_i} : (\mathcal{A}^{0,*}(U_i) \otimes B, \overline{\partial} + \boldsymbol{l}_x) \longrightarrow (\mathcal{A}^{0,*}(U_i) \otimes B, \overline{\partial})$ is an isomorphism of complexes, that lifts the isomorphism $e^{a_i} : \mathcal{O}_B(x)(U_i) \longrightarrow \mathcal{O}_X(U_i) \otimes B$.

Next, consider the isomorphism

$$\varphi_{|U_i} : \mathcal{O}_X(U_i) \otimes B \longrightarrow \mathcal{O}_X(U_i) \otimes B$$

defined as follows:

$$\varphi_{|U_i} = e^{a_i} \circ e^{-t_i} \circ \psi_{|U_i} \circ e^{-a_i} :$$

$$\mathcal{O}_X(U_i) \otimes B \xrightarrow{e^{-a_i}} \mathcal{O}_B(x)(U_i) \xrightarrow{\psi_{|U_i}} \mathcal{O}_B(y)(U_i) \xrightarrow{e^{-t_i}} \mathcal{O}_B(x)(U_i) \xrightarrow{e^{a_i}} \mathcal{O}_X(U_i) \otimes B.$$

Then, $\varphi_{|U_i}$ is an automorphism of $\mathcal{O}_X(U_i) \otimes B$ that restricts to the identity on $\mathcal{O}_X(U_i) \otimes A$. Therefore, Lemma I.2.6 implies the existence of $q_i \in \Gamma(U_i, \Theta_X) \otimes J$ such that $\varphi_{|U_i} = e^{q_i}$. In particular, $e^{a_i} \circ e^{-t_i} \circ \psi_{|U_i} \circ e^{-a_i} = e^{q_i}$; by Remark I.2.7, the automorphism e^{q_i} commutes with the other automorphisms and so

$$\psi_{|U_i} = e^{t_i + q_i}.$$

Let $s_i = t_i + q_i \in \mathcal{A}^{0,0}(U_i, \Theta_X) \otimes J$, then $e^{s_i} = \psi_{|U_i}$ and so we have locally lifted the isomorphism ψ.

Next, we prove that the automorphisms e^{s_i} can be glued together to obtain an automorphism e^s of $\mathcal{A}_X^{0,*} \otimes A$ that lifts ψ. Consider the intersection U_{ij}, then the isomorphisms coincide on $\mathcal{O}_B(x)(U_{ij})$, i.e., $e^{s_i}_{|U_{ij}} = \psi_{|U_{ij}} = e^{s_j}_{|U_{ij}} : \mathcal{O}_B(x)(U_{ij}) \longrightarrow \mathcal{O}_B(y)(U_{ij})$. Therefore, the isomorphism $e^{s_i - s_j}$ is the identity on $\mathcal{O}_B(x)(U_{ij})$. Since the action of $\mathcal{A}_X^{0,0}(\Theta_X) \otimes m_B$ on $A_X^{0,0}(U_{ij}) \otimes B$ is faithful on $\mathcal{O}_X(U_{ij}) \otimes B$, it follows that $(s_i - s_j)_{|U_{ij}} = 0$.

□

II.7.1. KS_X controls the infinitesimal deformations of X.

This section is devoted to prove that the Kodaira-Spencer algebra of a complex manifold X controls the infinitesimal deformations of X.

Theorem II.7.3. *Let X be a complex compact manifold and KS_X its Kodaira-Spencer algebra. Then there exists an isomorphism of functors*

$$\gamma' : \mathrm{Def}_{KS_X} \longrightarrow \mathrm{Def}_X,$$

defined in the following way: given a local Artinian $\mathbb{C}$-algebra (A, m_A) and a solution of the Maurer-Cartan equation $x \in A_X^{0,1}(\Theta_X) \otimes m_A$ we set

$$\mathcal{O}_A(x) = \ker(\mathcal{A}_X^{0,0} \otimes A \xrightarrow{\overline{\partial} + \boldsymbol{l}_x} \mathcal{A}_X^{0,1} \otimes A),$$

and the map $\mathcal{O}_A(x) \longrightarrow \mathcal{O}_X$ is induced by the projection $\mathcal{A}_X^{0,0} \otimes A \longrightarrow \mathcal{A}_X^{0,0} \otimes \mathbb{C} = \mathcal{A}_X^{0,0}$.

Remark II.7.4. As observed in Section I.2.1, a deformation of X can be interpreted as a morphism of sheaves of algebras $\mathcal{O}_A \longrightarrow \mathcal{O}_X$ such that $\mathcal{O}_A$ is flat over A and $\mathcal{O}_A \otimes_A \mathbb{C} \longrightarrow \mathcal{O}_X$ is an isomorphism.

Consequently, the first part of the following proof consists of showing the A-flatness of $\mathcal{O}_A(x)$ and the existence of an isomorphism $\mathcal{O}_A(x) \otimes_A \mathbb{C} \cong \mathcal{O}_X$.

Proof. For each $(A, m_A) \in \mathbf{Art}$ and $x \in MC_{KS_X}(A)$, we have defined

$$\mathcal{O}_A(x) = \ker(\mathcal{A}_X^{0,0} \otimes A \xrightarrow{\overline{\partial}+\boldsymbol{l}_x} \mathcal{A}_X^{0,1} \otimes A).$$

First of all, we observe that the projection π on the residue field $A/m_A \cong \mathbb{C}$ induces the following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_A(x) \arrow[r] \arrow[d, "\pi"] & \mathcal{A}_X^{0,0} \otimes A \arrow[r, "\overline{\partial}"] \arrow[d, "\pi"] & \cdots \arrow[r, "\overline{\partial}"] & \mathcal{A}_X^{0,n} \otimes A \arrow[r] \arrow[d, "\pi"] & 0 \\
 & \mathcal{O}_A(x) \otimes_A \mathbb{C} \arrow[r] \arrow[d] & \mathcal{A}_X^{0,0} \arrow[r, "\overline{\partial}"] \arrow[d, "id"] & \cdots \arrow[r, "\overline{\partial}"] & \mathcal{A}_X^{0,n} \arrow[r] \arrow[d, "id"] & 0 \\
0 \arrow[r] & \mathcal{O}_X \arrow[r] & \mathcal{A}_X^{0,0} \arrow[r, "\overline{\partial}"] & \cdots \arrow[r, "\overline{\partial}"] & \mathcal{A}_X^{0,n} \arrow[r] & 0.
\end{tikzcd}$$

Then π induces the morphism $\mathcal{O}_A(x) \longrightarrow \mathcal{O}_X$.

Using Lemma II.7.1, the Maurer-Cartan solution x is locally gauge equivalent to zero, therefore, there exist a cover $\mathcal{U} = \{U_i\}$ and elements $a_i \in A^{0,0}(U_i, \Theta_X) \otimes m_A$ such that $e^{a_i} * x_{|U_i} = 0$, for each i. Thus, by Lemma II.5.5, $e^{a_i} \circ (\overline{\partial} + \boldsymbol{l}_{x_{|U_i}}) \circ e^{-a_i} = e^{a_i} * (\overline{\partial} + \boldsymbol{l}_{x_{|U_i}}) = \overline{\partial} + e^{a_i} * \boldsymbol{l}_{x_{|U_i}} = \overline{\partial}$ and so we have the following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_A(x)(U_i) \arrow[r] \arrow[d, "e^{a_i}"] & \mathcal{A}_X^{0,0}(U_i) \otimes A \arrow[r, "\overline{\partial}+\boldsymbol{l}_{x_{|U_i}}"] \arrow[d, "e^{a_i}"] & \cdots \arrow[r, "\overline{\partial}+\boldsymbol{l}_{x_{|U_i}}"] & \mathcal{A}_X^{0,n}(U_i) \otimes A \arrow[r] \arrow[d, "e^{a_i}"] & 0 \\
0 \arrow[r] & \mathcal{O}_X(U_i) \otimes A \arrow[r] & \mathcal{A}_X^{0,0}(U_i) \otimes A \arrow[r, "\overline{\partial}"] & \cdots \arrow[r, "\overline{\partial}"] & \mathcal{A}_X^{0,n}(U_i) \otimes A \arrow[r] & 0,
\end{tikzcd}$$

where the vertical arrows are isomorphisms.

This implies that the deformation $\mathcal{O}_A(x)$ is locally trivial, i.e., $\mathcal{O}_A(x)(U_i) \cong \mathcal{O}_X(U_i) \otimes A$. Since $\mathcal{O}_X(U_i) \otimes A$ is flat over A, $\mathcal{O}_A(x)(U_i)$ is also A-flat. Since flatness is a local property, $\mathcal{O}_A(x)$ is A-flat.

Using the isomorphism $\mathcal{O}_A(x)(U_i) \cong \mathcal{O}_X(U_i) \otimes A$ we can also conclude that $\mathcal{O}_A(x)(U_i) \otimes_A \mathbb{C} \cong \mathcal{O}_X(U_i)$ and so $\mathcal{O}_A(x) \otimes_A \mathbb{C} \longrightarrow \mathcal{O}_X$ is an isomorphism.

Then it is well defined the following morphism of functors of Artin rings

$$\gamma : \mathrm{MC}_{KS_X} \longrightarrow \mathrm{Def}_X,$$

such that

$$\begin{aligned}\gamma(A) : \mathrm{MC}_{KS_X}(A) &\longrightarrow \mathrm{Def}_X(A)\\ x &\longmapsto \mathcal{O}_A(x).\end{aligned}$$

Next, we prove that the deformations $\mathcal{O}_A(x)$ and $\mathcal{O}_A(y)$ are isomorphic if and only if $x, y \in \mathrm{MC}_{KS_X}(A)$ are gauge equivalent.

Actually, if $\mathcal{O}_A(x) \cong \mathcal{O}_A(y)$, applying Proposition II.7.2, we can lift this isomorphism to an isomorphism of complexes e^s with $s \in A_X^{0,0} \otimes m_A$, i.e.,

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_A(x) \arrow[r] \arrow[d, "\cong"] & \mathcal{A}_X^{0,0} \otimes A \arrow[r, "\overline{\partial}+\boldsymbol{l}_x"] \arrow[d, "e^s"] & \cdots \arrow[r, "\overline{\partial}+\boldsymbol{l}_x"] & \mathcal{A}_X^{0,n} \otimes A \arrow[r] \arrow[d, "e^s"] & 0 \\
0 \arrow[r] & \mathcal{O}_A(y) \arrow[r] & \mathcal{A}_X^{0,0} \otimes A \arrow[r, "\overline{\partial}+\boldsymbol{l}_y"] & \cdots \arrow[r, "\overline{\partial}+\boldsymbol{l}_y"] & \mathcal{A}_X^{0,n} \otimes A \arrow[r] & 0.
\end{tikzcd}$$

The commutativity of the diagram and Lemma II.5.5 imply that $\overline{\partial} + \boldsymbol{l}_y = e^{-s} \circ (\overline{\partial} + \boldsymbol{l}_x) \circ e^s = \overline{\partial} + e^s * \boldsymbol{l}_x$. Therefore, $e^s * x = y$.

In conclusion, the map γ', induced by γ on $\mathrm{Def}_{KS_X} = MC_{KS_X}/gauge$, is a well defined injective morphism:

$$\gamma' : \mathrm{Def}_{KS_X} \hookrightarrow \mathrm{Def}_X\,.$$

To conclude that γ' is an isomorphism we prove that γ' is étale (and so γ' is surjective).

Using Corollary I.1.32, we need to prove that:
1) γ' induces a bijective map on the tangent spaces;
2) γ' induces an injective map on the obstruction spaces.

As to Def_{KS_X}, by Remark I.3.32, the tangent space is isomorphic to $H^1_{\overline{\partial}}(X, \Theta_X)$ and Lemma I.3.24 implies that the obstructions are naturally contained in $H^2_{\overline{\partial}}(X, \Theta_X)$. As to Def_X, Theorems I.2.8 and I.2.9 show that the tangent space is isomorphic to $\check{H}^1(X, \Theta_X)$ and the obstructions are naturally contained in $\check{H}^2(X, \Theta_X)$.

Then we will prove that the maps induced by γ' coincide with the Leray isomorphisms (see Theorem II.3.2 and Remark II.3.4).

1) Tangent Spaces. Let us prove that the map γ'_ε induced by γ' on the tangent space

$$\gamma'_\varepsilon : \mathrm{Def}_{KS_X}(\mathbb{C}[\varepsilon]) \longrightarrow \mathrm{Def}_X(\mathbb{C}[\varepsilon])$$

is the Leray isomorphism.

By Remark I.3.32, we have $\mathrm{Def}_{KS_X}(\mathbb{C}[\varepsilon]) = H^1(X, \Theta_X)$. Proceeding as in Remark II.3.4, there exists a Stein cover $\mathcal{U} = \{U_i\}$ so that we can associate with each $x \in \mathrm{Def}_{KS_X}(\mathbb{C}[\varepsilon])$ an element

$[\sigma] = [\{\sigma_{ij} = (a_i - a_j)_{|U_{ij}}\}] \in \check{H}^1(X, \Theta_X) \otimes \mathbb{C}\varepsilon$, with $x_{|U_i} = \overline{\partial} a_i$, that does not depend on the choice of a_i.

Next, let $\gamma(x) = \mathcal{O}_{\mathbb{C}[\varepsilon]}(x)$ be the deformation associated with x, i.e.,

$$0 \longrightarrow \mathcal{O}_{\mathbb{C}[\varepsilon]}(x) \longrightarrow \mathcal{A}_X^{0,0} \otimes \mathbb{C}[\varepsilon] \xrightarrow{\overline{\partial}+\boldsymbol{l}_x} \mathcal{A}_X^{0,1} \otimes \mathbb{C}[\varepsilon] \longrightarrow \cdots .$$

As before, the deformation $\mathcal{O}_{\mathbb{C}[\varepsilon]}(x)$ is locally trivial; thus, there exists $b_i \in \mathcal{A}^{0,0}(U_i, \Theta_X) \otimes \mathbb{C}\varepsilon$ such that $e^{b_i} * x_{|U_i} = 0$ and so

$$\mathcal{O}_{\mathbb{C}[\varepsilon]}(x)(U_i) \xrightarrow[\cong]{e^{b_i}} \mathcal{O}_X(U_i) \otimes \mathbb{C}[\varepsilon].$$

Proceeding as in the proof of Theorem I.2.8, for all i and j

$$\varphi_{ij} := e^{b_i - b_j} : \mathcal{O}_X(U_{ij}) \otimes \mathbb{C}[\varepsilon] \longrightarrow \mathcal{O}_X(U_{ij}) \otimes \mathbb{C}[\varepsilon]$$

is an automorphism of the trivial deformation $\mathcal{O}_X(U_{ij}) \otimes \mathbb{C}[\varepsilon]$ that restricts to the identity.

Applying Lemma I.2.6, the class $[\{\tau_{ij}\}] = [\{(b_i - b_j)_{|U_{ij}}\}] \in \check{H}^1(X, \Theta_X) \otimes \mathbb{C}\varepsilon$ is the Check 1-cocycle associated with the deformation $\gamma'_\varepsilon(x)$.

Since $e^{b_i} * x_{|U_i} = 0$, $\overline{\partial} b_i = x_{|U_i} = \overline{\partial} a_i$ and so $b_i = a_i + c_i$ with $c_i \in \Gamma(U_i, \Theta_X) \otimes \mathbb{C}\varepsilon$.

Therefore, $[\tau_{ij}] = [\{(b_i - b_j)_{|U_{ij}}\}] = [\sigma]$. This shows that γ'_ε coincides with the Leray's isomorphism.

2) Obstruction

Let

$$0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

be a small extension.

First we consider the obstruction class $[h]$ of $x \in \mathrm{Def}_{KS_X}$.

Let $x \in \mathrm{Def}_{KS_X}(A)$, and $\tilde{x} \in {KS_X}^1 \otimes m_B$ be a lifting of x. The obstruction class associated with x is $[h] \in H^2(KS_X) \otimes J$ with

$$h = \overline{\partial} \tilde{x} + \frac{1}{2}[\tilde{x}, \tilde{x}]$$

and this class does not depend on the choice of the lifting $\tilde{x}$ as it was shown in Lemma I.3.24.

Proceeding as in Remark II.3.4, there exists a Stein cover $\mathcal{U} = \{U_i\}$ such that the class $[\{\alpha_{ijk}\}] = [\{\rho_{jk} - \rho_{ik} + \rho_{ij}\}] \in \check{H}^2(X, \Theta_X) \otimes J$ is the class associated with h via the Leray's isomorphism, where $h_{|U_i} = \overline{\partial} \tau_i$ and $\overline{\partial} \rho_{ij} = (\tau_i - \tau_j)_{|U_{ij}}$. In particular, we note that $\rho_{ij} \in \Gamma(U_{ij}, \mathcal{A}^{0,0}(\Theta_X)) \otimes J$.

Since $h_{|U_i} = \overline{\partial} \tau_i$, x can be locally lifted to a solution of the Maurer-Cartan equation ,

$$\overline{x}_i = \tilde{x}_{|U_i} - \tau_i \in A^{0,1}(U_i, \Theta_X) \otimes m_B.$$

Indeed, on U_i we have

$$\alpha(\overline{x}) = \alpha(\tilde{x}) = x$$

and

$$\overline{\partial}\overline{x} + \frac{1}{2}[\overline{x},\overline{x}] = \overline{\partial}\tilde{x} - \overline{\partial}\tau_i + \frac{1}{2}[\tilde{x},\tilde{x}] = h_{|U_i} - \overline{\partial}\tau_i = 0.$$

Moreover, $e^{\rho_{ij}} * \overline{x}_{j|U_{ij}} = \overline{x}_{i|U_{ij}}$; more precisely (see Example I.3.28), we have

$$e^{\rho_{ij}} * \overline{x}_{j|U_{ij}} = e^{\rho_{ij}} * (\tilde{x}-\tau_j)_{|U_{ij}} = (-\overline{\partial}\rho_{ij} + \tilde{x} - \tau_j)_{|U_{ij}} = (\tilde{x}-\tau_i)_{|U_{ij}} = \overline{x}_{i|U_{ij}}.$$

As above, x is locally equivalent to zero; therefore, for each i there exists $a_i \in A^{0,0}(U_i, \Theta_X) \otimes m_A$, such that $e^{a_i} * x_{|U_i} = 0$.

Analogously, for each i, there exists $b_i \in A^{0,0}(U_i, \Theta_X) \otimes m_B$ that is a lifting of a_i such that

$$e^{b_i} * \overline{x}_i = 0.$$

Next, let $\gamma'(x) = \mathcal{O}_A(x)$ be the deformation of X induced by x. As above the deformation is locally trivial and so there exist a cover $\mathcal{U} = \{U_i\}$ and $a_i \in \mathcal{A}^{0,0}(U_i, T_X) \otimes m_A$ such that

$$\mathcal{O}_A(x)(U_i) \overset{e^{a_i}}{\cong} \mathcal{O}_X(U_i) \otimes A.$$

Let φ_{ij} be the following isomorphism

$$\varphi_{ij} : \mathcal{O}_X(U_{ij}) \otimes A \xrightarrow{e^{-a_j}} \mathcal{O}_A(x)(U_{ij}) \xrightarrow{e^{a_i}} \mathcal{O}_X(U_{ij}) \otimes A.$$

Proceeding as in the proof of Theorem I.2.9, since $b_i \in \mathcal{A}^{0,0}(U_i, \Theta_X) \otimes m_B$ are liftings of a_i, $\tilde{\varphi_{ij}} = e^{-b_j}e^{\rho_{ij}}e^{b_i} \in \operatorname{Aut}(\mathcal{O}_X(U_{ij}) \otimes B)$, defined as

$$\tilde{\varphi}_{ij} : \mathcal{O}_X(U_{ij}) \otimes B \xrightarrow{e^{-b_j}} \mathcal{O}_B(\overline{x}_j)(U_{ij}) \xrightarrow{e^{\rho_{ij}}} \mathcal{O}_B(\overline{x}_i)(U_{ij}) \xrightarrow{e^{b_i}} \mathcal{O}_X(U_{ij}) \otimes B,$$

is a lifting of φ_{ij}.

By Remark I.2.7, the automorphisms $e^{\rho_{ij}}$, $e^{\rho_{ik}}$ and $e^{\rho_{jk}}$ commute with the other automorphisms. Then $\Phi_{ijk} = \tilde{\varphi_{jk}}\tilde{\varphi_{ik}}^{-1}\tilde{\varphi_{ij}} = e^{\rho_{jk}-\rho_{ik}+\rho_{ij}}$ is an automorphism of the trivial deformation that restricts to the identity ($\Phi_{ijk|\mathcal{O}_X(U_{ijk})\otimes A} = id$). Therefore, by Lemma I.2.6, the element $[\{\alpha_{ijk}\}] = [\{\rho_{jk} - \rho_{ik} + \rho_{ij}\}] \in \check{H}^2(\mathcal{U}, \Theta_X) \otimes J$ is the obstruction class associated with $\gamma'(x)$. In conclusion, in the obstruction case too, the map induced by γ' coincides with the Leray's isomorphism.

□

II.7.2. Deformations of a product. As an application of Theorem II.7.3 we study deformations of a product of compact complex manifolds X and Y. The following remark will be used in Section IV.2.

REMARK II.7.5. In general, not all the deformations of the product $X \times Y$ are products of deformations of X and of Y.

The first example of this fact was given by Kodaira and Spencer in their work ([**17**, pag. 436]) where they provided one of the first examples of obstructed varieties. More precisely, they considered the product of the projective line and the complex torus of dimension $q \geq 2$

$\mathbb{P}^1 \times \mathbb{C}^q/G$ and proved that it is obstructed though the two manifolds are unobstructed.

A sufficient and necessary condition to have an isomorphism between products of deformations and deformations of the product is given by the lemma below.

LEMMA II.7.6. *The morphism*

$$F : \mathrm{Def}_X \times \mathrm{Def}_Y \longrightarrow \mathrm{Def}_{X\times Y}$$

is an isomorphism if and only if $H^1(\mathcal{O}_X) \otimes H^0(\Theta_Y) = H^1(\mathcal{O}_Y) \otimes H^0(\Theta_X) = 0$.

PROOF. By Theorem II.7.3, it suffices to prove that the morphism

$$F : \mathrm{Def}_{KS_X} \times \mathrm{Def}_{KS_Y} \longrightarrow \mathrm{Def}_{KS_{X\times Y}}$$

is an isomorphism.

Let p and q be the natural projections of the product $X \times Y$ onto X and Y, respectively. Consider the morphism of DGLAs

$$F : KS_X \times KS_Y \longrightarrow KS_{X\times Y},$$

$$(n_1, n_2) \longmapsto p^* n_1 + q^* n_2.$$

Thus, we have to prove that F induces an isomorphism of the associated deformation functors. Denote by $H^i(F)$ the induced map on cohomology.

By Theorem I.3.34, if $H^0(F)$ is surjective, $H^1(F)$ is bijective and $H^2(F)$ is injective, then F induces an isomorphism.

$H^0(KS_{X\times Y}) = H^0(X\times Y, \Theta_{X\times Y})$ and so, by the Kunneth's formula, $H^0(KS_{X\times Y}) = H^0(\Theta_X) \oplus H^0(\Theta_Y) = H^0(KS_X \times KS_Y)$. This implies that $H^0(F)$ is surjective.

Again by Kunneth's formula we get

$$H^1(KS_{X\times Y}) \cong H^1(\Theta_{X\times Y}) \cong$$

$$H^1(\Theta_X) \oplus H^1(\Theta_Y) \oplus H^1(\mathcal{O}_X) \otimes H^0(\Theta_Y) \oplus H^1(\mathcal{O}_Y) \otimes H^0(\Theta_X),$$

and

$$H^1(KS_X \times KS_Y) = H^1(\Theta_X) \oplus H^1(\Theta_Y).$$

Then the hypothesis implies that $H^1(F)$ is bijective.

Finally, reasoning as above we get that

$$H^2(F) : H^2(\Theta_X) \oplus H^2(\Theta_Y) \longrightarrow H^2(\Theta_{X\times Y})$$

is injective. This implies that F induce an isomorphism of deformation functors.

On the other hand, if F is an isomorphism of deformation functors, then $H^1(F)$ is a bijection onto the tangent spaces and so $H^1(\mathcal{O}_X) \otimes H^0(\Theta_Y) = H^1(\mathcal{O}_Y) \otimes H^0(\Theta_X) = 0$. □

CHAPTER III

Deformation functor of a pair of morphisms of DGLAs

In this chapter we give the key definition of deformation functor associated with a pair of morphisms of differential graded Lie algebras.

In the first section we define the (not extended) functors of Artin rings $\mathrm{MC}_{(h,g)}$ (Definition III.1.8) and $\mathrm{Def}_{(h,g)}$ (Definition III.1.12) associated with a pair $h : L \longrightarrow M$ and $g : N \longrightarrow M$. These functors will play an important role in the study of infinitesimal deformations of holomorphic maps in the next chapter.

Then Section III.2 is devoted introducing the extended deformation functors (Definition III.2.4). In particular, we define the functors $\widetilde{\mathrm{MC}}_{(h,g)}$ and $\widetilde{\mathrm{Def}}_{(h,g)}$ which are a generalization of the previous $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$.

We introduce the extended functors, since using their properties, we can show the existence of a DGLA $\mathrm{H}_{(h,g)}$ such that $\mathrm{Def}_{\mathrm{H}_{(h,g)}} \cong \mathrm{Def}_{(h,g)}$ (Theorem III.2.36).

III.1. Functors $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$

In this section we introduce the functors $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$ (Section III.1.2) associated with a pair $h : L \longrightarrow M$ and $g : N \longrightarrow M$ and we study some of their properties (Section III.1.3). First of all, we recall the definition of the mapping cone associated with morphisms of complexes.

Note. In this section, we suppose that M is concentrated in non negative degree. Since in the main application M will be the Kodaira-Spencer algebra of a manifold, this extra hypothesis is not restrictive. Anyway, in Section III.2 we will remove this hypothesis.

III.1.1. The mapping cone of a pair of morphisms.

The *suspension of the mapping cone of a morphism of complexes* $h : (L, d) \longrightarrow (M, d)$ is the differential graded vector space $(C_h^{\cdot}, \delta)$, where

$$C_h^i = L^i \oplus M^{i-1}$$

and the differential δ is

$$\delta(l, m) = (dl, -dm + h(l)) \in L^{i+1} \oplus M^i, \qquad \forall\, (l, m) \in L^i \oplus M^{i-1}.$$

Actually, $\delta^2(l,m)=\delta(dl,-dm+h(l))=(d^2l,d^2m-d(h(l))+h(d(l)))=(0,0)$.

Remark III.1.1. Consider the projection $\pi: M \longrightarrow \operatorname{coker}(h)$. Then there exists a morphism of complexes

$$\varphi:(C^{\cdot}_h,\delta)\longrightarrow(\operatorname{coker}(h)[-1],d_{[-1]}),$$

with

$$\varphi(l,m)=\pi(m), \qquad \forall\ (l,m)\in C^i_h.$$

Indeed,

$$\begin{tikzcd}
(l,m) \arrow[r, "\varphi"] \arrow[d, "\delta"'] & \pi(m) \arrow[d, "d_{[-1]}"] \\
(dl,-dm+h(l)) \arrow[r, "\varphi"] & -d\pi(m).
\end{tikzcd}$$

If h is injective then $(C^{\cdot}_h,\delta)$ and $(\operatorname{coker}(h)[-1],d_{[-1]})$ are quasi-isomorphic.

Remark III.1.2. Let $f: X\longrightarrow Y$ be a holomorphic map of complex compact manifolds and consider the induced map $f_*: A^{p,q}_X(\Theta_X)\longrightarrow A^{p,q}_X(f^*\Theta_Y)$ (see Section II.6). Then the cohomology groups $H^i(C^{\cdot}_{f_*})$ of the associated cone $(C^{\cdot}_{f_*},\delta)$ are isomorphic to the hyper-cohomology groups $\mathbb{H}^1\left(X,\mathcal{O}(\Theta_X)\xrightarrow{f_*}\mathcal{O}(f^*\Theta_Y)\right)$ (see [**11**, Section 5 of Chapter 3].

Next, suppose that $h:(L,d)\longrightarrow(M,d)$ and $g:(N,d)\longrightarrow(M,d)$ are morphisms of complexes:

$$\begin{tikzcd}
 & L \arrow[d, "h"] \\
N \arrow[r, "g"] & M.
\end{tikzcd}$$

Definition III.1.3. The *suspension of the mapping cone of a pair of morphisms* (h,g) is the differential graded vector space $(\mathrm{C}^{\cdot}_{(h,g)},D)$, where

$$\mathrm{C}^i_{(h,g)}=L^i\oplus N^i\oplus M^{i-1}$$

and the differential D is defined as follows

$$L^i\oplus N^i\oplus M^{i-1}\ni(l,n,m)\xrightarrow{D}(dl,dn,-dm-g(n)+h(l))\in L^{i+1}\oplus N^{i+1}\oplus M^i.$$

Actually, $D^2(l,n,m)=D(dl,dn,-dm-g(n)+h(l))=(d^2l,d^2n,d^2m+dg(n)-dh(l)-g(dn)+h(dl))=(0,0,0)$.

Remark III.1.4. By definition, $(\mathrm{C}^{\cdot}_{(h,g)},D)$ coincides with the suspended mapping cone associated with the morphism of complexes $h-g: L\oplus N\longrightarrow M$ (such that $(h-g)(l,n)=h(l)-g(n)$).

Moreover, the projection $\mathrm{C}^{\cdot}_{(h,g)}\longrightarrow L^{\cdot}\oplus N^{\cdot}$ is a morphism of complexes and so there exists the following exact sequence

$$\begin{tikzcd}0 \arrow[r] & (M^{\cdot -1}, -d) \arrow[r] & (C^{\cdot}_{(h,g)}, D) \arrow[r] & (L^{\cdot} \oplus N^{\cdot}, d) \arrow[r] & 0\end{tikzcd}$$

that induces

(8)

$$\cdots \longrightarrow H^i(\mathrm{C}^{\cdot}_{(h,g)}) \longrightarrow H^i(L^{\cdot} \oplus N^{\cdot}) \longrightarrow H^i(M^{\cdot}) \longrightarrow H^{i+1}(\mathrm{C}^{\cdot}_{(h,g)}) \longrightarrow \cdots .$$

Remark III.1.5. The complexes $\mathrm{C}^{\cdot}_{(h,g)}$ and $\mathrm{C}^{\cdot}_{(g,h)}$ are isomorphic. Actually, let $\gamma : \mathrm{C}^{\cdot}_{(h,g)} \longrightarrow \mathrm{C}^{\cdot}_{(g,h)}$ be defined as

$$\mathrm{C}^i_{(h,g)} \ni (l, n, m) \overset{\gamma}{\longmapsto} (-l, -n, m) \in \mathrm{C}^i_{(g,h)} .$$

Then

$$\begin{tikzcd}(l,n,m) \arrow[r, "\gamma"] \arrow[d, "D"'] & (-l,-n,m) \arrow[d, "\delta"] \\ (dl, dn, -dm - g(n) + h(l)) \arrow[r, "\gamma"] & (-dl, -dn, -dm - g(n) + h(l)),\end{tikzcd}$$

and so γ is a well defined morphism of complexes that is a quasi-isomorphism ($\gamma^2 = id$).

Lemma III.1.6. *Let $g : N \longrightarrow M$ and $h : L \longrightarrow M$ be morphisms of complexes with h injective, i.e., there exists the exact sequence of complexes*

$$\begin{tikzcd}0 \arrow[r] & L \arrow[r, "h"] & M \arrow[r, "\pi"] & \mathrm{coker}(h) \arrow[r] & 0 \\ & & N. \arrow[u, "g"'] & & \end{tikzcd}$$

Then $(\mathrm{C}^{\cdot}_{(h,g)}, D)$ is quasi-isomorphic to $(C^{\cdot}_{\pi\circ g}, \delta)$.

Proof. Let $\gamma : \mathrm{C}^{\cdot}_{(h,g)} \longrightarrow C^{\cdot}_{\pi\circ g}$ be defined as

$$\mathrm{C}^i_{(h,g)} \ni (l, n, m) \overset{\gamma}{\longmapsto} (-n, \pi(m)) \in C^i_{\pi\circ g},$$

then

$$\begin{tikzcd}(l,n,m) \arrow[r, "\gamma"] \arrow[d, "D"'] & (-n, \pi(m)) \arrow[d, "\delta"] \\ (dl, dn, -dm - g(n) + h(l)) \arrow[r, "\gamma"] & (-dn, -d\pi(m) - \pi(g(n))).\end{tikzcd}$$

Therefore, γ is a well defined morphism of complexes that is a quasi-isomorphism. The fact that γ induces isomorphisms in cohomology is an easy calculation but we state it for completeness.

Denote by the same γ the map induced in cohomology.

γ is injective. Suppose that $\gamma[(l, n, m)] = [(-n, \pi(m))]$ is zero in $H^i(C^{\cdot}_{\pi\circ g})$. Then $dl = dn = -dm - g(n) + l = 0$ and there exists $(b, c) \in N^{i-1} \times \mathrm{coker}(h)^{i-2}$ such that $-n = db$ and $\pi(m) = -dc + \pi\circ g(b)$. Let $m' \in M^{i-2}$ be a lifting of c, i.e., $\pi(m') = c$, $n' = -b \in N^{i-1}$ and $l' = m + dm' + g(n') \in M^{i-1}$. Then $l' \in L^{i-1}$, actually, $\pi(l') =$

$\pi(m)+\pi(dm')+\pi\circ g(n') = -dc+\pi\circ g(b)+dc-\pi\circ g(b) = 0$. Therefore, $D(l',n',m') = (dl',dn',-dm'-g(n')+l') = (dm-g(db),-db,-dm'+g(b)+(m+dm'+g(n'))) = (l,n,m)$.

γ is surjective. Let $[(n,t)] \in H^i(C^{\cdot}_{\pi\circ g})$; then $dn = 0$ and $-dt+\pi\circ g(n) = 0$. Let $m \in M^{i-1}$ be a lifting of t, i.e., $\pi(m) = t$ and $l = -g(n)+dm \in M^i$. Then $l \in L^i$; in fact, $\pi(l) = -\pi\circ g(n)+dt = 0$. Moreover, $(l,-n,m) \in H^i(\mathrm{C}^{\cdot}_{(h,g)})$ ($dl = dn = 0$ and $-dm+g(n)+l = 0$) and $\gamma[(l,-n,m)] = [n,\pi(m)] = [(n,t)]$.

□

Remark III.1.7. Let L, M and N be DGLAs and $h : L \longrightarrow M$ and $g : N \longrightarrow M$ morphisms of DGLAs. No canonical DGLA structure on $\mathrm{C}^{\cdot}_{(h,g)}$ can be defined such that the projection $\mathrm{C}^{\cdot}_{(h,g)} \longrightarrow L^{\cdot} \oplus N^{\cdot}$ is a morphism of DGLAs.

III.1.2. Definition of $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$.

Definition III.1.8. Let $h : L \longrightarrow M$ and $g : N \longrightarrow M$ be morphisms of differential graded Lie algebras:

$$\begin{tikzcd} & L \arrow[d, "h"] \\ N \arrow[r, "g"] & M. \end{tikzcd}$$

For each $(A, m_A) \in \mathbf{Art}$ the *Maurer-Cartan functor associated with the pair* (h, g) is defined as follows

$$\mathrm{MC}_{(h,g)} : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{MC}_{(h,g)}(A) = \{(x, y, e^p) \in (L^1 \otimes m_A) \times (N^1 \otimes m_A) \times \exp(M^0 \otimes m_A) | \\ dx + \frac{1}{2}[x, x] = 0,\ dy + \frac{1}{2}[y, y] = 0,\ g(y) = e^p * h(x)\}.$$

Remark III.1.9. In [**24**, Section 2], M. Manetti defined the functor MC_h associated with a morphism $h : L \longrightarrow M$ of DGLAs:

$$\mathrm{MC}_h : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{MC}_h(A) =$$

$$\{(x, e^p) \in (L^1 \otimes m_A) \times \exp(M^0 \otimes m_A) |\ dx + \frac{1}{2}[x, x] = 0,\ e^p * h(x) = 0\}.$$

Therefore, if we take $N = 0$ and $g = 0$, the new functor $\mathrm{MC}_{(h,g)}$ reduces to the old one MC_h.

By choosing $N = 0$ and $h = g = 0$, $\mathrm{MC}_{(h,g)}$ reduces to the Maurer-Cartan functor MC_L associated with the DGLA L (Definition I.3.17).

Remark III.1.10. As in the case of a differential graded Lie algebra (see Remark I.3.22), $\mathrm{MC}_{(h,g)}$ is a *homogeneous* functor, since $\mathrm{MC}_{(h,g)}(B \times_A C) \cong \mathrm{MC}_{(h,g)}(B) \times_{\mathrm{MC}_{(h,g)}(A)} \mathrm{MC}_{(h,g)}(C)$.

As in the case of a differential graded Lie algebra (see Definition I.3.26), we can define for each $(A, m_A) \in \mathbf{Art}$ a gauge action over $\mathrm{MC}_{(h,g)}(A)$.

DEFINITION III.1.11. The gauge action of $\exp(L^0 \otimes m_A) \times \exp(N^0 \otimes m_A)$ over $\mathrm{MC}_{(h,g)}(A)$ is given by:

$$(e^a, e^b) * (x, y, e^p) = (e^a * x, e^b * y, e^{g(b)} e^p e^{-h(a)}).$$

This is well defined since

$$e^{g(b)} e^p e^{-h(a)} * h(e^a * x) = e^{g(b)} e^p * h(x) = e^{g(b)} * g(y) = g(e^b * y).$$

and so $(e^a, e^b) * (x, y, e^p) \in \mathrm{MC}_{(h,g)}(A)$.

In conclusion, it makes sense to consider the following functor.

DEFINITION III.1.12. The *deformation functor associated with a pair* (h, g) of morphisms of differential graded Lie algebras is:

$$\mathrm{Def}_{(h,g)} : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{Def}_{(h,g)}(A) = \frac{\mathrm{MC}_{(h,g)}(A)}{\exp(L^0 \otimes m_A) \times \exp(N^0 \otimes m_A)}.$$

REMARK III.1.13. In [**24**, Section 2], M. Manetti defined the functor Def_h associated with a morphism $h : L \longrightarrow M$ of DGLAs:

$$\mathrm{Def}_h : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$\mathrm{Def}_h(A) = \frac{\mathrm{MC}_h(A)}{\exp(L^0 \otimes m_A) \times \exp(dM^{-1} \otimes m_A)},$$

where the gauge action of $\exp(L^0 \otimes m_A) \times \exp(dM^{-1} \otimes m_A)$ is given by the following formula

$$(e^a, e^{dm}) * (x, e^p) = (e^a * x, e^{dm} e^p e^{-h(a)}), \qquad \forall\, a \in L^0 \otimes m_A, m \in M^{-1} \otimes m_A.$$

Therefore, if we take $N = 0$ and $g = 0$, the new functor $\mathrm{Def}_{(h,g)}$ reduces to the old one Def_h (M is concentrated in non negative degree).

By choosing $N = M = 0$ and $h = g = 0$, $\mathrm{Def}_{(h,g)}$ reduces to the Maurer-Cartan functor Def_L associated with the DGLA L.

The name deformation functor is justified by the theorem below.

THEOREM III.1.14. $\mathrm{Def}_{(h,g)}$ *is a deformation functors, i.e., it satisfies the conditions of Definition I.1.10.*

PROOF. If $A = \mathbb{K}$, then $\mathrm{Def}_{(h,g)}(\mathbb{K}) = \{\text{one element}\}$; therefore, *ii)* of Definition I.1.10 holds, i.e., $\mathrm{Def}_{(h,g)}(B \times C) = \mathrm{Def}_{(h,g)}(B) \times \mathrm{Def}_{(h,g)}(C)$.

Let $\beta : B \longrightarrow A$ and $\gamma : C \longrightarrow A$ be morphisms in $\mathbf{Art}$ and $(v, w) \in \mathrm{Def}_{(h,g)}(B) \times_{\mathrm{Def}_{(h,g)}(A)} \mathrm{Def}_{(h,g)}(C)$. Then we are looking for a lifting $z \in \mathrm{Def}_{(h,g)}(B \times_A C)$, whenever $\beta : B \longrightarrow A$ is surjective.

Let $(x, y, e^p) \in \mathrm{MC}_{(h,g)}(B)$ and $(s, t, e^r) \in \mathrm{MC}_{(h,g)}(C)$ be liftings for v and w, respectively.

By hypothesis, $\beta(v)=\gamma(w)\in \mathrm{Def}_{(h,g)}(A)$; therefore, $\beta(x,y,e^p)$ and $\gamma(s,t,e^r)$ are gauge equivalent in $\mathrm{MC}_{(h,g)}(A)$, i.e., there exist $a\in L^0\otimes m_A$ and $b\in N^0\otimes m_A$ such that

$$e^a*\beta(x)=\gamma(s)\qquad e^b*\beta(y)=\gamma(t)\qquad e^{g(b)}e^{\beta(p)}e^{-h(a)}=e^{\gamma(r)}.$$

Let $c\in L^0\otimes m_B$ such that $\beta(c)=a$ and $d\in N^0\otimes m_B$ such that $\beta(d)=b$.

Up to substituting (x,y,e^p) with its gauge equivalent element $(e^c*x, e^d*y, e^{g(d)}e^pe^{-h(c)})$ (they both lift v), we can assume that[a] $\beta(x,y,e^p)=\gamma(s,t,e^r)\in \mathrm{MC}_{(h,g)}(A)$.

Since $\mathrm{MC}_{(h,g)}(B\times_A C)=\mathrm{MC}_{(h,g)}(B)\times_{\mathrm{MC}_{(h,g)}(A)}\mathrm{MC}_{(h,g)}(C)$, a lifting $z\in \mathrm{MC}_{(h,g)}(B\times_A C)$ is well defined and so it is sufficient to take its class $[z]\in \mathrm{Def}_{(h,g)}(B\times_A C)$.

□

REMARK III.1.15. Consider the functor $\mathrm{Def}_{(h,g)}$. Then the projection ϱ on the second factor:

$$\varrho:\mathrm{Def}_{(h,g)}\longrightarrow \mathrm{Def}_N,$$

$$\mathrm{Def}_{(h,g)}(A)\ni (x,y,e^p)\xrightarrow{\varrho} y\in \mathrm{Def}_N(A)$$

is a morphism of deformation functors.

REMARK III.1.16. If the morphism h *is injective*, then for each $(A,m_A)\in$ **Art** the functor $\mathrm{MC}_{(h,g)}$ has the following form:

$$\mathrm{MC}_{(h,g)}(A)=\{(x,e^p)\in (N^1\otimes m_A)\times \exp(M^0\otimes m_A)|$$

$$dx+\frac{1}{2}[x,x]=0,\ e^{-p}*g(x)\in L^1\otimes m_A\}.$$

In this case the gauge equivalence $\sim$ is given by

$$(x,e^p)\sim (e^b*x, e^{g(b)}e^pe^a),\qquad \text{with } a\in L^0\otimes m_A \text{ and } b\in N^0\otimes m_A.$$

LEMMA III.1.17. *The projection* $\pi:\mathrm{MC}_{(h,g)}\longrightarrow \mathrm{Def}_{(h,g)}$ *is a smooth morphism of functors.*

PROOF. Let $\beta:B\longrightarrow A$ be a surjection in **Art**; we prove that

$$\mathrm{MC}_{(h,g)}(B)\longrightarrow \mathrm{Def}_{(h,g)}(B)\times_{\mathrm{Def}_{(h,g)}(A)}\mathrm{MC}_{(h,g)}(A)$$

is surjective.

Let $((x,y,e^p),(l,n,e^m))\in \mathrm{Def}_{(h,g)}(B)\times_{\mathrm{Def}_{(h,g)}(A)}\mathrm{MC}_{(h,g)}(A)$; that is, the class of (l,n,m) and $\beta(x,y,e^p)$ are the same element in $\mathrm{Def}_{(h,g)}(A)$.

Then there exists $(a,b)\in \exp(L^0\otimes m_A)\times \exp(N^0\otimes m_A)$ such that

$$(l,n,e^m)=(e^a,e^b)*(\beta(x,y,e^p))=(e^a*\beta(x), e^b*\beta(y), e^{g(b)}e^{\beta(p)}e^{-h(a)}).$$

---

[a] $\beta(e^c*x, e^d*y, e^{g(d)}e^pe^{-h(c)})=(e^{\beta(c)}*\beta(x), e^{\beta(d)}*\beta(y), e^{g(\beta(d))}e^{\beta(p)}e^{-h(\beta(c))})=(e^a*\beta(x), e^b*\beta(y), e^{g(b)}e^{\beta(p)}e^{-h(a)})=\gamma(s,t,e^r)$

Let $c \in L^0 \otimes m_B$ be a lifting of a and $d \in N^0 \otimes m_B$ be a lifting of b. Then

$$t = (e^c * x, e^d * y, e^{g(d)} e^p e^{-h(c)})$$

lies in $\mathrm{MC}_{(h,g)}(B)$ and is a lifting of $((x, y, e^p), (l, n, e^m))$.

Actually, t is gauge equivalent to (x, y, e^p) and

$$\beta(t) = (e^{\beta(c)} * \beta(x), e^{\beta(d)} * \beta(y), e^{g(\beta(d))} e^{\beta(p)} e^{-h(\beta(c))}) =$$
$$(e^a * \beta(x), e^b * \beta(y), e^{g(b)} e^{\beta(p)} e^{-h(a)}) = (l, n, e^m).$$

$\square$

III.1.3. Tangent and obstruction spaces of $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$. By Definition III.1.8, *the tangent space of* $\mathrm{MC}_{(h,g)}$ is

$$\mathrm{MC}_{(h,g)}(\mathbb{K}[\varepsilon]) =$$
$$= \{(x, y, e^p) \in (L^1 \otimes \mathbb{K}\varepsilon) \times (N^1 \otimes \mathbb{K}\varepsilon) \times \exp(M^0 \otimes \mathbb{K}\varepsilon) |$$
$$dx = dy = 0, h(x) - g(y) - dp = 0\}$$
$$\cong \{(x, y, p) \in L^1 \times N^1 \times M^0 |\ dx = dy = 0, g(y) = h(x) - dp\} =$$
$$\ker(D : \mathrm{C}^1_{(h,g)} \longrightarrow \mathrm{C}^2_{(h,g)}).$$

By Definition III.1.12, *the tangent space of* $\mathrm{Def}_{(h,g)}$ is

$$\mathrm{Def}_{(h,g)}(\mathbb{K}[\varepsilon]) =$$
$$\frac{\{(x, y, p) \in L^1 \times N^1 \times M^0 |\ dx = dy = 0, g(y) = h(x) - dp\}}{\{(-da, -db, g(b) - h(a)) |\ a \in (L^0 \otimes \mathbb{K}[\varepsilon]), b \in (N^0 \otimes \mathbb{K}[\varepsilon])\}}.$$
$$\cong H^1(\mathrm{C}^{\cdot}_{(h,g)}).$$

Remark III.1.18. In the last equality we use the extra hypothesis that $M^{-1} = 0$.

As to the obstruction space of $\mathrm{MC}_{(h,g)}$, we prove below that it is naturally contained in $H^2(\mathrm{C}^{\cdot}_{(h,g)})$. Since $\pi : \mathrm{MC}_{(h,g)} \longrightarrow \mathrm{Def}_{(h,g)}$ is smooth, Corollary I.1.35 implies that the obstruction space of $\mathrm{Def}_{(h,g)}$ is also contained in $H^2(\mathrm{C}^{\cdot}_{(h,g)})$.

Lemma III.1.19. *$H^2(\mathrm{C}^{\cdot}_{(h,g)})$ is a complete obstruction space for* $\mathrm{MC}_{(h,g)}$.

Proof. Let

$$0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

be a small extension and $(x, y, e^p) \in \mathrm{MC}_{(h,g)}(A)$.

Since α is surjective there exist $\tilde{x} \in L^1 \otimes m_B$ that lifts x, $\tilde{y} \in N^1 \otimes m_B$ that lifts y, and $q \in M^0 \otimes m_B$ that lifts p. Let

$$l = d\tilde{x} + \frac{1}{2}[\tilde{x}, \tilde{x}] \in L^2 \otimes m_B$$

and

$$k = d\tilde{y} + \frac{1}{2}[\tilde{y}, \tilde{y}] \in N^2 \otimes m_B.$$

As in Lemma I.3.24, we can easily prove that $\alpha(l) = \alpha(k) = dl = dk = 0$; then $l \in H^2(L) \otimes J$ and $k \in H^2(N) \otimes J$.

Let $r = -g(\tilde{y}) + e^q * h(\tilde{x}) \in M^1 \otimes m_B$; in particular, $\alpha(r) = 0$ and so $r \in M^1 \otimes J$.

Next, we prove that $(l, k, r) \in Z^2(\mathrm{C}^{\cdot}_{(h,g)}) \otimes J$.

Since $dl = dk = 0$, it remains to prove that $-dr - g(k) + h(l) = 0$.

By definition, $h(\tilde{x}) = e^{-q} * (g(\tilde{y}) + r) = r + e^{-q} * g(\tilde{y})$ (in the last equalities we use Example I.3.28) and so

$$h(l) = d(h(\tilde{x})) + \frac{1}{2}[h(\tilde{x}), h(\tilde{x})] = dr + d(e^{-q} * g(\tilde{y})) + \frac{1}{2}[e^{-q} * g(\tilde{y}), e^{-q} * g(\tilde{y})].$$

Let $A = d(e^{-q} * g(\tilde{y}))$ and $B = [e^{-q} * g(\tilde{y}), e^{-q} * g(\tilde{y})]$. Therefore, it is sufficient to prove that $A + \dfrac{1}{2}B = g(k)$.

We have

$$B = [e^{-q} * g(\tilde{y}), e^{-q} * g(\tilde{y})] = [e^{[-q,]'}(d + g(\tilde{y})) - d, e^{[-q,]'}(d + g(\tilde{y})) - d]' =$$

$$[e^{[-q,]'}(d + g(\tilde{y})), e^{[-q,]'}(d + g(\tilde{y}))]' + [e^{[-q,]'}(d + g(\tilde{y})), -d]' + [-d, e^{[-q,]'}(d + g(\tilde{y}))]' =$$

$$e^{[-q,]'}[d + g(\tilde{y}), d + g(\tilde{y})]' - 2[d, e^{[-q,]'}(d + g(\tilde{y}))]' =$$

$$e^{[-q,]'}(2dg(\tilde{y}) + [g(\tilde{y}), g(\tilde{y})]) - 2[d, e^{[-q,]'}(d + g(\tilde{y})) - d]' = 2e^{[-q,]}(g(k)) - 2A.$$

By assumption, $g(k) \in M^2 \otimes J$ and so $e^{[-q,]}(g(k)) = g(k)$.

Therefore,

$$A + \frac{1}{2}B = A + \frac{1}{2}(2g(k) - 2A) = g(k)$$

and so $[(l, k, r)] \in H^2(\mathrm{C}^{\cdot}_{(h,g)}) \otimes J$.

This class does not depend on the choice of the liftings.

Actually, let $\tilde{x}' \in L^1 \otimes m_B$ be another lifting of x. Then $\tilde{x}' = \tilde{x} + j_x$, for some $j_x \in L^1 \otimes J$ and so

$$l' = d\tilde{x}' + \frac{1}{2}[\tilde{x}', \tilde{x}'] = d\tilde{x} + dj_x + \frac{1}{2}[\tilde{x}, \tilde{x}] = l + dj_x.$$

Analogously, if $\tilde{y}'$ is another lifting of y then there exists $j_y \in N^1 \otimes J$ such that

$$k' = k + dj_y.$$

Moreover,

$$r' = -g(\tilde{y}') + e^q * h(\tilde{x}') = -g(\tilde{y}) - g(j_y) + e^q * h(\tilde{x}) + h(j_x) = r - g(j_y) + h(j_x).$$

Therefore, $[(l', k', r')] = [(l + dj_x, k + dj_y, r - g(j_y) + h(j_x))] = [(l, k, r)] \in H^2(\mathrm{C}^{\cdot}_{(h,g)}) \otimes J$.

In conclusion, $[(l, k, r)] \in H^2(\mathrm{C}^{\cdot}_{(h,g)}) \otimes J$ is the obstruction class associated with the element $(x, y, e^p) \in \mathrm{MC}_{(h,g)}(A)$.

If this class vanishes, then there exists $(u, v, z) \in \mathrm{C}^1_{(h,g)} \otimes J$ such that $(du, dv, -dz - g(v) + h(u)) = (l, k, r)$. In this case, define $\overline{x} = \tilde{x} - u$,

$\overline{y} = \tilde{y} - v$ and $\overline{z} = q - z$. Then $(\overline{x}, \overline{y}, e^{\overline{z}}) \in \mathrm{MC}_{(h,g)}(B)$ and it lifts (x, y, e^p). Actually,

$$d\overline{x} + [\overline{x}, \overline{x}] = d\tilde{x} - du + [\tilde{x}, \tilde{x}] = l - du = 0,$$

$$d\overline{y} + [\overline{y}, \overline{y}] = d\tilde{y} - dv + [\tilde{y}, \tilde{y}] = k - dv = 0$$

and

$$\begin{gathered} g(\overline{y}) - e^{\overline{z}} * h(\overline{x}) = g(\tilde{y}) - g(v) - e^{q-z} * (h(\tilde{x}) - h(u)) = \text{[b]} \\ g(\tilde{y}) - g(v) - e^{q-z} * h(\tilde{x}) + h(u) = \\ (g(\tilde{y}) - e^q * h(\tilde{x})) - dz - g(v) + h(u) = -r + r = 0. \end{gathered}$$

□

III.1.4. Properties.

Lemma III.1.20. *Let $h : L \longrightarrow M$ and $g : N \longrightarrow M$ be morphisms of abelian DGLAs. Then the functor* $\mathrm{MC}_{(h,g)}$ *is smooth.*

Proof. We have to prove that for every surjection $\varphi : B \longrightarrow A \in$ **Art** the map

$$\mathrm{MC}_{(h,g)}(B) \longrightarrow \mathrm{MC}_{(h,g)}(A)$$

is surjective. By hypothesis, we have

$$\begin{gathered} \mathrm{MC}_{(h,g)}(A) = \{(x, y, e^p) \in (L^1 \otimes m_A) \times (N^1 \otimes m_A) \times \exp(M^0 \otimes m_A) | \\ dx = dy = 0, \qquad g(y) = e^p * h(x) = h(x) - dp\}. \end{gathered}$$

This implies that the Maurer-Cartan equation reduces to the linear equations $dx = dy = 0, g(y) + dp - h(x) = 0$; then $\mathrm{MC}_{(h,g)}(A) = Z^1(\mathrm{C}^{\cdot}_{(h,g)} \otimes m_A) = Z^1(\mathrm{C}^{\cdot}_{(h,g)}) \otimes m_A$.

Since $Z^1(\mathrm{C}^{\cdot}_{(h,g)} \otimes m_A) \twoheadrightarrow Z^1(\mathrm{C}^{\cdot}_{(h,g)} \otimes m_B)$, the lifting exists.

□

Remark III.1.21. Every commutative diagram of morphisms of DGLAs

$$\begin{tikzcd} & L \arrow[r, "\alpha'"] \arrow[d, "h"'] & P \arrow[d, "\eta"] \\ & M \arrow[r, "\alpha"] & Q \\ N \arrow[ur, "g"] \arrow[r, "\alpha''"] & R \arrow[ur, "\mu"'] & \end{tikzcd}$$

induces a *morphism* $\varphi^{\cdot}$ *of complexes*

$$\mathrm{C}^i_{(h,g)} \ni (l, n, m) \xmapsto{\varphi^i} (\alpha'(l), \alpha''(n), \alpha(m)) \in \mathrm{C}^i_{(\eta,\mu)}$$

and a natural *transformation* F of the associated deformation functors:

$$F : \mathrm{Def}_{(h,g)} \longrightarrow \mathrm{Def}_{(\eta,\mu)} .$$

---

[b] See Example I.3.28.

Then, we obtain the following proposition that is a generalization of [**24**, Proposition 2.3].

PROPOSITION III.1.22. *Let*

$$\begin{tikzcd} & L \arrow[r, "\alpha'"] \arrow[d, "h"'] & P \arrow[d, "\eta"] \\ & M \arrow[r, "\alpha"] & Q \\ N \arrow[ur, "g"] \arrow[r, "\alpha''"] & R \arrow[ur, "\mu"'] & \end{tikzcd}$$

be a commutative diagram of differential graded Lie algebras. If the functor $\mathrm{Def}_{(\eta,\mu)}$ *is smooth, then the obstruction space of* $\mathrm{Def}_{(h,g)}$ *is contained in the kernel of the map*

$$H^2(\mathrm{C}^{\cdot}_{(h,g)}) \longrightarrow H^2(\mathrm{C}^{\cdot}_{(\eta,\mu)}).$$

PROOF. The morphism $F : \mathrm{Def}_{(h,g)} \longrightarrow \mathrm{Def}_{(\eta,\mu)}$ induces a linear map between obstruction spaces. If $\mathrm{Def}_{(\eta,\mu)}$ is smooth, then its obstruction space is zero (Proposition I.1.31).

□

THEOREM III.1.23. *If* $\varphi^{\cdot} : \mathrm{C}^{\cdot}_{(h,g)} \longrightarrow \mathrm{C}^{\cdot}_{(\eta,\mu)}$ *is a quasi-isomorphism of complexes then* $F : \mathrm{Def}_{(h,g)} \longrightarrow \mathrm{Def}_{(\eta,\mu)}$ *is an isomorphism of functors.*

The proof of this theorem is postponed to the end of Section III.2.4.

Next, let

$$\begin{tikzcd} H \arrow[r, "\alpha"] \arrow[d, "\beta"'] & L \arrow[d, "h"] \\ N \arrow[r, "g"] & M \end{tikzcd}$$

be a commutative diagram of morphisms of DGLAs. Thus, it induces a morphism of complexes

$$H \xrightarrow{\varphi^{\cdot}} \mathrm{C}^{\cdot}_{(h,g)} \qquad \text{with} \qquad \varphi(x) = (\alpha(x), \beta(x), 0),$$

and a morphism of functors

$$\mathrm{Def}_H \xrightarrow{F} \mathrm{Def}_{(h,g)} \qquad \text{with} \qquad F(x) = (\alpha(x), \beta(x), 0).$$

THEOREM III.1.24. *With the notation above, if the morphism* $\varphi^{\cdot}$ *is a quasi-isomorphism then* F *is an isomorphism of functors.*

The proof of this theorem too is postponed to the end of Section III.2.4.

III.2. Extended Deformation Functors

In this section we study the extended deformation functors. In particular, we are interested in the functors $\widetilde{\mathrm{MC}}_{(h,g)}$ and $\widetilde{\mathrm{Def}}_{(h,g)}$ which are a generalization of the functors $\mathrm{MC}_{(h,g)}$ and $\mathrm{Def}_{(h,g)}$ introduced in Section III.1. Here, we remove the restrictive hypothesis of M concentrated in non negative degrees.

The main references for this chapter are [**21**], [**23**, Sections 5.7 and 5.8] and [**24**, Sections 6 and 7].

III.2.1. Notation. We denote by:

$\mathbf{C}$ the category of nilpotent (associative and commutative) differential graded algebras which are finite dimensional as $\mathbb{K}$-vector spaces.

$\mathbf{C}_0$ the full subcategory of $\mathbf{C}$ of algebras with trivial multiplication.

Example III.2.1. Define the complex $(\Omega = \Omega_0 \oplus \Omega_1, d)$, where $\Omega_0 = \mathbb{K}$, $\Omega_1 = \mathbb{K}[-1]$ and $d : \Omega_0 \longrightarrow \Omega_1$ is the canonical linear isomorphism $d(1[0]) = 1[-1]$. $\Omega \in \mathbf{C}_0$ and the projection $p : \Omega \longrightarrow \Omega_0 = \mathbb{K}$ and the inclusion $\Omega_1 \longrightarrow \Omega$ are morphisms in $\mathbf{C}_0$.

Moreover, $\Omega[n] = \mathbb{K}[n] \otimes \Omega$ is an acyclic complex in $\mathbf{C}_0$, for each $n \in \mathbb{N}$.

III.2.2. Definition of extended functors. Let $A \in \mathbf{C}$ and $J \subset A$ a differential ideal; then $J \in \mathbf{C}$ and the inclusion $J \longrightarrow A$ is a morphism of differential graded algebras.

Definition III.2.2. A *small extension* in $\mathbf{C}$ is a short exact sequence

$$0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

such that α is a morphism in $\mathbf{C}$ and J is an ideal of B such that $BJ = 0$; in addition, it is called *acyclic* if J is an acyclic complex, or equivalently α is a quasi-isomorphism.

Definition III.2.3. A covariant functor $F : \mathbf{C} \longrightarrow \mathbf{Set}$ is called a *predeformation functor* if the following conditions are satisfied:

III.2.3.1) $F(0) = \{*\}$ is the one point set.

III.2.3.2) For every $A, B \in \mathbf{C}$, the natural map

$$F(A \times B) \longrightarrow F(A) \times F(B)$$

is bijective.

III.2.3.3) For every surjective morphism $\alpha : A \longrightarrow C$ in $\mathbf{C}$, with C an acyclic complex in $\mathbf{C}_0$, the natural morphism

$$F(\ker(\alpha)) = F(A \times_C 0) \longrightarrow F(A) \times_{F(C)} F(0) = F(A)$$

is bijective.

III.2.3.4) For any pair of morphisms $\beta : B \longrightarrow A$ and $\gamma : C \longrightarrow A$ in $\mathbf{C}$, with β surjective, the natural map

$$F(B \times_A C) \longrightarrow F(B) \times_{F(A)} F(C)$$

is surjective.

III.2.3.5) For every small acyclic extension

$$0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

the induced map $F(B) \longrightarrow F(A)$ is surjective.

Definition III.2.4. A covariant functor $F : \mathbf{C} \longrightarrow \mathbf{Set}$ is called a *deformation functor* if it is a predeformation functor and $F(J) = 0$ for every acyclic complex $J \in \mathbf{C}_0$.

III.2.2.1. *Examples.* Let L be a differential graded Lie algebra and $A \in \mathbf{C}$. Then $L \otimes A$ has a natural structure of (nilpotent) DGLA given by:

$$(L \otimes A)^i = \bigoplus_{j \in \mathbb{Z}} L^j \otimes A^{i-j};$$

$$d(x \otimes a) = dx \otimes a + (-1)^{\deg(x)} x \otimes da;$$

$$[x \otimes a, y \otimes b] = (-1)^{\deg(a)\deg(y)} [x, y] \otimes ab.$$

Definition III.2.5. The *extended Maurer-Cartan* functor associated with a DGLA L is

$$\widetilde{\mathrm{MC}}_L(A) = \{x \in (L \otimes A)^1 |\, dx + \frac{1}{2}[x, x] = 0\}.$$

Lemma III.2.6. *$\widetilde{\mathrm{MC}}_L$ is a predeformation functor.*

Proof. See [**21**, Lemma 2.15]. We will also give a proof in Section III.2.4, since it is a particular case of Theorem III.2.20. □

Remark III.2.7. We note that, for each $C \in \mathbf{C}_0$, we have:

$$\widetilde{\mathrm{MC}}_L(C) = \{x \in (L \otimes C)^1 |\, dx = 0\} = Z^1(L \otimes C).$$

Definition III.2.8. The *extended deformation* functor associated with a DGLA L is

$$\widetilde{\mathrm{Def}}_L(A) = \frac{\{x \in (L \otimes A)^1 |\, dx + \frac{1}{2}[x, x] = 0\}}{\text{gauge action of } \exp{(L \otimes A)^0}},$$

where the gauge action of $a \in (L \otimes A)^0$ is the analogous of the not extended case, i.e.,

$$e^a * x := x + \sum_{n=0}^{\infty} \frac{[a, -]^n}{(n+1)!}([a, x] - da).$$

Lemma III.2.9. *$\widetilde{\mathrm{Def}}_L$ is a deformation functor.*

Proof. See [**21**, Theorem 2.16]. We will also give a proof in Section III.2.4, since it is a particular case of Theorem III.2.26.

□

Remark III.2.10. For each $C \in \mathbf{C}_0$ we note that:

$$\widetilde{\mathrm{Def}}_L(C) = \frac{\{x \in (L \otimes C)^1 |\, dx = 0\}}{\{da |\, a \in (L \otimes C)^0\}} = H^1(L \otimes C).$$

III.2.3. Properties. As in the not extended case, for every predeformation functor F and every $C \in \mathbf{C}_0$, there exists a natural structure of vector space on $F(A)$ with:

- addition given by the map $F(C\times C) = F(C)\times F(C) \xrightarrow{+} F(C)$ induced by $C \times C \xrightarrow{+} C$;
- scalar multiplication by s given by the map $F(C) \xrightarrow{\cdot s} F(C)$ induced by $C \xrightarrow{\cdot s} C$.

Remark III.2.11. 1) For any morphism $B \longrightarrow A$ in $\mathbf{C}_0$, the induced map $F(B) \longrightarrow F(A)$ is $\mathbb{C}$-linear.

2) For any natural transformation of predeformation functors $F \longrightarrow G$, the induced map $F(C) \longrightarrow G(C)$ is $\mathbb{C}$-linear for each $C \in \mathbf{C}_0$.

Definition III.2.12. Let F be a predeformation functor, the *tangent space* of F is the graded vector space $TF[1]$, where

$$TF = \bigoplus_{n\in\mathbb{Z}} T^nF, \quad T^{n+1}F = TF[1]^n = \operatorname{coker}(F(\Omega[n]) \xrightarrow{p} F(\mathbb{K}[n])), n \in \mathbb{Z}$$

and p is the linear map induced by the projection $\Omega[n] \longrightarrow \mathbb{K}[n]$ (see Example III.2.1).

In particular, if F is a deformation functor then $F(\Omega[n]) = 0$ for every n. Therefore, $T^{n+1}F = TF[1]^n = F(\mathbb{K}[\varepsilon])$, where ε is an indeterminate of degree $-n \in \mathbb{Z}$, such that $\varepsilon^2 = 0$.

Definition III.2.13. A natural transformation of predeformation functors $F \longrightarrow G$ is called a *quasi-isomorphism* if it induces isomorphisms on tangent spaces, i.e., $T^nF \cong T^nG$.

Theorem III.2.14. *(Inverse function theorem) A natural transformation of deformation functors is an isomorphism if and only if it is a quasi-isomorphism.*

Proof. See [**21**, Corollary 3.2] or [**23**, Corollary 5.72]. □

Theorem III.2.15. *(Manetti) Let F be a predeformation functor, then there exist a deformation functor F^+ and a natural transformation $\nu : F \longrightarrow F^+$, that is a quasi-isomorphism, such that for every deformation functor G and every natural transformation $\phi : F \longrightarrow G$ there exists a unique natural transformation $\psi : F^+ \longrightarrow G$ such that $\phi = \psi\nu$.*

Proof. See [**21**, Theorem 2.8]. □

Remark III.2.16. Given a natural transformation of predeformation functors $\alpha : F \longrightarrow G$, there is a natural transformation of associated deformation functors $\alpha^+ : F^+ \longrightarrow G^+$. Actually, let $\eta : G \longrightarrow G^+$. Then by composition we get $\beta = \eta \circ \alpha : F \longrightarrow G^+$ and so, by Theorem III.2.15, there exists $\alpha^+ : F^+ \longrightarrow G^+$.

THEOREM III.2.17. *(Manetti) Let S be a complex of vector spaces and assume that the functor*

$$\mathbf{C}_0 \longrightarrow \mathbf{Set},$$
$$C \longmapsto Z^1(S \otimes C)$$

is the restriction of a predeformation functor F. Then for every complex $C \in \mathbf{C}_0$, the equality $F^+(C) = H^1(S \otimes C)$ holds; in particular, $T^iF^+ = H^i(S)$.

PROOF. See [**21**, Lemma 2.10]. □

COROLLARY III.2.18. *(Manetti) For every differential graded Lie algebra L there exists a natural isomorphism $\widetilde{\mathrm{MC}}^+_L \cong \widetilde{\mathrm{Def}}_L$.*

PROOF. By Remark III.2.7 and Theorem III.2.17, we have that $T^i\widetilde{\mathrm{MC}}^+_L = H^i(L)$. Therefore, the natural projection $\widetilde{\mathrm{MC}}_L \longrightarrow \widetilde{\mathrm{Def}}_L$ induces (by Theorem III.2.15) a natural transformation $\widetilde{\mathrm{MC}}^+_L \longrightarrow \widetilde{\mathrm{Def}}_L$ which is an isomorphism on tangent spaces. □

III.2.4. Extended deformation functor of a pair of morphisms. Let L, M, N be DGLAs, and $h : L \longrightarrow M$ and $g : N \longrightarrow M$ be morphisms of DGLAs:

$$\begin{tikzcd} & L \arrow[d, "h"] \\ N \arrow[r, "g"'] & M. \end{tikzcd}$$

DEFINITION III.2.19. The *extended Maurer-Cartan functor associated with the pair* (h, g) is

$$\widetilde{\mathrm{MC}}_{(h,g)} : \mathbf{Art} \longrightarrow \mathbf{Set},$$
$$\widetilde{\mathrm{MC}}_{(h,g)}(A) = \{(x, y, e^p) \in (L \otimes A)^1 \times (N \otimes A)^1 \times \exp(M \otimes A)^0 |$$
$$dx + \frac{1}{2}[x, x] = 0,\ dy + \frac{1}{2}[y, y] = 0,\ g(y) = e^p * h(x)\}.$$

THEOREM III.2.20. $\widetilde{\mathrm{MC}}_{(h,g)}$ *is a predeformation functor.*

PROOF. $\widetilde{\mathrm{MC}}_{(h,g)}(0) = 0$ and so (III.2.3.1) is satisfied.
For any pair of morphisms $\beta : B \longrightarrow A$ and $\gamma : C \longrightarrow A$ in $\mathbf{C}$, we have

$$\widetilde{\mathrm{MC}}_{(h,g)}(B \times_A C) = \widetilde{\mathrm{MC}}_{(h,g)}(B) \times_{\widetilde{\mathrm{MC}}_{(h,g)}(A)} \widetilde{\mathrm{MC}}_{(h,g)}(C)$$

and so $\widetilde{\mathrm{MC}}_{(h,g)}$ satisfies properties (III.2.3.2),(III.2.3.3) and (III.2.3.4).
Let $0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$ be an acyclic small extension and $(x, y, e^p) \in \widetilde{\mathrm{MC}}_{(h,g)}(A)$. We want to prove the existence of a lifting $(\overline{x}, \overline{y}, e^{\overline{p}}) \in \widetilde{\mathrm{MC}}_{(h,g)}(B)$ such that the induced map $\widetilde{\mathrm{MC}}_{(h,g)}(B) \longrightarrow \widetilde{\mathrm{MC}}_{(h,g)}(A)$ is surjective.

Since α is surjective, there exists $(r, s, e^t) \in (L \otimes B)^1 \times (N \otimes B)^1 \times \exp(M \otimes B)^0$ such that $\alpha(r) = x, \alpha(s) = y$ and $\alpha(t) = p$.

Let $l \in (L \otimes J)^2$ and $k \in (N \otimes J)^2$ be defined as follows

$$l = dr + \frac{1}{2}[r, r] \quad \text{and} \quad k = ds + \frac{1}{2}[s, s].$$

Then

$$dl = \frac{1}{2}d[r, r] = [dr, r] = [l, r] - \frac{1}{2}[[r, r], r]$$

and the same holds for k, i.e.,

$$dk = [k, s] - \frac{1}{2}[[s, s], s].$$

Since $B \cdot J = 0$, we have $[l, r] = [k, s] = 0$; moreover, using Jacobi's identity (see Remark I.3.7), $[[r, r], r] = [[s, s], s] = 0$. This implies that $dl = dk = 0$.

By hypothesis, J is acyclic and so, by the Künneth formula, the complexes $L \otimes J$ and $N \otimes J$ are acyclic. Thus, there exist $w \in (L \otimes J)^1$ and $z \in (N \otimes J)^1$, such that $dw = l$ and $dz = k$.

Let

$$\overline{x} = r - w \in (L \otimes B)^1 \qquad \text{and} \qquad \overline{y} = s - z \in (N \otimes B)^1;$$

we have

$$\alpha(\overline{x}) = \alpha(r) - \alpha(w) = \alpha(r) = x, \qquad \alpha(\overline{y}) = \alpha(s) - \alpha(z) = \alpha(s) = y,$$

$$d\overline{x} + \frac{1}{2}[\overline{x}, \overline{x}] = dr - dw + \frac{1}{2}[r, r] = l - l = 0 \qquad \text{and} \qquad d\overline{y} + \frac{1}{2}[\overline{y}, \overline{y}] = 0.$$

Therefore, $\overline{x}$ and $\overline{y}$ lift x and y , respectively, and they satisfy the Maurer-Cartan equation.

Let $z = e^t * h(\overline{x}) - g(\overline{y}) \in (M \otimes B)^1$. Since $\alpha(z) = e^{\alpha(t)} * h(\alpha(\overline{x})) - g(\alpha(\overline{y})) = e^p * h(x) - g(y) = 0$, $z \in (M \otimes J)^1$. Moreover, $dz = 0$; indeed

$$2dz = 2d(e^t * h(\overline{x})) - 2d(g(\overline{y})) = -[e^t * h(\overline{x}), e^t * h(\overline{x})] + [g(\overline{y}), g(\overline{y})] =$$

$$-[e^t * h(\overline{x}), e^t * h(\overline{x})] + [g(\overline{y}), e^t * h(\overline{x})] - [g(\overline{y}), e^t * h(\overline{x})] + [g(\overline{y}), g(\overline{y})] =$$

$$-[e^t * h(\overline{x}) - g(\overline{y}), e^t * h(\overline{x})] - [g(\overline{y}), e^t * h(\overline{x}) - g(\overline{y})] = 0,$$

$(z = e^t * h(\overline{x}) - g(\overline{y}) \in (M \otimes J)^1$ and so $[e^t * h(\overline{x}) - g(\overline{y}), -] = 0)$.

Since $M \otimes J$ is acyclic, there exists $v \in (M \otimes J)^0$ such that $z = dv$. Therefore,

$$e^t * h(\overline{x}) = z + g(\overline{y}) = dv + g(\overline{y}) = e^{-v} * g(\overline{y})$$

and so

$$e^v e^t * h(\overline{x}) = g(\overline{y}).$$

This implies that $e^{\overline{p}} = e^v e^t$ lifts e^p.

Then the triple $(\overline{x}, \overline{y}, e^{\overline{p}}) \in \widetilde{\mathrm{MC}}_{(h,g)}(B)$ lifts $(x, y, e^p) \in \widetilde{\mathrm{MC}}_{(h,g)}(A)$ and so (III.2.3.5) holds.

$\square$

Proof of Lemma III.2.6. It suffices to apply the previous theorem with $M = N = 0$. $\square$

Remark III.2.21. If the DGLA M is concentrated in non negative degree, then for every $(A, m_A) \in \mathbf{Art}$, $\widetilde{\mathrm{MC}}_{(h,g)}(m_A) = \mathrm{MC}_{(h,g)}(A)$.

Applying Theorem III.2.15, we can conclude the existence of a deformation functor $\widetilde{\mathrm{MC}}^{+}_{(h,g)}$ associated with $\widetilde{\mathrm{MC}}_{(h,g)}$.

Proposition III.2.22. $T^i\widetilde{\mathrm{MC}}^{+}_{(h,g)} \cong H^i(\mathrm{C}^{\cdot}_{(h,g)})$.

Proof. For any $C \in \mathbf{C}_0$ we have

$$\widetilde{\mathrm{MC}}_{(h,g)}(C) = \{(l, n, e^m) \in (L \otimes C)^1 \times (N \otimes C)^1 \times \exp(M \otimes C)^0 | \\ dl = dn = 0,\ g(n) = e^m * h(l) = h(l) - dm\},$$

and

$$Z^1(\mathrm{C}^{\cdot}_{(h,g)} \otimes C) = \{(l, n, m) \in (L \otimes C)^1 \times (N \otimes C)^1 \times (M \otimes C)^0 | \\ dl = dn = -dm - g(n) + h(l) = 0\}.$$

Therefore, $\widetilde{\mathrm{MC}}_{(h,g)}(-)_{|\mathbf{C}_0} = Z^1(\mathrm{C}^{\cdot}_{(h,g)} \otimes -)_{|\mathbf{C}_0}$. Then we can apply Theorem III.2.17 to complete the proof. $\square$

Next, we consider on $\widetilde{\mathrm{MC}}_{(h,g)}(A)$ the following equivalence relation $\approx$:

$$(x_1, y_1, e^{p_1}) \ \approx\ (x_2, y_2, e^{p_2})$$

if and only if there exist $a \in (L \otimes A)^0$, $b \in (N \otimes A)^0$ and $c \in (M \otimes A)^{-1}$ such that

$$x_2 = e^a * x_1, \qquad y_2 = e^b * y_1$$

and

$$e^{p_2} = e^{g(b)} e^T e^{p_1} e^{-h(a)}, \quad \text{with} \quad T = dc + [g(y_1), c] \in Stab_A(g(y_1)).$$

Lemma III.2.23. *The relation $\approx$ is an equivalence relation.*

Proof. The reflexivity is obvious. As to the symmetry and the transitivity, we use the following property of the irrelevant stabilizers $Stab_A(-)$(see Section I.3.4): for each $x \in \widetilde{MC}_M(A)$, $a \in (M \otimes A)^0$ and $T = dc + [x, c] \in Stab_A(x)$ ($c \in (M \otimes A)^{-1}$), there exists $f \in (M \otimes A)^{-1}$ such that

$$e^a e^T = e^{T'} e^a$$

where $T' = df + [y, f] \in Stab_A(y)$ and $y = e^a * x$.

Symmetry. Let $(x_1, y_1, e^{p_1}) \approx (x_2, y_2, e^{p_2})$. Therefore, there exist $a \in (L \otimes A)^0$, $b \in (N \otimes A)^0$ and $c \in (M \otimes A)^{-1}$ such that

$$x_2 = e^a * x_1, \qquad y_2 = e^b * y_1$$

and

$$e^{p_2} = e^{g(b)}e^T e^{p_1} e^{-h(a)}, \quad \text{with} \quad T = dc + [g(y_1), c].$$

This implies the existence of $f \in (M \otimes A)^{-1}$ such that

$$e^{p_2} = e^{T'}e^{g(b)}e^{p_1}e^{-h(a)} \quad \text{with} \quad T' = df + [g(y_2), f].$$

Thus, by choosing $\alpha = -a \in (L \otimes A)^0$, $\beta = -b \in (N \otimes A)^0$ and $\gamma = -f \in (M \otimes A)^{-1}$ yields

$$x_1 = e^{\alpha} * x_2, \qquad y_1 = e^{\beta} * y_2$$

and

$$e^{p_1} = e^{-g(b)}e^{-T'}e^{p_2}e^{h(a)} = e^{g(\beta)}e^{T''}e^{p_2}e^{-h(\alpha)} \quad \text{with} \quad T'' = d\gamma + [g(y_1), \gamma].$$

Then $(x_2, y_2, e^{p_2}) \approx (x_1, y_1, e^{p_1})$.

Transitivity. Suppose

$$(x_1, y_1, e^{p_1}) \approx (x_2, y_2, e^{p_2}) \quad \text{and} \quad (x_2, y_2, e^{p_2}) \approx (x_3, y_3, e^{p_3}).$$

Therefore, there exist $a_1, a_2 \in (L \otimes A)^0$, $b_1, b_2 \in (N \otimes A)^0$ and $c_1, c_2 \in (M \otimes A)^{-1}$ such that

$$x_2 = e^{a_1} * x_1, \qquad y_2 = e^{b_1} * y_1 \qquad e^{p_2} = e^{g(b_1)}e^{T_1}e^{p_1}e^{-h(a_1)}$$

and

$$x_3 = e^{a_2} * x_2, \qquad y_3 = e^{b_2} * y_2 \qquad e^{p_3} = e^{g(b_2)}e^{T_2}e^{p_2}e^{-h(a_2)}$$

with $T_1 = dc_1 + [g(y_1), c_1]$ and $T_2 = dc_2 + [g(y_2), c_2]$. Then

$$e^{p_3} = e^{g(b_2)}e^{T_2}e^{g(b_1)}e^{T_1}e^{p_1}e^{-h(a_1)}e^{-h(a_2)} =$$
$$e^{g(b_2)}e^{g(b_1)}e^{T_2{}'}e^{T_1}e^{p_1}e^{-h(a_1)}e^{-h(a_2)},$$

for some $c' \in (M \otimes A)^{-1}$ and $T_2{}' = dc' + [g(y_1), c']$.

Since $Stab_A(g(y_1))$ is a subgroup, there exists $c \in (M \otimes A)^{-1}$ such that $e^{T_2{}'}e^{T_1} = e^T$ with $T = dc + [g(y_1), c]$.

Let $a = a_2 \bullet a_1 \in (L \otimes A)^0$, $b = b_2 \bullet b_1 \in (N \otimes A)^0$ and $c \in (M \otimes A)^{-1}$ as above. Then

$$x_3 = e^a * x_1, \qquad y_3 = e^b * y_3, \qquad e^{p_3} = e^{g(b)}e^T e^{p_1}e^{-h(a)}$$

and so $(x_1, y_1, e^{p_1}) \approx (x_3, y_3, e^{p_3})$.

□

REMARK III.2.24. We note that this equivalence relation generalizes the equivalence relation induced by the gauge action given in Definition III.1.11, when M is concentrated in non negative degree.

DEFINITION III.2.25. Define the functor

$$\widetilde{\mathrm{Def}}_{(h,g)} : \mathbf{C} \longrightarrow \mathbf{Set},$$

$$\widetilde{\mathrm{Def}}_{(h,g)}(A) = \widetilde{\mathrm{MC}}_{(h,g)}(A)/\approx .$$

THEOREM III.2.26. $\widetilde{\mathrm{Def}}_{(h,g)} : \mathbf{C} \longrightarrow \mathbf{Set}$ *is a deformation functor with* $T^i\widetilde{\mathrm{Def}}_{(h,g)} = H^i(C^{\cdot}_{(h,g)})$.

Proof. We first prove that $\widetilde{\mathrm{Def}}_{(h,g)}$ is a predeformation functor.

Since $\widetilde{\mathrm{Def}}_{(h,g)}$ is the quotient of the predeformation functor $\widetilde{\mathrm{MC}}_{(h,g)}$, conditions (III.2.3.1) and (III.2.3.5) are verified.

An easy calculation shows that $\widetilde{\mathrm{Def}}_{(h,g)}$ satisfies (III.2.3.2).

Next, we verify condition (III.2.3.4). Let $\beta : B \longrightarrow A$ and $\gamma : C \longrightarrow A$ be morphisms in $\mathbf{C}$, with β surjective. We prove that the natural map $\widetilde{\mathrm{Def}}_{(h,g)}(B \times_A C) \longrightarrow \widetilde{\mathrm{Def}}_{(h,g)}(B) \times_{\widetilde{\mathrm{Def}}_{(h,g)}(A)} \widetilde{\mathrm{Def}}_{(h,g)}(C)$ is surjective.

Let $(x_1, y_1, e^{p_1}) \in \widetilde{\mathrm{MC}}_{(h,g)}(B)$ and $(x_2, y_2, e^{p_2}) \in \widetilde{\mathrm{MC}}_{(h,g)}(C)$, such that $\beta(x_1, y_1, e^{p_1})$ and $\gamma(x_2, y_2, e^{p_2})$ are the same element in $\widetilde{\mathrm{Def}}_{(h,g)}(A)$.

Then, there exist $a \in (L \otimes A)^0$, $b \in (N \otimes A)^0$ and $c \in (M \otimes A)^{-1}$ such that

$$\gamma(x_2) = e^a * \beta(x_1), \qquad \gamma(y_2) = e^b * \beta(y_1)$$

and

$$e^{\gamma(p_2)} = e^{g(b)} e^T e^{\beta(p_1)} e^{-h(a)} \quad \text{with } T = dc + [g(\beta(y_1)), c].$$

Let $\tilde{a} \in (L \otimes B)^0$ be a lifting of a, $\tilde{b} \in (N \otimes B)^0$ a lifting of b and $\tilde{c} \in (M \otimes B)^{-1}$ a lifting of c, so that $\tilde{T} = d\tilde{c} + [g(y_1), \tilde{c}]$ lifts T.

Up to substituting (x_1, y_1, e^{p_1}) with its gauge equivalent $(e^{\tilde{a}} * x_1, e^{\tilde{b}} * y_1, e^{g(\tilde{b})} e^{\tilde{T}} e^{p_1} e^{h(\tilde{a})})$, we can suppose that[c] $\beta(x_1, y_1, e^{p_1}) = \gamma(x_2, y_2, e^{p_2}) \in \widetilde{\mathrm{MC}}_{(h,g)}(A)$.

Then $((x_1, y_1, e^{p_1}), (x_2, y_2, e^{p_2})) \in \widetilde{\mathrm{MC}}_{(h,g)}(B) \times_{\widetilde{\mathrm{MC}}_{(h,g)}(A)} \widetilde{\mathrm{MC}}_{(h,g)}(C)$ and so, since $\widetilde{\mathrm{MC}}_{(h,g)}$ is a predeformation functor, there exists a lifting in $\widetilde{\mathrm{MC}}_{(h,g)}(B \times_A C)$. Next, it is sufficient to take its equivalence class in $\widetilde{\mathrm{Def}}_{(h,g)}$.

Finally, we prove that condition (III.2.3.3) is satisfied.

Let $\alpha : A \longrightarrow C$ be a surjection with $C \in \mathbf{C}_0$ an acyclic complex. Let (x_1, y_1, e^{p_1}) and $(x_2, y_2, e^{p_2}) \in \widetilde{\mathrm{Def}}_{(h,g)}(\ker \alpha)$ (in particular, $g(y_1) = e^{p_1} * h(x_1)$ and $g(y_2) = e^{p_2} * h(x_2)$) be gauge equivalent as elements in $\widetilde{\mathrm{MC}}_{(h,g)}(A)$, i.e., there exist $a \in (L \otimes A)^0$, $b \in (N \otimes A)^0$ and $c \in (M \otimes A)^{-1}$ such that

$$x_2 = e^a * x_1, \qquad y_2 = e^b * y_1,$$

$$e^{p_2} = e^{g(b)} e^T e^{p_1} e^{-h(a)}, \quad \text{with} \qquad T = dc + [g(y_1), c].$$

We are looking for $\overline{a} \in (L \otimes \ker \alpha)^0$, $\overline{b} \in (N \otimes \ker \alpha)^0$ and $\overline{c} \in (M \otimes \ker \alpha)^{-1}$ such that $\overline{T} = d\overline{c} + [g(y_1), \overline{c}]$ and

$$x_2 = e^{\overline{a}} * x_1 m \qquad y_2 = e^{\overline{b}} * y_1, \qquad e^{p_2} = e^{g(\overline{b})} e^{\overline{T}} e^{p_1} e^{-h(\overline{a})}.$$

---

[c] $\beta(e^{\tilde{a}} * x_1, e^{\tilde{b}} * y_1, e^{g(b)} e^{\tilde{T}} e^{p_1} e^{h(\tilde{a})}) = (e^{\beta(\tilde{a})} * \beta(x_1), e^{\beta(\tilde{b})} * \beta(y_1), e^{g(\beta(\tilde{b}))} e^{\beta(\tilde{T})} e^{\beta(p_1)} e^{h(\beta(\tilde{a}))}) = (e^a * \beta(x_1), e^b * \beta(y_1), e^{g(b)} e^T e^{\beta(p_1)} e^{-h(a)}) = (\gamma(x_2), \gamma(y_2), e^{\gamma(p_2)}) = \gamma(x_2, y_2, e^{p_2})$

Since $L\otimes C$ and $N\otimes C$ are abelian DGLAs and $\alpha(x_i)=\alpha(y_i)=0$, for $i=1,2$, we have

$$0=\alpha(x_2)=e^{\alpha(a)}*\alpha(x_1)=-d\alpha(a)$$

and

$$0=\alpha(y_2)=e^{\alpha(b)}*\alpha(y_1)=-d\alpha(y).$$

Moreover, $L\otimes C$ and $N\otimes C$ are acyclic; therefore, there exist $l\in(L\otimes A)^{-1}$ and $k\in(N\otimes A)^{-1}$ such that $d\alpha(l)=-\alpha(a)$ and $d\alpha(k)=-\alpha(b)$. This implies $dl+a\in(L\otimes\ker\alpha)^0$ and $dk+b\in(N\otimes\ker\alpha)^0$.

Set $w_1=dl+[x_1,l]\in Stab_A(x_1)$, $w_2=dk+[y_1,k]\in Stab_A(y_1)$ and define $\overline{a}=a\bullet w_1$ and $\overline{b}=b\bullet w_2$.

We claim that $\overline{a}\in(L\otimes\ker\alpha)^0$ and $\overline{b}\in(N\otimes\ker\alpha)^0$. Actually,

$$\overline{a}=a\bullet w_1\equiv a+w_1\equiv a+dl\ (mod\,[L\otimes A,L\otimes A]);$$

since $A\cdot A\subset\ker\alpha$, we conclude $\overline{a}\equiv a+dl\equiv 0\ (mod\,L\otimes\ker\alpha)$. An analogous calculation implies that $\overline{b}\in(N\otimes\ker\alpha)^0$.

Moreover, we note that

$$e^{\overline{a}}*x_1=e^ae^{w_1}*x_1=x_2\qquad\text{and}\qquad e^{\overline{b}}*y_1=e^be^{w_2}*y_1=y_2.$$

As to e^{p_1}, since $e^aStab_A(x)e^{-a}=Stab_A(y)$ with $y=e^a*x$, we have

$$e^{-g(w_2)}e^Te^{p_1}e^{h(w_1)}=e^Se^{p_1}$$

for some $S=df+[g(y_1),f]$ with $f\in(M\otimes A)^{-1}$.

Thus,

$$e^{p_2}=e^{g(b)}e^Te^{p_1}e^{-h(a)}=$$
$$e^{g(b)}e^{g(w_2)}e^{-g(w_2)}e^Te^{p_1}e^{h(w_1)}e^{-h(w_1)}e^{-h(a)}=e^{g(\overline{b})}e^Se^{p_1}e^{-h(\overline{a})}.$$

This implies that $e^S=e^{-g(\overline{b})}e^{p_2}e^{h(\overline{a})}e^{-p_1}$ lies in the subgroup $\exp((M\otimes\ker\alpha)^0)$ or equivalently $S=df+[g(y_1),f]\in(M\otimes\ker\alpha)^0$.

Since C is acyclic, the inclusion $M\otimes\ker\alpha\longrightarrow M\otimes A$ is a quasi-isomorphism and it remains a quasi-isomorphism if we consider the deformed differentials $d(-)+[g(y_1),-]$ on both $M\otimes\ker\alpha$ and $M\otimes A$ ($g(y_1)$ satisfies the Maurer-Cartan equation and so by Remark I.3.18 $d(-)+[g(y_1),-]$ is a differential).

Therefore, since the class of S is trivial in $M\otimes A$, it is also trivial in $M\otimes\ker\alpha$, i.e., there exists $\overline{c}\in(M\otimes\ker\alpha)^{-1}$ such that $S=\overline{T}=d\overline{c}+[g(y_1),\overline{c}]$.

In conclusion, $e^{p_2}=e^{g(\overline{b})}e^{\overline{T}}e^{p_1}e^{-h(\overline{a})}$ and so $\widetilde{\operatorname{Def}}_{(h,g)}$ is a predeformation functor.

Next, we prove that $\widetilde{\operatorname{Def}}_{(h,g)}$ is a deformation functor.

Let $C\in\mathbf{C}_0$, then

$$\widetilde{\operatorname{MC}}_{(h,g)}(C)=\{(l,n,e^m)\in(L\otimes C)^1\times(N\otimes C)^1\times\exp(M\otimes C)^0|$$
$$dl=dn=0,\ g(n)=e^m*h(l)=h(l)-dm\}=Z^1(\mathrm{C}_{(h,g)}\otimes C).$$

and

$$\widetilde{\mathrm{Def}}_{(h,g)}(C) = \widetilde{\mathrm{MC}}_{(h,g)}(C) \,/\, \{(-da, -db, dc+g(b)-h(a)) \mid a \in (L\otimes C)^0,\ b \in (N\otimes C)^0,\ c \in (M\otimes C)^{-1}\}. \tag{9}$$

Therefore, $\widetilde{\mathrm{Def}}_{(h,g)}(C)$ is isomorphic to the first cohomology group of the suspended cone of the pair of morphisms $h : L \otimes C \longrightarrow M \otimes C$ and $g : N \otimes C \longrightarrow M \otimes C$.

If C is also acyclic, then $\widetilde{\mathrm{Def}}_{(h,g)}(C) = 0$. This implies that $\widetilde{\mathrm{Def}}_{(h,g)}$ satisfies the condition of Definition III.2.4 and so it is a deformation functor.

Finally, equation (9) also implies that $T^i\widetilde{\mathrm{Def}}_{(h,g)} \cong H^i(C^{\cdot}_{(h,g)})$.

□

Proof of Lemma III.2.9. It suffices to apply the previous theorem with $M = N = 0$. □

Theorem III.2.27. $\widetilde{\mathrm{Def}}_{(h,g)} \cong \widetilde{\mathrm{MC}}^{+}_{(h,g)}$.

Proof. The projection onto the quotient $\widetilde{\mathrm{MC}}_{(h,g)} \longrightarrow \widetilde{\mathrm{Def}}_{(h,g)}$ induces, by Theorem III.2.15, a map $\widetilde{\mathrm{MC}}^{+}_{(h,g)} \longrightarrow \widetilde{\mathrm{Def}}_{(h,g)}$ that is a quasi-isomorphism by Proposition III.2.22 and Theorem III.2.26. □

Corollary III.2.28. *Let M be concentrated in non negative degree. Then for every $(A, m_A) \in$ **Art** we have $\widetilde{\mathrm{Def}}_{(h,g)}(m_A) = \mathrm{Def}_{(h,g)}(A)$.*

Proof. Obvious. □

Remark III.2.29. Every commutative diagram of differential graded Lie algebras

$$\begin{tikzcd}
& L \arrow[r, "\alpha'"] \arrow[d, "h"'] & P \arrow[d, "\eta"] \\
& M \arrow[r, "\alpha"] & Q \\
N \arrow[ur, "g"] \arrow[r, "\alpha''"] & R \arrow[ur, "\mu"'] &
\end{tikzcd}$$

induces a natural transformation $\widetilde{F}$ of the associated deformation functors:

$$\widetilde{F} : \widetilde{\mathrm{Def}}_{(h,g)} \longrightarrow \widetilde{\mathrm{Def}}_{(\eta,\mu)}.$$

Moreover, the inverse function Theorem III.2.14 implies that $\widetilde{F}$ is an isomorphism if and only if the map $(\alpha', \alpha, \alpha'')$ induces a quasi-isomorphism of complexes $\varphi^{\cdot} : \mathrm{C}^{\cdot}_{(h,g)} \longrightarrow \mathrm{C}^{\cdot}_{(\eta,\mu)}$.

Proof of Theorem III.1.23. It is sufficient to apply the previous remark and Corollary III.2.28.

□

Proof of Theorem III.1.24. It suffices to apply the inverse function Theorem III.2.14 and Corollary III.2.28.

□

III.2.5. Fibred product. In Example I.3.14, we have defined a DGLA structure on $M[t, dt] = M \otimes \mathbb{C}[t, dt]$ and for each $a \in \mathbb{C}$ an evaluation morphism e_a which is a surjective quasi-isomorphism:

$$e_a : M[t, dt] \longrightarrow M, \qquad e_a(\sum m_i t^i + n_i t^i dt) = \sum m_i a^i.$$

Define $K \subset L \times N \times M[t, dt] \times M[s, ds]$ as follows

$$K = \{(l, n, m_1(t, dt), m_2(s, ds)) |\, h(l) = e_1(m_2(s, ds)), g(n) = e_0(m_1(t, dt))\}.$$

K is a DGLA with bracket and differential δ defined as the natural ones on each component.

Define the following morphisms of DGLAs:

$$e_0 : K \longrightarrow M, \qquad (l, n, m_1(t, dt), m_2(s, ds)) \longmapsto e_0(m_1(t, dt))$$

and

$$e_1 : K \longrightarrow M, \qquad (l, n, m_1(t, dt), m_2(s, ds)) \longmapsto e_1(m_2(s, ds)).$$

Then we can construct the following simplicial diagram of DGLAs

$$\begin{tikzcd}
L \times N \arrow[r, "G"] \arrow[d, shift right, "h"'] \arrow[d, shift left, "g"] & K \subset L \times N \times M[t, dt] \times M[s, ds] \arrow[d, shift right, "e_1"'] \arrow[d, shift left, "e_0"] \\
M \arrow[r, "id"] & M
\end{tikzcd} \tag{10}$$

with:

$$\begin{tikzcd}
(l, n) \arrow[r, "G"] \arrow[d, "g"] \arrow[dd, "h"'] & (l, n, g(n), h(l)) \arrow[d, "e_0"] \arrow[dd, "e_1"'] \\
g(n) \arrow[r, "id"] & g(n) \\
h(l) \arrow[r, "id"] & h(l).
\end{tikzcd}$$

The diagram is commutative in a simplicial meaning and G is a quasi-isomorphism.

Lemma III.2.30. *The complexes* $C^{\cdot}_{e_1 - e_0}$ *and* $C^{\cdot}_{h-g}$ *are quasi-isomorphic.*

Proof. Consider the following commutative diagram of complexes

$$\begin{tikzcd}
0 \arrow[r] & (M^{\cdot -1}, -d) \arrow[r] \arrow[d, "id"] & C^{\cdot}_{h-g} \arrow[r] \arrow[d, "\alpha"] & (L^{\cdot} \oplus N^{\cdot}, d) \arrow[r] \arrow[d, "G"] & 0 \\
0 \arrow[r] & (M^{\cdot -1}, -d) \arrow[r] & C^{\cdot}_{e_1-e_0} \arrow[r] & (K^{\cdot}, \delta) \arrow[r] & 0,
\end{tikzcd}$$

with $\alpha(l,n,m) = (l,n,g(n),h(l),m)$, for $(l,n,m) \in C^{\cdot}_{h-g}$. Since id and G are quasi-isomorphism, α is a quasi-isomorphisms. $\square$

PROPOSITION III.2.31. $\widetilde{\mathrm{Def}}_{(h,g)} \cong \widetilde{\mathrm{Def}}_{(e_1,e_0)}$.

PROOF. Since G is a morphism of DGLAs, using diagram (10), we can define a morphism of Maurer-Cartan functors:

$$G' : \widetilde{\mathrm{MC}}_{(h,g)} \longrightarrow \widetilde{\mathrm{MC}}_{(e_1,e_0)},$$

$$(l,n,m) \longmapsto (G(l,n),m), \qquad \text{with } G(l,n) = (l,n,g(n),h(l)).$$

It is well-defined since $G(l,n) \in K$ satisfies the Maurer-Cartan equation and

$$e^m * e_1(G(l,n)) = e^m * e_1(h(l)) = e^m * h(l) = g(n) = e_0(g(n)) = e_0(G(l,n)).$$

Therefore, G' induces a morphism between the associated deformation functors $\widetilde{\mathrm{Def}}_{(h,g)}$ and $\widetilde{\mathrm{Def}}_{(e_1,e_0)}$ that is a quasi-isomorphism by Lemma III.2.30 (and so an isomorphism by Theorem III.2.14). $\square$

III.2.5.1. *Definition of* $\mathrm{H}_{(h,g)}$, *properties and barycentric subdivision.* Let $H \subset K$ be defined as follow

$$H = \{(l,n,m_1(t,dt),m_2(s,ds)) \in K |\ e_1(m_1(t,dt)) = e_0(m_2(s,ds))\},$$

or written in detail

$$H = \{(l,n,m_1(t,dt),m_2(s,ds)) \in L \times N \times M[t,dt] \times M[s,ds] \mid$$
$$h(l) = e_1(m_2(s,ds)),\ g(n) = e_0(m_1(t,dt)),\ e_1(m_1(t,dt)) = e_0(m_2(s,ds))\}.$$

Let $k = (l,n,m_1(t,dt),m_2(s,ds)) \in K$. Then the pair $m_1(t,dt)$ and $m_2(s,ds)$ have fixed values at one of the extremes of the unit interval. More precisely, the value of $m_1(t,dt)$ is fixed at the origin and $m_2(s,ds)$ is fixed at 1, i.e.,



If k also lies in H, then there are conditions on the other extremes: the value of $m_1(t,dt)$ at 1 has to coincide with the value of $m_2(s,ds)$ at 0.

Let

$$\mathrm{H}_{(h,g)} = \{(l,n,m(t,dt)) \in L \times N \times M[t,dt] \mid h(l) = e_1(m(t,dt)),\ g(n) = e_0(m(t,dt))\}. \tag{11}$$

Since e_i are morphisms of DGLAs, it is clear that $\mathrm{H}_{(h,g)}$ is a DGLA.

Moreover, considering the barycentric subdivision we get an injective quasi-isomorphism

$$\mathrm{H}_{(h,g)} \hookrightarrow H,$$
$$(l, n, m(t, dt)) \longmapsto (l, n, m\,(\frac{1}{2}\,t, dt), m\,(\frac{s+1}{2}, ds)).$$

Definition III.2.32. $\mathrm{H}_{(h,g)}$ is the *differential graded Lie algebra associated with the pair* (h, g).

Proposition III.2.33. $\mathrm{H}_{(h,g)}$ *is quasi-isomorphic to the complex* $C^{\cdot}_{e_1-e_0}$.

Proof. It is sufficient to consider the following commutative diagram of complexes

$$\begin{tikzcd}
\mathrm{H}_{(h,g)} \arrow[r, hook] \arrow[d] & K \arrow[d, "e_1-e_0"] \\
0 \arrow[r] & M,
\end{tikzcd}$$

where $e_1 - e_0$ is surjective. □

Proposition III.2.34. *Let* $h : L \longrightarrow M$ *and* $g : N \longrightarrow M$ *be morphisms of DGLAs. If the morphism* $g - h : N \times L \longrightarrow M$ *is surjective, then* $\widetilde{\mathrm{Def}}_{L\times_N M}$ *is isomorphic to* $\widetilde{\mathrm{Def}}_{(h,g)}$.

Proof. We recall that, by definition,

$$\widetilde{\mathrm{MC}}_{L\times_M N}(A) =$$
$$\{(l, n) \in (L\otimes A)^1\times(N\otimes A)^1 | dl+\frac{1}{2}[l, l] = 0,\ dn+\frac{1}{2}[n, n] = 0,\ h(l) = g(n)\},$$

and

$$\widetilde{\mathrm{MC}}_{(h,g)}(A) = \{(l, n, m) \in (L \otimes A)^1 \times (N \otimes A)^1 \times (M \otimes A)^0 |$$
$$dl + \frac{1}{2}[l, l] = 0,\ dn + \frac{1}{2}[n, n] = 0,\ g(n) = e^m * h(l)\}.$$

Moreover, $T^i\widetilde{\mathrm{Def}}_{L\times_M N} \cong H^i(L \times_M N)$ and $T^i\widetilde{\mathrm{Def}}_{(h,g)} \cong H^i(C^{\cdot}_{(h,g)})$.

Let

$$\psi : \widetilde{\mathrm{MC}}_{L\times_M N}(A) \longrightarrow \widetilde{\mathrm{MC}}_{(h,g)}(A),$$
$$(l, n) \longmapsto (l, n, 0),$$

and $\varphi : \widetilde{\mathrm{Def}}_{L\times_N M} \longrightarrow \widetilde{\mathrm{Def}}_{(h,g)}$ the induced map between the associated extended deformation functors. Then φ is a quasi-isomorphism.

For completeness we state all details, denoting by the same φ the map induced on cohomology.

φ is injective. Suppose that $\varphi([(l, n)]) = [(l, n, 0)] = 0$ in $H^i(C^{\cdot}_{(h,g)})$. Then, $[(l, n, 0)] = (dr, ds, -dt-g(s)+h(r))$ with $r \in (L)^{i-1}$, $s \in (N)^{i-1}$ and $t \in (M)^{i-2}$.

Since $g-h$ is surjective, there exist $p \in L^{i-2}$ and $q \in N^{i-2}$ such that $g(q)-h(p)=-t$ and so $g(dq)-h(dp)=-dt$. Let $(l',n')=(r-dp,s-dq)$; then $h(l')=h(r)-h(dp)=g(s)+dt-dt-g(dq)=g(n')$.

Therefore, $(l',n') \in L \times_M N$ and $d(l',n')=(dr,ds)=(l,n)$ and so $[(l,n)]=0 \in H^i(L \times_M N)$.

φ is surjective. Let $[(l,n,m)] \in H^i(C^{\cdot}_{(h,g)})$, i.e., $dl=dn=0$ and $-dm-g(n)+h(l)=0$. Since the class $[(l,n,m)]$ coincides with the class $[l+dr,n+ds,m-dt-g(s)+h(r)] \in H^i(C^{\cdot}_{(h,g)})$, we are looking for $r \in L^{i-1}$, $s \in N^{i-1}$ and $t \in M^{i-2}$ such that $(l+dr,n+ds) \in H^i(L \times_M N)$ and $m-dt-g(s)+h(r)=0$ (thus, $[(l,n,m)] \in \operatorname{Im}(\varphi)$).

Since $g-h$ is surjective, there exist $r \in L^{i-1}$ and $s \in N^{i-1}$, such that $g(s)-h(r)=m-dt$ and so $g(ds)-h(dr)=dm$. Therefore, $h(l+dr)=h(l)+g(ds)-dm=h(l)+g(ds)-h(l)+g(n)=g(n+ds)$, that is, $(l+dr,n+ds) \in L \times_M N$ and $\varphi([(l+dr,n+ds)])=[(l,n,m)]$. $\square$

Remark III.2.35. Let $\alpha_0,\alpha_1 : L \longrightarrow M$ be morphisms of DGLAs with $\alpha_0-\alpha_1$ surjective and $T=\{t \in L \mid \alpha_0(t)=\alpha_1(t)\}$ (T is called the equalizer of α_0 and α_1). In this particular case, the previous proposition implies $\widetilde{\operatorname{Def}}_T \cong \widetilde{\operatorname{Def}}_{(\alpha_1,\alpha_0)}$.

In conclusion, we have the following theorem.

Theorem III.2.36. $\widetilde{\operatorname{Def}}_{\mathrm{H}_{(h,g)}} \cong \widetilde{\operatorname{Def}}_{(h,g)}$.

Proof. It is sufficient to apply Proposition III.2.34 to $e_0,e_1 : K \longrightarrow M$ (with e_0-e_1 surjective) to conclude that $\widetilde{\operatorname{Def}}_{\mathrm{H}_{(h,g)}} \cong \widetilde{\operatorname{Def}}_{(e_1,e_0)}$; then Proposition III.2.31 is used to complete the proof. $\square$

CHAPTER IV

Deformations of holomorphic maps

This chapter is devoted to the main topic of this thesis: infinitesimal deformations of holomorphic maps of complex compact manifolds.

These deformations were first studied during the 70s by E. Horikawa in his works ([**14**] and [**15**]) and then by M. Namba [**27**] and Z. Ran [**28**].

Our purpose is to study these deformations using a technique based on differential graded Lie algebras.

Note. Unless otherwise specified, X *and* Y *are compact complex connected smooth varieties.*

IV.1. Deformation functor $\operatorname{Def}(f)$ of holomorphic maps

The main references for this section are [**14**] and [**27**, Section 3.6].

Definition IV.1.1. Let $f : X \longrightarrow Y$ be a holomorphic map and $A \in \mathbf{Art}$. An *infinitesimal deformation of* f *with fixed domain and codomain over* $\operatorname{Spec}(A)$ is a commutative diagram

$$\begin{tikzcd}
X \times S \arrow[rr, "\mathcal{F}"] \arrow[dr] & & Y \times S \arrow[dl] \\
& S, &
\end{tikzcd}$$

where $S = \operatorname{Spec}(A)$, the morphisms onto S are the projections, $\mathcal{F}$ is a holomorphic map and f coincides with the restriction of $\mathcal{F}$ to the fibers over the closed point of S.

If $A = \mathbb{K}[\varepsilon]$ we have a *first order deformation of* f *with fixed domain and codomain.*

Two infinitesimal deformations of f with fixed domain and codomain

$$\begin{tikzcd}
X \times S \arrow[rr, "\mathcal{F}"] \arrow[dr] & & Y \times S \arrow[dl] \\
& S &
\end{tikzcd} \quad \text{and} \quad \begin{tikzcd}
X \times S \arrow[rr, "\mathcal{F}'"] \arrow[dr] & & Y \times S \arrow[dl] \\
& S &
\end{tikzcd}$$

are equivalent if there exist automorphisms $\phi : X \longrightarrow X$ and $\psi : Y \longrightarrow Y$ such that the following diagram is commutative:

$$\begin{tikzcd}
X \times S \arrow[r, "\mathcal{F}"] \arrow[d, "{(\phi,id)}"'] & Y \times S \arrow[d, "{(\psi,id)}"] \\
X \times S \arrow[r, "\mathcal{F}'"] & Y \times S.
\end{tikzcd}$$

Definition IV.1.2. The *infinitesimal deformation functor* $\mathrm{Def}(X \xrightarrow{f} Y)$ of infinitesimal deformations of a holomorphic map f with fixed domain and codomain is defined as follows:

$$\mathrm{Def}(X \xrightarrow{f} Y) : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$A \longmapsto \mathrm{Def}(X \xrightarrow{f} Y)(A) = \left\{ \begin{array}{c} \text{isomorphism classes of} \\ \text{infinitesimal deformations of } f \\ \text{with fixed domain and codomain} \\ \text{over } \mathrm{Spec}(A) \end{array} \right\}.$$

Remark IV.1.3. When the domain and codomain are fixed, an infinitesimal deformation of a holomorphic map f can be viewed as an infinitesimal deformation of the graph of the map f in the product $X \times Y$, with $X \times Y$ fixed.

In this case, we are just deforming the map f. In general, we can also deform both the domain and the codomain.

Definition IV.1.4. Let $f : X \longrightarrow Y$ be a holomorphic map and $A \in \mathbf{Art}$. An *infinitesimal deformation of f over* $\mathrm{Spec}(A)$ is a commutative diagram of complex spaces

$$\begin{tikzcd}
X_A \arrow[rr, "\mathcal{F}"] \arrow[dr, "\pi"'] & & Y_A \arrow[dl, "\mu"] \\
& S, &
\end{tikzcd}$$

where $S = \mathrm{Spec}(A)$, (X_A, π, S) and (Y_A, μ, S) are infinitesimal deformations of X and Y, respectively (Definition I.2.1), $\mathcal{F}$ is a holomorphic map that restricted to the fibers over the closed point of S coincides with f.

If $A = \mathbb{K}[\varepsilon]$ we have a *first order deformation* of f.

Definition IV.1.5. Let

$$\begin{tikzcd}
X_A \arrow[rr, "\mathcal{F}"] \arrow[dr, "\pi"'] & & Y_A \arrow[dl, "\mu"] \\
& S &
\end{tikzcd} \quad \text{and} \quad \begin{tikzcd}
X'_A \arrow[rr, "\mathcal{F}'"] \arrow[dr, "\pi'"'] & & Y'_A \arrow[dl, "\mu'"] \\
& S &
\end{tikzcd}$$

be two infinitesimal deformations of f. They are *equivalent* if there exist bi-holomorphic maps $\phi : X_A \longrightarrow X'_A$ and $\psi : Y_A \longrightarrow Y'_A$ (that are equivalence of infinitesimal deformations of X and Y, respectively) such that the following diagram is commutative:

$$\begin{tikzcd}
X_A \arrow[r, "\mathcal{F}"] \arrow[d, "\phi"'] & Y_A \arrow[d, "\psi"] \\
X'_A \arrow[r, "\mathcal{F}'"] & Y'_A.
\end{tikzcd}$$

Definition IV.1.6. The *functor of infinitesimal deformations* of a holomorphic map $f : X \longrightarrow Y$ is

$$\mathrm{Def}(f) : \mathbf{Art} \longrightarrow \mathbf{Set},$$

$$A \longmapsto \mathrm{Def}(f)(A) = \left\{ \begin{array}{c} \text{isomorphism classes of} \\ \text{infinitesimal deformations of} \\ f \text{ over } \mathrm{Spec}(A) \end{array} \right\}.$$

Proposition IV.1.7. Def(f) *is a deformation functor, since it satisfies the conditions of Definition I.1.10.*

Proof. It follows from the fact that the functors Def_X and Def_Y of infinitesimal deformations of X and Y are deformation functors. ☐

Remark IV.1.8. In this general case, the infinitesimal deformations of f can be interpreted as infinitesimal deformations $\widetilde{\Gamma}$ of the graph Γ of the map f in the product $X \times Y$, such that the induced deformations $\widetilde{X \times Y}$ of $X \times Y$ are products of infinitesimal deformations of X and of Y. Since not all the deformations of a product are product of deformations (see Remark II.7.5), we are not just considering the deformations of the graph in the product.

Moreover, with this interpretation, two infinitesimal deformations $\widetilde{\Gamma} \subset \widetilde{X \times Y}$ and $\widetilde{\Gamma}' \subset \widetilde{X \times Y}'$ are equivalent if there exists an isomorphism $\phi : \widetilde{X \times Y} \longrightarrow \widetilde{X \times Y}'$ of infinitesimal deformations of $X \times Y$ such that $\phi(\widetilde{\Gamma}) = \widetilde{\Gamma}'$.

IV.1.1. Tangent and obstruction spaces of Def(f)**.** Let $\mathcal{U} = \{U_i\}$ and $\mathcal{W} = \{W_j\}$ be Stein open covers of X and Y, respectively, such that $f(U_i) \subset V_i$ for each i. For any integer $p \geq 0$, let

$$\check{C}^p(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y) = \check{C}^p(\mathcal{U}, \Theta_X) \oplus \check{C}^p(\mathcal{W}, \Theta_Y) \oplus \check{C}^{p-1}(\mathcal{U}, f^*\Theta_Y),$$

(with $\check{C}^{-1}(\mathcal{U}, f^*\Theta_Y) = 0$). Define a linear map

$$\check{D} : \check{C}^p(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y) \longrightarrow \check{C}^{p+1}(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y)$$

$$(x, y, z) \longmapsto (\check{\delta}x, \check{\delta}y, \check{\delta}z + (-1)^p(f_*x - f^*y)).$$

Using the equalities $f_*\check{\delta} = \check{\delta}f_*$ and $f^*\check{\delta} = \check{\delta}f^*$, we conclude that $\check{D}$ is a differential ($\check{D} \circ \check{D} = 0$). Therefore, the cohomology $\mathbb{C}$-vector spaces $\check{H}^p(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y)$ are well defined.

Lemma IV.1.9. $\check{H}^{\cdot}(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y)$ *does not depend on the choice of the covers and so we denote it by* $\check{H}^{\cdot}(\Theta_X, \Theta_Y, f^*\Theta_Y)$.

Proof. The following linear maps are well defined:

$$\check{H}^p(\mathcal{U},\Theta_X)\oplus\check{H}^p(\mathcal{V},\Theta_Y)\longrightarrow\check{H}^p(\mathcal{U},f^*\Theta_Y)$$

$$(\{n_1\},\{n_2\})\longmapsto\{f_*n_1-f^*n_2\};$$

$$\check{H}^p(\mathcal{U},f^*\Theta_Y)\longrightarrow\check{H}^{p+1}(\mathcal{U},\mathcal{W},\Theta_X,\Theta_Y,f^*\Theta_Y)$$

$$\{a\}\longmapsto\{(0,0,a)\};$$

$$H^p(\mathcal{U},\mathcal{W},\Theta_X,\Theta_Y,f^*\Theta_Y)\longrightarrow\check{H}^p(\mathcal{U},\Theta_X)\oplus\check{H}^p(\mathcal{V},\Theta_Y)$$

$$\{(n_1,n_2,a)\}\longmapsto(\{n_1\},\{n_2\}).$$

Then the sequence below is exact:

$$\cdots\longrightarrow\check{H}^p(\mathcal{U},\Theta_X)\oplus\check{H}^p(\mathcal{V},\Theta_Y)\longrightarrow\check{H}^p(\mathcal{U},f^*\Theta_Y)\longrightarrow$$

$$\longrightarrow\check{H}^{p+1}(\mathcal{U},\mathcal{W},\Theta_X,\Theta_Y,f^*\Theta_Y)\longrightarrow$$

$$\longrightarrow\check{H}^{p+1}(\mathcal{U},\Theta_X)\oplus\check{H}^{p+1}(\mathcal{V},\Theta_Y)\longrightarrow\check{H}^{p+1}(\mathcal{U},f^*\Theta_Y)\longrightarrow\cdots.$$

Moreover, $\check{H}^{\cdot}(\mathcal{U},\Theta_X)$, $\check{H}^{\cdot}(\mathcal{U},\Theta_X)$ and $\check{H}^{\cdot}(\mathcal{U},f^*\Theta_Y)$ does not depend on the choice of the Stein open covers $\mathcal{U}$ and $\mathcal{V}$. Hence applying the so called "five lemma", $\check{H}^{\cdot}(\mathcal{U},\mathcal{W},\Theta_X,\Theta_Y,f^*\Theta_Y)$ does not depend on the choice of the covers. ☐

In [**14**], E. Horikawa used the vector space $\check{H}^{\cdot}(\Theta_X,\Theta_Y,f^*\Theta_Y)$ to describe the tangent and obstruction spaces of the deformation functor Def(f).

Theorem IV.1.10. *$\check{H}^1(\Theta_X,\Theta_Y,f^*\Theta_Y)$ is in 1-1 correspondence with the first order deformations of $f:X\longrightarrow Y$.*

The obstruction space of the functor Def(f) *is naturally contained in $\check{H}^2(\Theta_X,\Theta_Y,f^*\Theta_Y)$.*

Proof. See [**27**, Section 3.6].

☐

Remark IV.1.11. Consider a first order deformation f_ε of f: in particular, we are considering first order deformations X_ε and Y_ε of X and Y. Using Theorem I.2.8, we associate with X_ε a class $x\in\check{H}^1(X,\Theta_X)$ and to Y_ε a class $y\in\check{H}^1(Y,\Theta_Y)$.

Therefore, the class in $\check{H}^1(\Theta_X,\Theta_Y,f^*\Theta_Y)$ associated with f_ε is $[(x,y,z)]$ with $z\in\check{C}^0(\mathcal{U},f^*\Theta_Y)$ such that $\check{\delta}z=f_*x-f^*y$.

Analogously, let $0\longrightarrow J\longrightarrow B\longrightarrow A\longrightarrow 0$ be a small extension and f_A an infinitesimal deformation of f over Spec(A). If $h\in\check{H}^2(X,\Theta_X)$ and $k\in\check{H}^2(Y,\Theta_Y)$ are the obstruction classes associated with X_A and Y_A, respectively, then the obstruction class in $\check{H}^2(\Theta_X,\Theta_Y,f^*\Theta_Y)$ associated with f_A is $[(h,k,r)]$, with $r\in\check{C}^1(\mathcal{U},f^*\Theta_Y)$ such that $\check{\delta}r=-(f_*h-f^*k)$.

Remark IV.1.12. Using Čech's cohomology, we have defined the $\mathbb{C}$-vector space $\check{H}^*(\Theta_X, \Theta_Y, f^*\Theta_Y)$. For convenience we reinterpret it, using Dolbeault's cohomology.

Let $(B^{\cdot}, D_{\overline{\partial}})$ be the complex below

$$B^p = A_X^{(0,p)}(\Theta_X) \oplus A_Y^{(0,p)}(\Theta_Y) \oplus A_X^{(0,p-1)}(f^*\Theta_Y)$$

and

$$D_{\overline{\partial}} : B^p \longrightarrow B^{p+1}, \qquad (x, y, z) \longmapsto (\overline{\partial}x, \overline{\partial}y, \overline{\partial}z + (-1)^p(f_*x - f^*y)).$$

Lemma IV.1.13. *The complexes* $(\check{C}^{\cdot}(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y), \check{D})$ *and* $(B^{\cdot}, D_{\overline{\partial}})$ *are quasi-isomorphic.*

Proof. Let $\mathcal{U} = \{U_i\}$ and $\mathcal{W} = \{W_j\}$ as above and denote by $\phi_1 : \check{C}^*(\mathcal{U}, \Theta_X) \longrightarrow A_X^{0,*}(\Theta_X)$, $\phi_2 : \check{C}^*(\mathcal{V}, \Theta_Y) \longrightarrow A_Y^{0,*}(\Theta_Y)$ and $\phi_3 : \check{C}^*(\mathcal{U}, f^*\Theta_X) \longrightarrow A_X^{0,*}(f^*\Theta_X)$ the quasi-isomorphisms of complexes of Leray's theorem, defined in Section II.3.1. We recall that $\phi_i\check{\delta} = \overline{\partial}\phi_i$, $f_*\phi_i = \phi_i f_*$ and $f^*\phi_i = \phi_i f^*$, for each $i = 1, 2, 3$ (see Section II.6).

Next, define the following morphism

$$\gamma : (\check{C}^{\cdot}(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y), \check{D}) \longrightarrow (B^{\cdot}, D_{\overline{\partial}}),$$

$$\gamma(x, y, z) = (\phi_1(x), \phi_2(y), \phi_3(z)).$$

Then γ is a morphism of complexes:; actually, for any $(x, y, z) \in \check{C}^p(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y)$

$$\begin{tikzcd}
(x, y, z) \arrow[r, "\gamma"] \arrow[d, "\check{D}"'] & (\phi_1(x), \phi_2(y), \phi_3(z)) \arrow[d, "D_{\overline{\partial}}"] \\
(\check{\delta}x, \check{\delta}y, \check{\delta}z + (-1)^p(f_*x - f^*y)) \arrow[r] & b
\end{tikzcd}$$

where $b = (\overline{\partial}\phi_1(x), \overline{\partial}\phi_2(y), \overline{\partial}\phi_3(z) + (-1)^p(f_*\phi_1(x) - f^*\phi_2(y)))$.

Moreover, we have the following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \check{C}^{\cdot-1}(\mathcal{U}, f^*\Theta_Y) \arrow[r] \arrow[d, "\phi_3"] & \check{\mathcal{C}}^{\cdot} \arrow[r] \arrow[d, "\gamma"] & \check{C}^{\cdot}(\mathcal{U}, \Theta_X) \oplus \check{C}^{\cdot}(\mathcal{W}, \Theta_Y) \arrow[r] \arrow[d, "(\phi_1, \phi_2)"] & 0 \\
0 \arrow[r] & A_X^{(0,\cdot-1)}(f^*\Theta_Y) \arrow[r] & B^{\cdot} \arrow[r] & A_X^{(0,\cdot)}(\Theta_X) \oplus A_Y^{(0,\cdot)}(\Theta_Y) \arrow[r] & 0
\end{tikzcd}$$

where $\check{\mathcal{C}}^{\cdot}$ stands for $\check{C}^{\cdot}(\mathcal{U}, \mathcal{W}, \Theta_X, \Theta_Y, f^*\Theta_Y)$.

Since ϕ_3 and (ϕ_1, ϕ_2) are quasi-isomorphisms, γ is a quasi-isomorphism.

□

IV.2. Infinitesimal deformations of holomorphic maps

Let $f : X \longrightarrow Y$ be a holomorphic map and Γ its graph in $Z := X \times Y$.

Let

$$F : X \longrightarrow \Gamma \subseteq Z := X \times Y,$$
$$x \longmapsto (x, f(x)),$$

and $p : Z \longrightarrow X$ and $q : Z \longrightarrow Y$ the natural projections.

Then we have the following commutative diagram:

$$\begin{tikzcd}
X \arrow[rr, "F"] \arrow[dr, "id"'] \arrow[drrr, "f"] & & Z \arrow[dl, "p"'] \arrow[dr, "q"] & \\
& X & & Y.
\end{tikzcd}$$

In particular, $F^* \circ p^* = id$ and $F^* \circ q^* = f^*$.

Since $\Theta_Z = p^*\Theta_X \oplus q^*\Theta_Y$, it follows that $F^*(\Theta_Z) = \Theta_X \oplus f^*\Theta_Y$.

Define the morphism $\gamma : \Theta_Z \longrightarrow f^*\Theta_Y$ as the product

$$\gamma : \Theta_Z \xrightarrow{F^*} \Theta_X \oplus f^*\Theta_Y \xrightarrow{(f^*,-id)} f^*\Theta_Y$$

and let π be the following surjective morphism:

$$\mathcal{A}^{0,j}_Z(\Theta_Z) \xrightarrow{\pi} \mathcal{A}^{0,j}_X(f^*\Theta_Y) \longrightarrow 0,$$

$$\pi(\omega\, u) = F^*(\omega)\gamma(u), \qquad \forall\ \omega \in \mathcal{A}^{0,j}_Z,\ u \in \Theta_Z.$$

Since each $u \in \Theta_Z$ can be written as $u = p^*v_1 + q^*v_2$, for some $v \in \Theta_X$ and $w \in \Theta_Y$, we also have

$$\pi(\omega u) = F^*(\omega)(f_*(v_1) - f^*(v_2)).$$

Since $F^*\overline{\partial} = \overline{\partial}F^*$, π is a morphism of complexes.

For the reader's convenience, we give an explicit description of the map π.

Let $\mathcal{U} = \{U_i\}_{i\in I}$ and $\mathcal{V} = \{V_i\}_{i\in I}$ be finite Stein open covers of X and Y, respectively, such that $f(U_i) \subset V_i$ (U_i is allowed to be empty). Moreover, let $z = (z_1, z_2, \dots, z_n)$ on U_i and $w = (w_1, w_2, \dots w_m)$ on V_i be local holomorphic coordinate systems for each $i \in I$. As in Section II.6, if $v_1 = \displaystyle\sum_{i=1}^{n} \varphi_i(z)\frac{\partial}{\partial z_i}$ and $v_2 = \displaystyle\sum_{h=1}^{m} \psi_h(w)\frac{\partial}{\partial w_h}$ then $f_*v_1 = \displaystyle\sum_{i=1}^{n} \varphi_i(z)\sum_{h=1}^{m}\frac{\partial f_h}{\partial z_i}\frac{\partial}{\partial w_h}$ and $f^*v_2 = \displaystyle\sum_{h=1}^{m} \psi_h(f(z))\frac{\partial}{\partial w_h}$. Let K and J be multi-indexes of length k and j, respectively, and fix $\omega = \Phi(z,w)d\overline{z}_K \wedge d\overline{w}_J \in \mathcal{A}^{0,k+j}_Z$. Then

$$\pi(\omega u) = F^*(\omega)(f_*(v_1) - f^*(v_2)) =$$

$$\Phi(z, f(z)) d\overline{z}_K \wedge \overline{\partial} f_J \sum_{h=1}^{m} \left(\sum_{i=1}^{n} \varphi_i(z) \frac{\partial f_h}{\partial z_i} - \psi_h(f(z)) \right) \frac{\partial}{\partial w_h}.$$

Let $\mathcal{L}$ be the kernel of π:

$$0 \longrightarrow \mathcal{L} \xrightarrow{h} \mathcal{A}_Z^{0,*}(\Theta_Z) \xrightarrow{\pi} \mathcal{A}_X^{0,*}(f^*\Theta_Y) \longrightarrow 0 \tag{12}$$

and $h : \mathcal{L} \longrightarrow \mathcal{A}_Z^{0,*}(\Theta_Z)$ the inclusion.

Lemma IV.2.1. *$\mathcal{L}$ is a sheaf of differential graded subalgebras of $\mathcal{A}_Z^{0,*}(\Theta_Z)$ and h is a morphism of differential graded Lie algebras.*

Proof. There is a canonical isomorphism between the normal bundle $N_{\Gamma|Z}$ of Γ in Z and the pull-back f^*T_Y.

Therefore, (12) reduces to

$$0 \longrightarrow \mathcal{L} \xrightarrow{h} \mathcal{A}_Z^{0,*}(\Theta_Z) \xrightarrow{\pi} \mathcal{A}_\Gamma^{0,*}(N_{\Gamma|Z}) \longrightarrow 0.$$

Then, by Lemma II.5.7, $\mathcal{L}$ is a sheaf of differential graded subalgebras of $A_Z^{0,*}(\Theta_Z)$.

□

Let L be the differential graded Lie algebra of global sections of $\mathcal{L}$.

Let M be the Kodaira-Spencer algebra of the product $X \times Y$: $M = KS_{X\times Y} = KS_Z$ and $h : L \longrightarrow M$ be the inclusion.

Let $N = KS_X \times KS_Y$ be the product of the Kodaira-Spencer algebras of X and of Y and $g : KS_X \times KS_Y \longrightarrow KS_{X\times Y}$ be given by $g = p^* + q^*$, i.e., $g(n_1, n_2) = p^*n_1 + q^*n_2$ (for $n = (n_1, n_2)$ we also use the notation $g(n)$).

Therefore, we get a diagram

$$\begin{tikzcd} & L \arrow[d, hook, "h"] \\ N = KS_X \times KS_Y \arrow[r, "g=(p^*,q^*)"] & M = KS_{X\times Y}. \end{tikzcd} \tag{13}$$

Remark IV.2.2. Given the morphisms of DGLAs $h : L \longrightarrow KS_{X\times Y}$ and $g : KS_X \times KS_Y \longrightarrow KX_{X\times Y}$ we can consider the complex $(\mathrm{C}^\cdot_{(h,g)}, D)$, with $C^i_{(h,g)} = L^i \oplus KS^i_X \oplus KS^i_Y \oplus KS^{i-1}_{X\times Y}$ and differential $D(l, n_1, n_2, m) = (\overline{\partial} l, \overline{\partial} n_1, \overline{\partial} n_2, -\overline{\partial} m - p^*n_1 - q^*n_2 + h(l))$.

Using the morphism $\pi : KS_{X\times Y} \longrightarrow A_X^{0,*}(f^*\Theta_Y)$ we can define a morphism

$$\beta : (\mathrm{C}^\cdot_{(h,g)}, D) \longrightarrow (B^\cdot, D_{\overline{\partial}}),$$

$$\beta(l, n_1, n_2, m) = (n_1, n_2, (-1)^i \pi(m)) \qquad \forall\ (l, n_1, n_2, m) \in \mathrm{C}^i_{(h,g)}.$$

Proposition IV.2.3. *$\beta : (\mathrm{C}^\cdot_{(h,g)}, D) \longrightarrow (B^\cdot, D_{\overline{\partial}})$ is a morphism of complexes which is a quasi-isomorphism.*

PROOF. β commutes with the differentials, i.e., $\beta \circ D = D_{\overline{\partial}} \circ \beta$; in fact, for each $(l, n_1, n_2, m) \in C^i_{(h,g)}$ we have the following commutative diagram

$$\begin{tikzcd}
(l, n_1, n_2, m) \arrow[r, "\beta"] \arrow[d, "D"'] & (n_1, n_2, (-1)^i\pi(m)) \arrow[d, "D_{\overline{\partial}}"] \\
c \arrow[r, "\beta"] & b,
\end{tikzcd}$$

where $c = (\overline{\partial} l, \overline{\partial} n_1, \overline{\partial} n_2, -\overline{\partial} m - p^*n_1 - q^*n_2 + h(l))$ and $b = \beta(c) = (\overline{\partial} n_1, \overline{\partial} n_2, (-1)^i(\overline{\partial}\pi(m) + f_*n_1 - f^*n_2)) = D_{\overline{\partial}}(n_1, n_2, (-1)^i\pi(m)))$.

Therefore, β induces a map in cohomology that we again call β that is a quasi-isomorphism. The proof is standard but we write it.

β is injective. Let $[(l, n_1, n_2, m)] \in H^i(C^{\cdot}_{(h,g)})$[a] be a class such that $\beta([(l, n_1, n_2, m)]) = [(n_1, n_2, (-1)^i\pi(m))]$ is zero in $H^i(B^{\cdot})$.

Then there exists $(a_1, a_2, b) \in B^{i-1}$ such that $n_1 = \overline{\partial} a_1$, $n_2 = \overline{\partial} a_2$, and $(-1)^i\pi(m) = \overline{\partial} b + (-1)^{i-1}(f_*a_1 - f^*a_2)$, so that $\pi(m) = (-1)^i\overline{\partial} b - f_*a_1 + f^*a_2$.

Let $n'_1 = a_1 \in KS^{i-1}_X$ and $n'_2 = a_2 \in KS^{i-1}_Y$; then $\overline{\partial} n'_1 = n_1$ and $\overline{\partial} n'_2 = n_2$.

Let $z \in KS^{i-2}_{X\times Y}$ be a lifting of $-b$ (i.e., $\pi(z) = -b$) and $l' = m + (-1)^i\overline{\partial} z + p^*n'_1 + q^*n'_2 \in KS_{X\times Y}i - 1$. Then $l' \in (L)^{i-1}$; indeed $\pi(l') = \pi(m) - (-1)^i\overline{\partial} b + f_*a_1 - f^*a_2 = 0$; moreover, $\overline{\partial} l' = \overline{\partial} m + p^*n_1 + q^*n_2 = l$. Therefore, $[(l, n_1, n_2, m)] = [\overline{\partial}(l', n'_1, n'_2, (-1)^i z)]$ is zero in $H^i(C^{\cdot}_{(h,g)})$.

β is surjective. Let $[(a_1, a_2, b)] \in H^i(B^{\cdot})$. Then $\overline{\partial} a_1 = \overline{\partial} a_2 = 0$ and $\overline{\partial} b + (-1)^i(f_*a_1 - f^*a_2) = 0$. Let $m \in KS^{i-1}_{X\times Y}$ be a lifting of $(-1)^i b$, i.e., $\pi(m) = (-1)^i b$. Let $n_1 = a_1 \in KS^i_X$ and $n_2 = a_2 \in KS^i_Y$. Finally, let $l = \overline{\partial} m + p^*a_1 + q^*a_2 \in KS^i_{X\times Y}$. Then $l \in L^i$; in fact $\pi(l) = \pi(\overline{\partial} m) + f_*a_1 - f^*a_2 = (-1)^i\overline{\partial} b + f_*a_1 - f^*a_2 = 0$.

Since $\overline{\partial} l = \overline{\partial} n_1 = \overline{\partial} n_2 = 0$ and $-\overline{\partial} m - p^*n_1 - q^*n_2 + l = 0$, $[(l, n_1, n_2, m)] \in H^i(C^{\cdot}_{(h,g)})$ and $\beta[(l, n_1, n_2, m)] = [(n_1, n_2, (-1)^i\pi(m))] = [(a_1, a_2, b)] \in H^i(B)$. ☐

IV.2.1. $\mathrm{Def}_{(h,g)}$ is isomorphic to $\mathrm{Def}(f)$. Using the notation above and diagram (13), consider the functor $\mathrm{Def}_{(h,g)}$. Since h is injective, by Remark III.1.16, for each $(A, m_A) \in \mathbf{Art}$ we have

$$\mathrm{Def}_{(h,g)}(A) = \{(n, e^m) \in (N^1 \otimes m_A) \times \exp(M^0 \otimes m_A) |$$
$$dn + \frac{1}{2}[n, n] = 0,\ e^{-m} * g(n) \in L^1 \otimes m_A\}/gauge.$$

REMARK IV.2.4. Let $(n, e^m) \in \mathrm{Def}_{(h,g)}$. In particular, $n = (n_1, n_2)$ satisfies the Maurer-Cartan equation and so $n_1 \in \mathrm{MC}_{KS_X}$ and $n_2 \in$

---

[a] In particular, this implies that $-\overline{\partial} m - p^*n_1 - q^*n_2 + l = 0$

MC_{KS_Y}. Therefore, there are associated with n infinitesimals deformations X_A of X (induced by n_1) and Y_A of Y (induced by n_2). Moreover, since $g(n)$ satisfies Maurer-Cartan equation in $M = KS_{X\times Y}$, it defines an infinitesimal deformation $(X\times Y)_A$ of $X\times Y$. By construction, the deformation $(X\times Y)_A$ is the product of the deformations X_A and Y_A.

Let $i^* : \mathcal{A}_Z^{0,*} \longrightarrow \mathcal{A}_\Gamma^{0,*}$ be the restriction morphism and let $I = \ker i^* \cap \mathcal{O}_Z$ be the holomorphic ideal sheaf of the graph $\Gamma$of f in Z.

Consider an infinitesimal deformation of the holomorphic map f over $\operatorname{Spec}(A)$ as an infinitesimal deformation $\widetilde{\Gamma}$ of Γ over $\operatorname{Spec}(A)$ and $\widetilde{Z}$ of Z over $\operatorname{Spec}(A)$, with $\widetilde{Z}$ product of deformations of X and Y over $\operatorname{Spec}(A)$.

By applying Remark IV.2.4 and Theorem II.7.3, the condition on the deformation $\widetilde{Z}$ is equivalent to requiring $\mathcal{O}_{\widetilde{Z}} = \mathcal{O}_A(g(n))$, for some Maurer-Cartan element $n \in KS_X \times KS_Y$.

The deformation $\widetilde{\Gamma}$ of the graph corresponds to an infinitesimal deformation $I_A \subset \mathcal{O}_{\widetilde{Z}}$ of the holomorphic ideal sheaf I over $\operatorname{Spec}(A)$, that is, I_A is a sheaf flat over A such that $I_A \otimes_A \mathbb{C} \cong I$.

In conclusion, to give an infinitesimal deformation of f over $\operatorname{Spec}(A)$ (an element in $\operatorname{Def}(f)(A)$) it is sufficient to give an ideal sheaf $I_A \subset \mathcal{O}_A(g(n))$ (for some $n \in \mathrm{MC}_{KS_X\times KS_Y}$) with I_A A-flat and $I_A \otimes_A \mathbb{C} \cong I$.

Theorem IV.2.5. *Let h, g and i^* be as above. Then there exists an isomorphism of functors*

$$\gamma : \operatorname{Def}_{(h,g)} \longrightarrow \operatorname{Def}(f).$$

Given a local Artinian $\mathbb{C}$-algebra A and an element $(n, e^m) \in \mathrm{MC}_{(h,g)}(A)$, we define a deformation of f over $\operatorname{Spec}(A)$ *as a deformation $I_A(n, e^m)$ of the holomorphic ideal sheaf of the graph of f in the following way*

$$\gamma(n, e^m) = I_A(n, e^m) := (\ker(\mathcal{A}_Z^{0,0}\otimes A \xrightarrow{\overline{\partial}+\boldsymbol{l}_{g(n)}} \mathcal{A}_Z^{0,1}\otimes A))\cap e^m(\ker i^*\otimes A) =$$

$$= \mathcal{O}_A(g(n)) \cap e^m(\ker i^* \otimes A),$$

where $\mathcal{O}_A(g(n))$ is the infinitesimal deformation of Z that corresponds to $g(n) \in MC_{KS_{X\times Y}}$ (see Theorem II.7.3).

Proof. For each $(n, e^m) \in \mathrm{MC}_{(h,g)}(A)$ we have defined

$$I_A(n, e^m) = \mathcal{O}_A(g(n)) \cap e^m(\ker i^* \otimes A).$$

First of all, we verify that this sheaf $I_A(n, e^m) \subset \mathcal{O}_A(g(n))$ defines an infinitesimal deformation of f; therefore, we need to prove that $I_A(n, e^m)$ is flat over A and $I_A \otimes_A \mathbb{C} \cong I$. It is equivalent to verify these properties for $e^{-m} I_A(n, e^m)$.

Applying Lemma II.5.5 yields

$$e^{-m}(\mathcal{O}_A(g(n))) = \ker(\overline{\partial} + e^{-m} * g(n) : \mathcal{A}_Z^{0,0} \otimes A \longrightarrow \mathcal{A}_Z^{0,1} \otimes A)$$

and also

$$e^{-m}I_A(n,e^m)=e^{-m}(\mathcal{O}_A(g(n)))\cap(\ker i^*\otimes A)=$$

$$=\ker(\overline{\partial}+e^{-m}*g(n))\cap(\ker i^*\otimes A).$$

Since flatness is a local property, we can assume that Z is a Stein manifold. Then $H^1(Z,\Theta_Z)=0$ and $H^0(Z,\Theta_Z)\longrightarrow H^0(Z,N_{\Gamma|Z})$ is surjective. Since the following sequence is exact

$$\cdots\longrightarrow H^0(Z,\Theta_Z)\longrightarrow H^0(Z,N_{\Gamma|Z})\longrightarrow H^1(Z,L)\longrightarrow H^1(Z,\Theta_Z)\longrightarrow\cdots,$$

we conclude that $H^1(L)=0$ or, equivalently, that the tangent space of the functor Def_L is trivial. Therefore, by Corollary I.1.21, Def_L is the trivial functor.

This implies the existence of $\nu\in L^0\otimes m_A$ such that $e^{-m}*g(n)=e^{\nu}*0$ (by hypothesis, $e^{-m}*g(n)$ is a solution of the Maurer-Cartan equation in L). Moreover, we recall that if $a\in L^0\otimes m_A$ then $e^a(\ker i^*\otimes A)=\ker i^*\otimes A$ (see Section II.5.1).

Therefore,

$$e^{-m}I_A(n,e^m)=\ker(\overline{\partial}+e^{\nu}*0)\cap(\ker i^*\otimes A)=\mathcal{O}_A(e^{\nu}*0)\cap(\ker i^*\otimes A)$$

$$=e^{\nu}(\mathcal{O}_A(0))\cap e^{\nu}(\ker i^*\otimes A)=e^{\nu}(I\otimes A).$$

Thus, $I_A(n,e^m)$ defines a deformation of f and the morphism

$$\gamma:\mathrm{MC}_{(h,g)}\longrightarrow\operatorname{Def}(f)$$

is well defined, such that

$$\gamma(A):\mathrm{MC}_{(h,g)}(A)\longrightarrow\operatorname{Def}(f)(A)$$

$$(n,e^m)\longmapsto\gamma(n,e^m)=I_A(n,e^m).$$

Moreover, γ is well defined on $\operatorname{Def}_{(h,g)}(A)=\mathrm{MC}_{(h,g)}(A)/gauge$. Actually, for each $a\in L^0\otimes m_A$ and $b\in N^0\otimes m_A$, we have

$$\gamma(e^b*n,e^{g(b)}e^me^a)=\mathcal{O}_A(e^{g(b)}*g(n))\cap e^{g(b)}e^me^a(\ker i^*\otimes A)=$$

$$e^{g(b)}\mathcal{O}_A(g(n))\cap e^{g(b)}e^m(\ker i^*\otimes A)=e^{g(b)}\gamma(n,e^m).$$

This implies that the deformations $\gamma(n,e^m)$ and $\gamma(e^b*n,e^{g(b)}e^me^a)$ are isomorphic (see Remark IV.1.8).

In conclusion, $\gamma:\operatorname{Def}_{(f,g)}\longrightarrow\operatorname{Def}(f)$ is a well defined natural transformation of functors.

In order to prove that γ is an isomorphism it is sufficient to prove that

i) γ is injective;
ii) γ induces a bijective map on the tangent spaces;
iii) γ induces an injective map on the obstruction spaces.

Actually, by Corollary I.1.32, conditions *ii)* and *iii)* imply that γ is étale and so surjective.

i) γ *is injective*. Suppose that $\gamma(n, e^m) = \gamma(r, e^s)$, i.e., the deformations induced by (n, e^m) and (r, e^s), respectively, are isomorphic. We want to conclude that (n, e^m) is gauge equivalent to (r, e^s), i.e., there exist $a \in L^0 \otimes m_A$ and $b \in N^0 \otimes m_A$ such that $e^b * r = n$ and $e^{g(b)}e^s e^a = e^m$.

By hypothesis, $\gamma(n, e^m)$ and $\gamma(r, e^s)$ are isomorphic deformations; then, in particular, the deformations induced on Z are isomorphic. This implies that there exists $b \in N^0 \otimes m_A$ such that $e^b * r = n$ and so $e^{g(b)}(\mathcal{O}_A(g(r))) = \mathcal{O}_A(g(n))$. Up to substituting (r, e^s) with its equivalent $(e^b * r, e^{g(b)}e^s)$, we can assume to be in the following situation

$$\mathcal{O}_A(g(n)) \cap e^m(\ker i^* \otimes A) = \mathcal{O}_A(g(n)) \cap e^{m'}(\ker i^* \otimes A).$$

Let $e^a = e^{-m'}e^m$, then

$$e^a(e^{-m}(\mathcal{O}_A(g(n))) \cap (\ker i^* \otimes A)) = e^{-m'}(\mathcal{O}_A(g(n))) \cap (\ker i^* \otimes A).$$

In particular, $e^a(e^{-m}(\mathcal{O}_A(g(n))) \cap (\ker i^* \otimes A)) \subseteq \ker i^* \otimes A$.

Next, we prove, by induction, that $a \in L^0 \otimes m_A$ (thus $e^m = e^{m'}e^a = e^{g(b)}e^s e^a$).

Let $z_1, \ldots, z_n$ be holomorphic coordinates on Z such that $Z \supset \Gamma = \{z_{t+1} = \cdots = z_n = 0\}$. Consider the projection on the residue field

$$e^{-m}(\mathcal{O}_A(g(n))) \cap (\ker i^* \otimes A) \longrightarrow \mathcal{O}_Z \cap \ker i^*.$$

Then $z_i \in \ker i^* \cap \mathcal{O}_Z$, for $i > t$. Since $e^{-m}(\mathcal{O}_A(g(n))) \cap (\ker i^* \otimes A)$ is flat over A, we can lift z_i to $\tilde{z}_i = z_i + \varphi_i \in e^{-m}(\mathcal{O}_A(g(n))) \cap (\ker i^* \otimes A)$, with $\varphi_i \in \ker i^* \otimes m_A$. By hypothesis,

$$e^a(\tilde{z}_i) = e^a(z_i) + e^a(\varphi_i) \in \ker i^* \otimes A. \tag{14}$$

By Lemma II.5.7, in order to prove that $a \in L^0 \otimes m_A$ it is sufficient to verify that $e^a(z_i) \in \ker i^* \otimes A$ and so, by (14), that $e^a(\varphi_i) \in \ker i^* \otimes A$.

If $A = \mathbb{C}[\varepsilon]$, then $\varphi_i \in \ker i^* \otimes \mathbb{C}\varepsilon$, $a \in \mathcal{A}_Z^{0,0} \otimes \mathbb{C}\varepsilon$, this implies $e^a(\varphi_i) = \varphi_i \in \ker i^* \otimes \mathbb{C}\varepsilon$.

Next, let $0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$ be a small extension. By hypothesis, $\alpha(a) \in L^0 \otimes m_A$, that is, $\alpha(a) = \displaystyle\sum_{j=1}^{n} \overline{a}_j \frac{\partial}{\partial z_j}$ with $\overline{a}_j \in \ker i^* \otimes m_A$ for $j > t$.

Let a'_j be liftings of $\overline{a}_j$. Then $a'_j \in \ker i^* \otimes m_B$ for $j > t$, $a' = \displaystyle\sum_{j=1}^{n} a'_j \frac{\partial}{\partial z_j} \in L^0 \otimes m_B$ and $e^{a'}(\varphi_i) \in \ker i^* \otimes m_B$. Since $\alpha(a) = \alpha(a')$, then $a = a' + j$ with $j \in M^0 \otimes J$. This implies that $e^a(\varphi_i) = e^{a'+j}(\varphi_i) = e^{a'}(\varphi_i) \in \ker i^* \otimes m_B$.

As to tangent and obstruction spaces, by Theorem IV.1.10 and Lemma IV.1.13, the tangent space of $\mathrm{Def}(f)$ is isomorphic to $H^1(B^\cdot)$ and the obstruction space is naturally contained in $H^2(B^\cdot)$, where $(B^\cdot, D_{\overline{\partial}})$ is the complex with $B^p = A_X^{(0,p)}(\Theta_X) \oplus A_Y^{(0,p)}(\Theta_Y) \oplus A_X^{(0,p-1)}(f^*\Theta_Y)$ and $D_{\overline{\partial}}(x,y,z) = (\overline{\partial}x, \overline{\partial}y, \overline{\partial}z + (-1)^p(f_*x - f^*y))$, for each $(x,y,z) \in B^p$. As to the functor $\mathrm{Def}_{(h,g)}$, in Section III.1.3, we have proved that the tangent space is $H^1(\mathrm{C}^\cdot_{(h,g)})$ and the obstruction space is naturally contained in $H^2(\mathrm{C}^\cdot_{(h,g)})$, where $\mathrm{C}^\cdot_{(h,g)}$ is the suspended cone associated with the pair (h,g) (Section III.1.1). Moreover, Proposition IV.2.3 shows the existence of a quasi-isomorphism β between the previous complexes $B^\cdot$ and $\mathrm{C}^\cdot_{(h,g)}$.

Next, we want to prove that the maps induced by γ on tangent and obstruction spaces coincide with the isomorphism induced by β.

ii) γ induces a bijection on the tangent spaces (we prove that γ coincides with the isomorphism β).

Let $(n_1, n_2, m) \in \mathrm{Def}_{(h,g)}(\mathbb{C}[\varepsilon])$. Then $\overline{\partial}n_1 = \overline{\partial}n_2 = 0$, $\overline{\partial}m + g(n) = \overline{\partial}m + p^*n_1 + q^*n_2 \in H^1(L)$ and so (n_1,n_2,m) determines the class $[(g(n) + \overline{\partial}m, n_1, n_2, m)] \in H^1(C^\cdot_{(h,g)})$. Moreover, we note that $\overline{\partial}\pi(m) + f_*n_1 - f^*n_2 = 0$ and $\beta[(g(n) + \overline{\partial}m, n_1, n_2, m)] = [(n_1, n_2, -\pi(m))]$.

Next, consider the first order deformation of f induced by (n_1,n_2,m), i.e., $\gamma(n_1,n_2,m) = \mathcal{O}_A(g(n)) \cap e^m(\ker i^* \otimes A)$. Using Theorem IV.1.10 and Remark IV.1.11, we associate with $\gamma(n_1,n_2,m)$ the class [b] $[(n_1,n_2,a)] \in H^1(B^\cdot)$, such that $\overline{\partial}a = f_*n_1 - f^*n_2$. Then $\overline{\partial}a = -\overline{\partial}\pi(m)$ and so $[(n_1,n_2,a)] = [(n_1,n_2,-\pi(m))] = \beta([(g(n)+\overline{\partial}m, n_1, n_2, m)]) \in H^1(B^\cdot)$.

This implies that the map induced by γ on the tangent spaces coincides with the isomorphism β.

iii) γ' induces an injective map on the obstruction spaces. In this case too, we prove that the map induced by γ on the obstruction space coincides with β and so it is injective.

Actually, let

$$0 \longrightarrow J \longrightarrow B \xrightarrow{\alpha} A \longrightarrow 0$$

be a small extension. The obstruction class associated with $(n_1,n_2,m) \in \mathrm{Def}_{(h,g)}(A)$ is the class $[(k,h_1,h_2,r)] \in H^2(\mathrm{C}^\cdot_{(h,g)}) \otimes J$ defined in Lemma III.1.19. We note that $\overline{\partial}r + p^*h_1 + q^*h_2 \in H^2(L))$ and so $\pi(\overline{\partial}r) = -f_*h_1 + f^*h_2$.

Next, again by Theorem IV.1.10 and Remark IV.1.11, the obstruction class associated with the deformation $\gamma(n_1,n_2,m)$ of f induced by

---

[b] By Theorem II.7.3, the class associated with the first order deformation of X induced by n_1 is n_1 itself and the class associated with the first order deformation of Y induced by n_2 is n_2 itself.

(n_1, n_2, m) is $[(h_1, h_2, a)] \in H^2(B^{\cdot}) \otimes J$ with $\overline{\partial} a = -(f_* h_1 - f^* h_2)$ and as above $[(h_1, h_2, a)] = \beta([(k, h_1, h_2, r)]) \in H^2(B^{\cdot}) \otimes J$.

$\square$

In conclusion, by suitably choosing L, M, and h, g, Theorem IV.2.5 shows that the infinitesimal deformation functor $\mathrm{Def}(f)$ associated with a holomorphic map f is isomorphic to the functor $\mathrm{Def}_{(h,g)}$.

Theorem IV.2.6. *Let $f : X \longrightarrow Y$ be a holomorphic map. Then the DGLA $\mathrm{H}_{(h,g)}$ associated with the above morphisms $h : L \hookrightarrow KS_{X \times Y}$ and $g = (p^*, q^*) : KS_X \times KS_Y \longrightarrow KS_{X \times Y}$ (see Definition III.2.32) controls infinitesimal deformations of f:*

$$\mathrm{Def}_{\mathrm{H}_{(h,g)}} \cong \mathrm{Def}(f).$$

Proof. It is sufficient to apply Theorem IV.2.5, Corollary III.2.28 and Theorem III.2.36. $\square$

Remark IV.2.7. This theorem is very interesting from the point of view of "guiding principle", since it shows the existence of a DGLA which controls the geometric problem of infinitesimal deformations of holomorphic maps.

Anyway, in Chapter V we will see that in the applications it is more convenient to use the functor $\mathrm{Def}_{(h,g)}$ than $\mathrm{Def}_{\mathrm{H}_{(h,g)}}$.

Remark IV.2.8. Consider the diagram

$$\begin{tikzcd}
 & L \arrow[d, "h"] & \\
KS_X \times KS_Y \arrow[r, "g"] \arrow[rrd, bend right=10, "\pi \circ g"'] & KS_{X \times Y} \arrow[rd, "\pi"] & \\
 & & A_X^{0,*}(f^* T_Y).
\end{tikzcd}$$

Since h is injective, Lemma III.1.6 implies the existence of a quasi-isomorphism of complexes $(\mathrm{C}^{\cdot}_{(h,g)}, D)$ and $(C^{\cdot}_{\pi \circ g}, \check{\delta})$.

Therefore, we get the following exact sequence

(15)

$$\cdots \longrightarrow H^1(C^{\cdot}_{\pi\circ g}) \xrightarrow{\varrho^1} H^1(X, \Theta_X) \oplus H^1(Y, \Theta_Y) \longrightarrow H^1(X, f^*\Theta_Y) \longrightarrow$$

$$\longrightarrow H^2(C^{\cdot}_{\pi\circ g}) \xrightarrow{\varrho^2} H^2(X, \Theta_X) \oplus H^2(Y, \Theta_Y) \longrightarrow H^2(X, f^*\Theta_Y) \longrightarrow \cdots$$

where ϱ^1 and ϱ^2 are the projections on the first factors and they are induced by the projection morphism $\varrho : \mathrm{Def}(f) \longrightarrow \mathrm{Def}_{KS_X \times KS_Y}$ (see Remark III.1.15).

In particular, $\varrho : \mathrm{Def}(f) \longrightarrow \mathrm{Def}_{KS_X \times KS_Y}$ associates with an infinitesimal deformation of f the induced infinitesimal deformations of X and Y.

Then ϱ^1 associates with a first order deformation of f the induced first order deformations of X and Y and ϱ^2 is a morphism of obstruction theory: the obstruction to deform f is mapped to the induced obstructions to deform X and Y (see also Remark IV.2.4).

In [**28**], Z. Ran studied the infinitesimal deformations of a holomorphic map $f : X \longrightarrow Y$ of singular compact complex spaces. He introduced some algebraic objects T^i_f, $i = 1, 2$, that classify the deformations of a map f and obtained the following exact sequence

(16)

$$T^1_f \longrightarrow \operatorname{Ext}^1_{\mathcal{O}_X}(\Omega_X, \mathcal{O}_X) \oplus \operatorname{Ext}^1_{\mathcal{O}_Y}(\Omega_Y, \mathcal{O}_Y) \longrightarrow \operatorname{Ext}^1_{\mathcal{O}_X}(f^*\Omega_Y, \mathcal{O}_X) \longrightarrow$$

$$\longrightarrow T^2_f \longrightarrow \operatorname{Ext}^2_{\mathcal{O}_X}(\Omega_X, \mathcal{O}_X) \oplus \operatorname{Ext}^2_{\mathcal{O}_Y}(\Omega_Y, \mathcal{O}_Y) \longrightarrow \operatorname{Ext}^2_{\mathcal{O}_X}(f^*\Omega_Y, \mathcal{O}_X).$$

LEMMA IV.2.9. *If X and Y are compact complex manifold, then the exact sequence (16) reduces to (15).*

PROOF. If X and Y are smooth, then Ω_X and Ω_Y are locally free. Then applying the spectral sequence associated with Ext (see [**9**, Lemme 7.4.1]) we get $\operatorname{Ext}^i_{\mathcal{O}_X}(\Omega_X, \mathcal{O}_X) \cong H^i(X, \Theta_X)$, $\operatorname{Ext}^i_{\mathcal{O}_Y}(\Omega_Y, \mathcal{O}_Y) \cong H^i(Y, \Theta_Y)$ and $\operatorname{Ext}^i_{\mathcal{O}_X}(f^*\Omega_Y, \mathcal{O}_X) \cong H^i(X, f^*\Theta_Y)$.

□

CHAPTER V

Semiregularity maps

In the previous chapter we have studied the infinitesimal deformations of holomorphic maps.

More precisely, let $f : X \longrightarrow Y$ be a holomorphic map of compact complex manifolds and $\Gamma \subset X \times Y$ its graph.

Consider the DGLAs $M = KS_{X\times Y}$, $N = KS_X \times KS_Y$ and the morphism $g = (p^*, q^*) : KS_X \times KS_Y \longrightarrow KS_{X\times Y}$, where p and q are the projections of $X \times Y$ onto X and Y, respectively.

Moreover, let $L = A^{0,*}_{X\times Y}(-log\,\Gamma)$ be the DGLA defined by the following exact sequence (see Section IV.2):

$$0 \longrightarrow L \xrightarrow{h} KS_{X\times Y} \xrightarrow{\pi} A^{0,*}_X(f^*\Theta_Y) \longrightarrow 0.$$

Then we get the following diagram

(17)
$$\begin{tikzcd}
 & L \arrow[d, "h"] \\
KS_X \times KS_Y \arrow[r, "(p^*,q^*)"] & KS_{X\times Y}.
\end{tikzcd}$$

Theorem IV.2.5 shows the existence of an isomorphism between the functor $\mathrm{Def}(f)$ of infinitesimal deformations of f and the functor $\mathrm{Def}_{(h,g)}$ associated with the pair of morphisms (h, g):

$$\mathrm{Def}(f) \cong \mathrm{Def}_{(h,g)} \qquad \text{(Theorem IV.2.5)}.$$

Furthermore, Theorem IV.2.6 gives an explicit description of the DGLA $H_{(h,g)}$ (Definition III.2.32) that controls the infinitesimal deformations of f:

$$\mathrm{Def}(f) \cong \mathrm{Def}_{H(h,g)} \qquad \text{(Theorem IV.2.6)}.$$

In particular, if $i : X \hookrightarrow Y$ is an inclusion, we can find an easy description of the DGLA associated with $\mathrm{Def}(i)$ (see Section V.4). Actually, let L' be the the DGLA L' introduced in Section II.5.1 (see also [**24**, Section 5])

$$0 \longrightarrow L' \longrightarrow A^{0,*}_Y(\Theta_Y) \xrightarrow{\pi'} A^{0,*}_X(N_{X|Y}) \longrightarrow 0.$$

Thus, L' controls the deformations of i (Corollary V.4.1).

In general, without an easy description of $H_{(h,g)}$ it is convenient to use the deformation functor associated with the previous diagram (17).

For example, if we want to study the infinitesimal deformations of f with fixed domain or fixed codomain, it is sufficient to consider diagram (17) with $N = KS_Y$ (Section V.1) or $N = KS_X$ (Section V.2), respectively.

Anyway, the main application of the techniques developed in the previous chapters concerns the study of obstructions to deforming a holomorphic map f and the "semiregularity" maps.

In general, we can find a vector space V (most of the times a cohomology vector space) that contains the obstruction space, but we do not know an explicit description of the elements which are obstructions.

Then the idea is to restrict the vector space V, for example by defining a map on V, the so called "semiregularity" map, that contains the obstructions in the kernel.

In particular, let $f : X \longrightarrow Y$ be a holomorphic map and consider the infinitesimal deformations of f with fixed codomain Y. In [**14**] E. Horikawa proved the following theorem.

THEOREM V.0.10. *(Horikawa) Let $f : X \longrightarrow Y$ be a holomorphic map and consider the functor of infinitesimal deformations of f with fixed codomain Y. Then the tangent space is the hypercohomology vector space $\mathbb{H}^1\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right)$ and the obstruction space is contained in $\mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right)$.*

Using the techniques introduced in the previous section, we can improve this result in the case of Kähler manifolds, defining a map that contains the obstruction space in the kernel.

THEOREM. *Let $f : X \longrightarrow Y$ be a holomorphic map of compact Kähler manifolds. Let $p = dimY - dimX$. Then the obstruction space of deformations of f (with fixed Y) is contained in the kernel of a map*

$$\sigma : \mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

The previous map is the generalization of the semiregularity map defined by Bloch (see [**3**] or [**24**, Section 9]) obtained when f is the inclusion map $X \hookrightarrow Y$, i.e.,

$$\sigma : H^1(X, N_{X|Y}) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

The proof of this theorem is postponed to Section V.1.2 (Corollary V.1.5), where we also give an explicit description of the map σ.

V.1. Deformations with fixed codomain

This section is devoted to studying infinitesimal deformations of a holomorphic map $f : X \longrightarrow Y$, with fixed codomain Y.

In this case the DGLA N reduces to KS_X and so diagram (17) reduces to

$$\begin{tikzcd}
 & L \arrow[d, "h"] & \\
A_X^{0,*}(\Theta_X) \arrow[r, "p^*"] \arrow[rrd, bend right=10, "f_*"'] & A_{X\times Y}^{0,*}(\Theta_{X\times Y}) \arrow[rd, "\pi"] & \\
 & & A_X^{0,*}(f^*\Theta_Y).
\end{tikzcd}$$

where f_* is the product $\pi \circ p^*$ (see Section IV.2).

Using this diagram and Theorem IV.2.5, we can easily prove Theorem V.0.10 due to E. Horikawa.

Proposition V.1.1. *The tangent space of the infinitesimal deformation functor of a holomorphic map $f : X \longrightarrow Y$, with Y fixed, is $H^1(C^{\cdot}_{f_*})$ and the obstruction space is naturally contained in $H^2(C^{\cdot}_{f_*})$.*

Proof. Theorem IV.2.5 implies that the infinitesimal deformation functor of f, with Y fixed, is isomorphic to $\mathrm{Def}_{(h,p^*)}$. Therefore, the tangent space is $H^1(C^{\cdot}_{(h,p^*)})$ and the obstruction space is naturally contained in $H^2(C^{\cdot}_{(h,p^*)})$. Since h is injective, Lemma III.1.6 implies that, for each i, $H^i(C^{\cdot}_{(h,p^*)}) \cong H^i(C^{\cdot}_{\pi\circ p^*}) = H^i(C^{\cdot}_{f_*})$. □

As we already announced we improve this theorem in the case of Kähler manifolds (see Section V.1.2).

The next section is devoted to some preliminary lemmas.

V.1.1. Preliminaries.

Let Z be a complex manifold.

Then $KS_Z = A_Z^{0,*}(\Theta_Z)$ is the Kodaira-Spencer algebra of Z and in Section II.5 we have defined the contraction map $\boldsymbol{i}$:

$$\boldsymbol{i} : KS_Z \longrightarrow \mathrm{Hom}^*(A_Z, A_Z),$$

$$\boldsymbol{i}_a(\omega) = a \lrcorner \omega$$

for any $a \in KS_Z$ and $\omega \in A_Z^{*,*}$.

Therefore, $\boldsymbol{i}(A_Z^{0,j}(\Theta_Z)) \subset \oplus_{h,l} \mathrm{Hom}^0(A_Z^{h,l}, A_Z^{h-1,l+j}) \subset \mathrm{Hom}^{j-1}(A_Z, A_Z)$.

In order to interpret $\boldsymbol{i}$ as a morphism of DGLAs, the key idea, due to M. Manetti [**24**, Section 8], is to substitute $\mathrm{Hom}^*(A_Z, A_Z)$ with the graded vector space $\mathrm{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right) = \bigoplus_i \mathrm{Hom}^{i-1}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right)$ (see Example I.3.4). Consider on $\mathrm{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right)$ the following differential δ and bracket $\{\ ,\ \}$:

$$\delta(f) = -\overline{\partial} f - (-1)^{\deg(f)} f \overline{\partial},$$

$$\{f, g\} = f\partial g - (-1)^{\deg(f)\deg(g)} g\partial f.$$

Proposition V.1.2. $\operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right)$ *is a DGLA and the linear map*

$$\boldsymbol{i} : A_Z^{0,*}(T_Z) \longrightarrow \operatorname{Htp}\left(\ker(\partial), \frac{A_Z}{\partial A_Z}\right)$$

is a morphism of DGLAs.

Proof. See [**24**, Proposition 8.1]. An easy calculation shows that the graded vector space $\operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right)$ is a DGLA.

Moreover, $\boldsymbol{i}$ is a linear map that preserves degree and commutes with differential and bracket. Actually, using Cartan fomulas[a] (see Lemma II.5.1), we get

$$\boldsymbol{i}_{\tilde{d}a} = -[\overline{\partial}, \boldsymbol{i}_a] = -\overline{\partial}\boldsymbol{i}_a + (-1)^{\overline{a}-1}\boldsymbol{i}_a\overline{\partial}$$

and, by definition of δ,

$$\delta(\boldsymbol{i}_a) = -\overline{\partial}\boldsymbol{i}_a - (-1)^{\overline{a}}\boldsymbol{i}_a\overline{\partial}.$$

Therefore, $\boldsymbol{i}_{\tilde{d}a} = \delta(\boldsymbol{i}_a)$. As to the bracket, again by Cartan fomulas, we have

$$\boldsymbol{i}_{[a,b]} = [\boldsymbol{i}_a, [\partial, \boldsymbol{i}_b]] = [\boldsymbol{i}_a, \partial\boldsymbol{i}_b - (-1)^{\overline{b}-1}\boldsymbol{i}_b\partial] =$$
$$\boldsymbol{i}_a\partial\boldsymbol{i}_b - (-1)^{\overline{b}-1}\boldsymbol{i}_a\boldsymbol{i}_b\partial - (-1)^{(\overline{a}-1)\overline{b}}(\partial\boldsymbol{i}_b\boldsymbol{i}_a - (-1)^{\overline{b}-1}\boldsymbol{i}_b\partial\boldsymbol{i}_a) =^{\text{b}}$$
$$\boldsymbol{i}_a\partial\boldsymbol{i}_b + (-1)^{\overline{a}\overline{b}-1}\boldsymbol{i}_b\partial\boldsymbol{i}_a$$

and, by definition of { , },

$$\{\boldsymbol{i}_a, \boldsymbol{i}_b\} = \boldsymbol{i}_a\partial\boldsymbol{i}_b - (-1)^{\overline{a}\,\overline{b}}\boldsymbol{i}_b\partial\boldsymbol{i}_a.$$

□

Next, let $f : X \longrightarrow Y$ be a holomorphic map, fix $Z = X \times Y$ and Γ the graph of f in Z. Let $I_\Gamma \subset A_Z$ be the space of the differential forms vanishing on Γ and $L \subset KS_Z$ be the DGLA defined as in Lemma IV.2.1:

$$0 \longrightarrow L \longrightarrow KS_Z \longrightarrow A_X^{0,*}(f^*\Theta_Y) \longrightarrow 0.$$

We recall that

$$L \subset \{a \in A_Z^{0,*}(\Theta_Z) |\ \boldsymbol{i}_a(I_\Gamma) \subset I_\Gamma\};$$

moreover,

$$p^*A_X^{0,*}(\Theta_X) \subset \{a \in A_Z^{0,*}(\Theta_Z) |\ \boldsymbol{i}_a(q^*A_Y) = 0\},$$

where p and q are the projection of Z onto X and Y, respectively.

In conclusion, we can define the following commutative diagram of morphisms of DGLAs:

---

[a] $\boldsymbol{i}_{\tilde{d}a} = -[\overline{\partial}, \boldsymbol{i}_a],\ \boldsymbol{i}_{[a,b]} = [\boldsymbol{i}_a, [\partial, \boldsymbol{i}_b]] = [[\boldsymbol{i}_a, \partial], \boldsymbol{i}_b],\ [\boldsymbol{i}_a, \boldsymbol{i}_b] = 0.$

[b] $\boldsymbol{i}_a\boldsymbol{i}_b\partial$ and $\partial\boldsymbol{i}_b\boldsymbol{i}_a$ are zero in $\operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right)$

(18)

$$\begin{tikzcd}
L \arrow[r] \arrow[d, hook, "h"] & K = \left\{ f \in \operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right) \mid f(I_\Gamma \cap \ker(\partial)) \subset \dfrac{I_\Gamma}{I_\Gamma \cap \partial A_Z} \right\} \arrow[d, hook, "\eta"] \\
A_Z^{0,*}(\Theta_Z) \arrow[r] & \operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right) \\
A_X^{0,*}(\Theta_X) \arrow[r] \arrow[u, "p^*"] & J = \left\{ f \in \operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right) \mid f(\ker(\partial) \cap q^* A_Y) = 0 \right\}, \arrow[u, hook, "\mu"']
\end{tikzcd}$$

where the horizontal morphisms are all given by $\boldsymbol{i}$.

Therefore, diagram (18) induces a morphism of deformation functors:

$$\mathcal{I} : \operatorname{Def}_{(h,p^*)} \longrightarrow \operatorname{Def}_{(\eta,\mu)} .$$

LEMMA V.1.3. *If the differential graded vector spaces* $(\partial A_Z, \overline{\partial})$, $(\partial A_\Gamma, \overline{\partial})$ *and* $(\partial A_Z \cap q^* A_Y, \overline{\partial})$ *are acyclic, then the functor* $\operatorname{Def}_{(\eta,\mu)}$ *is unobstructed. In particular, the obstruction space of* $\operatorname{Def}_{(h,p^*)}$ *is naturally contained in the kernel of the map*

$$H^2(C^{\cdot}_{(h,p^*)}) \xrightarrow{\mathcal{I}} H^2(C^{\cdot}_{(\eta,\mu)}).$$

PROOF. This proof is an extension of the proof of [**24**, Lemma 8.2].

The projection $\ker(\partial) \to \ker(\partial)/\partial A_Z$ induces a commutative diagram

(19)

$$\begin{tikzcd}
K \arrow[d, "\eta"] & \{f \in K | f(\partial A_Z) = 0\} \arrow[l, "\alpha"'] \arrow[d, "\eta'"'] \\
\operatorname{Htp}\left(\ker(\partial), \dfrac{A_Z}{\partial A_Z}\right) & \operatorname{Htp}\left(\dfrac{\ker(\partial)}{\partial A_Z}, \dfrac{A_Z}{\partial A_Z}\right) \arrow[l, "\beta"'] \\
J \arrow[u, "\mu"] & \{f \in J | f(\partial A_Z) = 0\}. \arrow[l, "\gamma"'] \arrow[u, "\mu'"']
\end{tikzcd}$$

Since ∂A_Z is acyclic, β is a quasi-isomorphism of DGLAs. Since

$$\operatorname{coker}(\alpha) = \{f \in \operatorname{Htp}\left(\partial A_Z, \frac{A_Z}{\partial A_Z}\right) | f(I_\Gamma \cap \partial A_Z) \subset \frac{I_\Gamma}{I_\Gamma \cap \partial A_Z}\},$$

there exists an exact sequence

$$0 \to \operatorname{Htp}\left(\frac{\partial A_Z}{I_\Gamma \cap \partial A_Z}, \frac{A_Z}{\partial A_Z}\right) \to \operatorname{coker}(\alpha) \to \operatorname{Htp}\left(I_\Gamma \cap \partial A_Z, \frac{I_\Gamma}{I_\Gamma \cap \partial A_Z}\right) \to 0.$$

Moreover, the exact sequence

$$0 \longrightarrow I_\Gamma \cap A_Z \longrightarrow \partial A_Z \longrightarrow \partial A_\Gamma \longrightarrow 0$$

implies that $I_\Gamma \cap A_Z$ and $\dfrac{\partial A_Z}{I_\Gamma \cap \partial A_Z} = \partial A_\Gamma$ are acyclic. By Example I.3.3, this implies that the complexes $\operatorname{Htp}\left(\dfrac{\partial A_Z}{I_\Gamma \cap \partial A_Z}, \dfrac{A_Z}{\partial A_Z}\right)$ and $\operatorname{Htp}\left(I_\Gamma \cap \partial A_Z, \dfrac{I_\Gamma}{I_\Gamma \cap \partial A_Z}\right)$ are acyclic and so the same holds for $\operatorname{coker}(\alpha)$; i.e., α is a quasi-isomorphism.

As to γ, we have

$$\operatorname{coker}(\gamma) =$$
$$\left\{f \in \operatorname{Htp}\left(\partial A_Z, \frac{A_Z}{\partial A_Z}\right) \mid f(\partial A_Z \cap q^* A_Y) = 0\right\} =$$
$$\operatorname{Htp}\left(\frac{\partial A_Z}{\partial A_Z \cap q^* A_Y}, \frac{A_Z}{\partial A_Z}\right).$$

By hypothesis, $\partial A_Z \cap q^* A_Y$ and ∂A_Z are acyclic and so the same holds for $\dfrac{\partial A_Z}{\partial A_Z \cap q^* A_Y}$. Then $\operatorname{coker}(\gamma)$ is acyclic, i.e., γ is also a quasi-isomorphism.

Therefore, Theorem III.1.23 implies the existence of an isomorphism of functors $\operatorname{Def}_{(\eta,\mu)} \cong \operatorname{Def}_{(\eta',\mu')}$.

We note that the elements of the three algebras of the second column of (19) vanish on ∂A_Z. Then, by definition of the bracket $\{\ ,\ \}$, these algebras are abelian. Therefore, Lemma III.1.20 implies that the functor $\operatorname{Def}_{(\eta,\mu)} \cong \operatorname{Def}_{(\eta',\mu')}$ is smooth. Finally, Proposition III.1.22 guarantees that the obstruction space lies in the kernel of $H^2(C^\cdot_{(h,p^*)}) \xrightarrow{\mathcal{I}} H^2(C^\cdot_{(\eta,\mu)})$. □

V.1.2. Semiregularity for deformations with fixed codomain.

With the notation of the previous section we have the following theorem.

THEOREM V.1.4. *Let $f : X \longrightarrow Y$ be a holomorphic map of compact Kähler manifolds. Then the obstruction space to the infinitesimal deformations of f with fixed codomain is contained in the kernel of the following map*

$$H^2(C^\cdot_{f_*}) \xrightarrow{\mathcal{I}'} H^1\left(\operatorname{Htp}(I_\Gamma \cap \ker(\partial) \cap q^* A_Y, A_\Gamma)\right).$$

PROOF. Lemma II.2.2 implies that the complexes $(\partial A_Z, \overline{\partial})$, $(\partial A_\Gamma, \overline{\partial})$ and $(\partial A_Z \cap q^* A_Y, \overline{\partial})$ are acyclic. Then we can apply Lemma V.1.3 to conclude that the obstruction space lies in the kernel of the following map

$$H^2(C^\cdot_{(h,p^*)}) \xrightarrow{\mathcal{I}} H^2(C^\cdot_{(\eta,\mu)}).$$

Since h is injective, by Lemma III.1.6, $H^2(C^{\cdot}_{(h,p^*)}) \cong H^2(C^{\cdot}_{\pi\circ p^*}) \cong H^2(C^{\cdot}_{f_*})$. Then the obstructions lie in the kernel of $\mathcal{I} : H^2(C^{\cdot}_{f_*}) \longrightarrow H^2(C^{\cdot}_{(\eta,\mu)})$.

As to $H^2(C^{\cdot}_{(\eta,\mu)})$, consider

$$K = \left\{ f \in \mathrm{Htp}\left(\ker(\partial), \frac{A_Z}{\partial A_Z}\right) \mid f(I_\Gamma \cap \ker(\partial)) \subset \frac{I_\Gamma}{I_\Gamma \cap \partial A_Z} \right\}$$

and the exact sequence

$$0 \longrightarrow K \xrightarrow{\eta} \mathrm{Htp}\left(\ker(\partial), \frac{A_Z}{\partial A_Z}\right) \xrightarrow{\pi'} \mathrm{coker}(\eta) \longrightarrow 0,$$

with $\mathrm{coker}(\eta) = \mathrm{Htp}\left(I_\Gamma \cap \ker(\partial), \dfrac{A_\Gamma}{\partial A_\Gamma}\right)$.

Applying again Lemma III.1.6, there exists an isomorphism $H^2(C^{\cdot}_{(\eta,\mu)}) \cong H^2(C^{\cdot}_{\pi'\circ\mu})$. By Remark III.1.1 there also exists a map $\mathcal{I}'' : H^2(C^{\cdot}_{\pi'\circ\mu}) \longrightarrow H^1(\mathrm{coker}(\pi' \circ \mu))$.

More precisely,

$$J = \left\{ f \in \mathrm{Htp}\left(\ker(\partial), \frac{A_Z}{\partial A_Z}\right) \mid f(\ker(\partial) \cap q^* A_Y) = 0 \right\},$$

$$\pi' \circ \mu : J \longrightarrow \mathrm{Htp}\left(I_\Gamma \cap \ker(\partial), \frac{A_\Gamma}{\partial A_\Gamma}\right)$$

and

$$\mathrm{coker}(\pi' \circ \mu) = \mathrm{Htp}(I_\Gamma \cap \ker(\partial) \cap q^* A_Y, \frac{A_\Gamma}{\partial A_\Gamma}).$$

Finally, since the complex ∂A_Γ is acyclic, the projection

$$H^1(\mathrm{Htp}(I_\Gamma \cap \ker(\partial) \cap q^* A_Y, \frac{A_\Gamma}{\partial A_\Gamma}) \longrightarrow H^1(\mathrm{Htp}(I_\Gamma \cap \ker(\partial) \cap q^* A_Y, A_\Gamma))$$

is an isomorphism.

Therefore, the obstruction space is contained in the kernel of $\mathcal{I}' : H^2(C^{\cdot}_{f_*}) \longrightarrow H^1(\mathrm{Htp}(I_\Gamma \cap \ker(\partial) \cap q^* A_Y, A_\Gamma))$, i.e.,

$$\begin{tikzcd}
H^2(C^{\cdot}_{f_*}) \arrow[r, "\mathcal{I}"] \arrow[drr, "\mathcal{I}'"] & H^2(C^{\cdot}_{\pi'\circ\mu}) \arrow[r, "\mathcal{I}''"] & H^1\left(\mathrm{Htp}(I_\Gamma \cap \ker(\partial) \cap q^* A_Y, \frac{A_\Gamma}{\partial A_\Gamma})\right) \arrow[d, "\cong"] \\
 & & \oplus_i \mathrm{Hom}(H^i(I_\Gamma \cap \ker(\partial) \cap q^* A_Y), H^i(A_\Gamma)).
\end{tikzcd}$$

□

Corollary V.1.5. *Let $f : X \longrightarrow Y$ be a holomorphic map of compact Kähler manifolds. Let $p = dimY - dimX$. Then the obstruction* *space to the infinitesimal deformations of f with fixed Y is contained in the kernel of the map*

$$\sigma : \mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

PROOF. Let $n = dimX$, $p = dimY - dimX$ and $\mathcal{H}$ be the space of harmonic forms on Y of type $(n+1, n-1)$. By Dolbeault's theorem and Serre's duality we obtain the equalities $\mathcal{H}^\nu = (H^{n-1}(Y, \Omega_Y^{n+1}))^\nu = H^{p+1}(Y, \Omega_Y^{p-1})$.

Let $\omega \in \mathcal{H}$ such that $f^*\omega = 0$. By considering the contraction with ω we define a morphism of complexes

$$(A_X^{0,*}(f^*\Theta_Y), \overline{\partial}) \xrightarrow{\lrcorner\omega} (A_X^{n,*+n-1}, \overline{\partial}),$$

$$\lrcorner\omega(\phi f^*\chi) = \phi f^*(\chi\lrcorner\omega) \in A_X^{n,p+n-1} \qquad \forall\ \phi f^*\chi \in A_X^{0,p}(f^*\Theta_Y).$$

Actually, since $\overline{\partial}\omega = 0$, then $\overline{\partial}(\phi f^*(\chi\lrcorner\omega)) = (\overline{\partial}\phi) f^*(\chi\lrcorner\omega) = \lrcorner\omega(\overline{\partial}\phi f^*\chi)$.

In particular, since $f^*\omega = 0$, using the identity of Lemma II.6.1[c], we have the following commutative diagram

$$\begin{tikzcd}
A_X^{0,*}(f^*\Theta_Y) \arrow[r, "\lrcorner\omega"] & A_X^{n,*+n-1} \\
A_X^{0,*}(\Theta_X) \arrow[r] \arrow[u, "f_*"] & 0. \arrow[u, "\alpha"']
\end{tikzcd}$$

Then we get a morphism between the second cohomology groups of the cones associated with the morphisms f_* and α:

$$H^2(C^{\cdot}_{f_*}) \longrightarrow H^2(C^{\cdot}_{\alpha}) \cong H^n(X, \Omega_X^n).$$

By taking the product of the previous morphism with the integration on X, we get

$$\sigma : \mathbb{H}^2\left(X, \mathcal{O}(\Theta_X) \xrightarrow{f_*} \mathcal{O}(f^*\Theta_Y)\right) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

Since $q^*\mathcal{H}$ is contained in $I_\Gamma \cap \ker\overline{\partial} \cap \ker\partial \cap q^*A_Y$, we conclude the proof applying Theorem V.1.4.

$\square$

REMARK V.1.6. We recall that, as already observed in Remark II.2.3, the Kähler hypothesis is just used to have the $\partial\overline{\partial}$-lemma on A_X,A_Y, $A_{X\times Y}$ and A_Γ.

---

[c] $f^*(\chi\lrcorner\omega) = \eta\lrcorner f^*\omega$, for each $\omega \in A_Y^{*,*}$, $\chi \in A_Y^{0,*}(\Theta_Y)$ and $\eta \in A_X^{0,*}(\Theta_X)$ such that $f^*\chi = f_*\eta$.

V.2. Deformations of a map with fixed domain

In this section we study infinitesimal deformations of a holomorphic map $f : X \longrightarrow Y$, with fixed domain X.

In this case the DGLA N reduces to KS_Y and so diagram (17) reduces to

$$\begin{tikzcd}
& L \arrow[d, "h"] & \\
A_Y^{0,*}(\Theta_Y) \arrow[r, "q^*"] \arrow[rrd, bend right=10, "-f^*"'] & A_{X\times Y}^{0,*}(\Theta_{X\times Y}) \arrow[rd, "\pi"] & \\
& & A_X^{0,*}(f^*\Theta_Y).
\end{tikzcd}$$

where $-f^*$ is the product $\pi \circ q^*$.

Using this diagram and Theorem IV.2.5, analogously to the case of fixed codomain we can prove the following proposition.

Proposition V.2.1. *The tangent space of the infinitesimal deformation functor of a holomorphic map $f : X \longrightarrow Y$, with X fixed, is $H^1(C^{\cdot}_{f^*})$ and the obstruction space is naturally contained in $H^2(C^{\cdot}_{f^*})$.*

Proof. Analogous to the proof of Proposition V.1.1. □

V.3. Deformations with fixed domain and codomain

Let $f : X \longrightarrow Y$ be a holomorphic map and consider the infinitesimal deformations of f with fixed codomain and domain (see Definition IV.1.1).

As we have already observed in Remark IV.1.3, these deformations can be interpreted as infinitesimal deformations of the graph Γ in $X\times Y$, with $X \times Y$ fixed.

Therefore, in diagram (17), we do not need to consider the DGLA $N = KS_X \times KS_Y$ and g. Thus, the functor $\mathrm{Def}_{(h,g)}$ reduces to Def_h with

$$h : L = A_{X\times Y}^{0,*}(-log\,\Gamma) \hookrightarrow M = KS_{X\times Y}.$$

This implies that Theorem IV.2.5 reduces to:

Corollary V.3.1. *Let $f : X \longrightarrow Y$ be a holomorphic map. Then the functor* $\mathrm{Def}(X \xrightarrow{f} Y)$ *of infinitesimal deformations of f with fixed codomain and domain is isomorphic to* Def_h*:*

$$Def_h \cong \mathrm{Def}(X \xrightarrow{f} Y).$$

Proof. Apply Theorem IV.2.5 with $N = g = 0$. □

Remark V.3.2. This corollary is equivalent to [**24**, Theorem 5.2].

V.3.1. Semiregularity map. As to the obstructions to deform a map f fixing both X and Y, Lemma V.1.3 has the form below.

LEMMA V.3.3. *If the differential graded vector spaces* $(\partial A_Z, \overline{\partial})$ *and* $(\partial A_\Gamma, \overline{\partial})$ *are acyclic, then the functor* Def_η *is unobstructed. In particular, the obstruction space of* Def_h *is naturally contained in the kernel of the map*

$$H^2(C_h^\cdot) \xrightarrow{\mathcal{I}} H^2(C_\eta^\cdot) \cong \oplus_i \operatorname{Hom}\left(H^i(I_\Gamma \cap \ker(\partial)), H^i(\frac{A_Z}{\partial A_Z})\right).$$

PROOF. See [**24**, Lemma 8.2]. It follows from Lemma V.1.3, with $N = J = 0$.

□

We note that $H^2(C_h^\cdot) \cong H^1(X, f^*\Theta_Y)$ (see Remark III.1.1). Moreover, we have an analogue of Theorem V.1.4 and then Corollary V.1.5 becomes:

COROLLARY V.3.4. *Let* $f : X \longrightarrow Y$ *be a holomorphic map of compact Kähler manifolds. Let* $p = dimY - dimX$*. Then the obstruction space to the infinitesimal deformations of* f*, with fixed* X *and* Y*, is contained in the kernel of the map*

$$\sigma : H^1(X, f^*\Theta_Y) \longrightarrow H^{p+1}(Y, \Omega_Y^{p-1}).$$

PROOF. See [**24**, Corollary 9.2]. It follows from Lemma V.3.3 and Corollary V.1.5.

□

V.4. Deformations of the inclusion map

In this section, we focus our attention on the infinitesimal deformations of an inclusion $i : X \hookrightarrow Y$ of compact complex manifolds. The DGLAs L, M, N and the morphisms g, h and π are as before.

Consider the DGLA L' introduced in Section II.5.1 (see also [**24**, Section 5])

$$0 \longrightarrow L' \longrightarrow A_Y^{0,*}(\Theta_Y) \xrightarrow{\pi'} A_X^{0,*}(N_{X|Y}) \longrightarrow 0.$$

COROLLARY V.4.1. L' *controls the infinitesimal deformations of the inclusion* $i : X \hookrightarrow Y$*.*

The proof is postponed to the end of this section after two preliminary lemmas.

LEMMA V.4.2. *If* $i : X \hookrightarrow Y$ *is the inclusion, then the morphism* $g - h : L \times N \longrightarrow M$ *is surjective.*

PROOF. We want to prove that for each $\phi \in M = A_{X\times Y}^{0,*}(\Theta_{X\times Y})$ there exist $n_1 \in A_X^{0,*}(\Theta_X)$ and $n_2 \in A_Y^{0,*}(\Theta_Y)$ such that $g(n_1, n_2) - \phi = p^*n_1 + q^*n_2 - \phi \in L = \ker \pi$, that is, $\pi(\phi) = i_*n_1 - i^*n_2 \in A_X^{0,*}(i^*\Theta_Y) =$

$A_X^{0,*}(\Theta_{Y|X})$. Then the proof immediately follows from the fact that the restriction morphism $i^* : A_Y^{0,*}(\Theta_Y) \longrightarrow A_X^{0,*}(\Theta_{Y|X})$ is surjective. $\square$

Lemma V.4.3. *With the notation above, $L \times_M N \cong L'$.*

Proof. By definition, $L\times_M N = \{(l, n_1, n_2) \in L\times KS_X\times KS_Y \mid h(l) = p^*n_1 + q^*n_2\}$ and so $0 = \pi h(l) = i_*n_1 - i^*n_2$.

Define $\gamma : L \times_M N \longrightarrow KS_Y$ as the projection onto KS_Y.

Then γ is an injective morphism of DGLAs with L' as image.

Actually, suppose that $\gamma(l, n_1, n_2) = n_2 = 0$; then $i_*n_1 = 0$ and so $n_1 = n_2 = l = 0$.

As to the image, consider the following exact sequences

$$\begin{tikzcd}
0 \arrow[r] & L' \arrow[r] \arrow[d] & A_Y^{0,*}(\Theta_Y) \arrow[r, "\pi'"] \arrow[d, "i^*"] & A_X^{0,*}(N_{X|Y}) \arrow[r] \arrow[d, "\cong"'] & 0 \\
0 \arrow[r] & A_X^{0,*}(\Theta_X) \arrow[r, "i_*"] & A_Y^{0,*}(\Theta_{Y|X}) \arrow[r, "\beta"] & A_X^{0,*}(N_{X|Y}) \arrow[r] & 0.
\end{tikzcd}$$

Let $\gamma(l, n_1, n_2) = n_2 \in KS_Y$; then $\beta(i^*n_2) = \beta(i_*n_1) = 0$ and so $\pi'(n_2) = 0$. This implies that $\gamma(l, n_1, n_2) \in L'$. $\square$

Proof of Corollary V.4.1. By Theorem IV.2.5, the infinitesimal deformation functor $\mathrm{Def}(i)$ is isomorphic to the functor $\mathrm{Def}_{(h,g)}$. By Lemma V.4.2 and Proposition III.2.34, $\mathrm{Def}_{L\times_M N} \cong \mathrm{Def}_{(h,g)}$. Finally, Lemma V.4.3 implies that $\mathrm{Def}_{L'} \cong \mathrm{Def}_{L\times_M N}$. Then $\mathrm{Def}(i) \cong \mathrm{Def}_{L'}$. $\square$

V.4.1. Example. We can generalize the case of one inclusion $i : X \hookrightarrow Y$ considering more subvarieties.

For example, let X be a manifold of dimension n and $D_1, \ldots, D_m$ smooth hypersurfaces, with $0 < m \leq n-2$. Moreover, assume that $D_1, \ldots, D_m$ intersect transversally in a smooth subvariety S.

Define

$$\mathrm{Def}_{X;D_1,\ldots,D_m} : \mathbf{Art} \longrightarrow \mathbf{Set}$$

as the functor of infinitesimal deformations of the holomorphic map

$$f : \overset{\circ}{\bigcup} D_i \longrightarrow X,$$

where $f_{|D_i}$ is the inclusion. Equivalently, for any $A \in \mathbf{Art}$, $\mathrm{Def}_{X;D_1,\ldots,D_m}(A)$ provides an infinitesimal deformation X_A of X over $\mathrm{Spec}(A)$ and an infinitesimal deformations $\mathcal{D}_i \subset X_A$ of the D_i's.

Let $\Theta_X(-log\, D) \subset \Theta_X$ be the subsheaf of vector fields that are tangent to D_i, for every i, and $N_{D_i|X}$ be the normal bundle of D_i in X.

Define $L' := A_X^{0,*}(\Theta_X(-log\, D))$ as in Section II.5.1 and $D = \cup_i D_i \subset X$. Then L' is a DGLA and we have the following exact sequence

$$0 \longrightarrow L' \longrightarrow A_X^{0,*}(\Theta_X) \xrightarrow{\pi'} A_D^{0,*}(\oplus_i N_{D_i|X}) \longrightarrow 0.$$

Denoting for convenience $D^\circ = \overset{\circ}{\bigcup} D_i$, we define $M = KS_{D^\circ \times X}$, $N = KS_{D^\circ} \times KS_X$ and the morphism

$$g = (p^*, q^*) : N = KS_{D^\circ} \longrightarrow M = KS_{D^\circ \times X},$$

where p and q are the projections of $D^\circ \times X$ onto D° and X, respectively.

Finally, let L be the DGLA defined, as in Section IV.2, by the following exact sequence:

$$0 \longrightarrow L \xrightarrow{h} M \xrightarrow{\pi} A^{0,*}_{D^\circ}(f^*\Theta_X) \longrightarrow 0.$$

Corollary V.4.4. *L' controls the functor* $\mathrm{Def}_{X;D_1,\ldots,D_m}$*. In particular, the tangent space of* $\mathrm{Def}_{X;D_1,\ldots,D_m}$ *is $H^1(X, \Theta_X(-log\, D))$ and the obstruction space is naturally contained in $H^2(X, \Theta_X(-log\, D))$.*

Proof. With the notation above, applying Theorem IV.2.5, there exists an isomorphism of functors $\mathrm{Def}_{X;D_1,\ldots,D_m} \cong \mathrm{Def}_{(h,g)}$. Then proceeding as in the case of inclusion it suffices to prove the two steps below.

Step 1. The morphism $g - h : L \times N \longrightarrow M$ such that $(g-h)(l,n) = g(n) - h(l)$ is surjective (analogous of Lemma V.4.2).
Step 2 . $L' \cong L \times_M N$ (analogous of Lemma V.4.3).

Actually, *Step* 1 and Proposition III.2.34 imply that $\mathrm{Def}_{X;D_1,\ldots,D_m} \cong \mathrm{Def}_{(h,g)} \cong \mathrm{Def}_{L\times_M N}$. Finally, *Step* 2 implies $\mathrm{Def}_{X;D_1,\ldots,D_m} \cong \mathrm{Def}_{L'}$.

Proof of Step 1. We want to prove that for each $\phi \in M = A^{0,*}_{D^\circ \times X}(\Theta_{D^\circ \times X})$ there exist $n_1 \in KS_{D^\circ} = A^{0,*}_{D^\circ}(\Theta_{D^\circ})$ and $n_2 \in A^{0,*}_X(\Theta_X)$ such that $g(n_1, n_2) - \phi = p^*n_1 + q^*n_2 - \phi \in L = \ker \pi$, or equivalently $\pi(\phi) = f_*n_1 - f^*n_2 \in A^{0,*}_{D^\circ}(f^*\Theta_X)$. Then we have to prove that $(f^*, -f_*) : A^{0,*}_X(\Theta_X) \times A^{0,*}_{D^\circ}(\Theta_{D^\circ}) \longrightarrow A^{0,*}_{D^\circ}(f^*\Theta_X)$ is surjective and this follows from the hypothesis on D_i.

Proof of Step 2. By definition, $L \times_M N = \{(l, n_1, n_2) \in L \times KS_{D^\circ} \times KS_X \mid h(l) = p^*n_1 + q^*n_2\}$ and so $0 = \pi h(l) = f_*n_1 - f^*n_2$.

Define $\gamma : L \times_M N \longrightarrow KS_X$ as the projection onto KS_X.

Then γ is an injective morphism of DGLAs and its image is $L' = A^{0,*}_X(\Theta_X(-log\, D))$.

Actually, suppose that $\gamma(l, n_1, n_2) = n_2 = 0$; then $f_*n_1 = 0$ and so $n_1 = n_2 = l = 0$.

As to the image, consider the following exact sequences

$$\begin{tikzcd}
0 \arrow[r] & L' \arrow[r] \arrow[d] & A^{0,*}_X(\Theta_X) \arrow[r, "\pi'"] \arrow[d, "f^*"] & A_D(\oplus_i N_{D_i|X}) \arrow[r] \arrow[d, "\cong"'] & 0 \\
0 \arrow[r] & A^{0,*}_{D^\circ}(\Theta_{D^\circ}) \arrow[r, "f_*"] & A^{0,*}_{D^\circ}(f^*\Theta_X) \arrow[r, "\beta"] & A_D(\oplus_i N_{D_i|X}) \arrow[r] & 0.
\end{tikzcd}$$

Let $\gamma(l, n_1, n_2) = n_2 \in KS_X$, then $\beta(f^* n_2) = \beta(f_* n_1) = 0$ and so $\pi'(n_2) = 0$. This implies that $\gamma(l, n_1, n_2) \in L'$.

□

V.4.2. Semiregularity map for the inclusion. Let $i : X \hookrightarrow Y$ be the inclusion of a submanifold X in Y and L' the DGLA (defined in Section II.5.1):

$$0 \longrightarrow L' \longrightarrow A_Y^{0,*}(\Theta_Y) \xrightarrow{\pi'} A_X^{0,*}(N_{X|Y}) \longrightarrow 0.$$

In Corollary V.4.1, we proved that L' controls the infinitesimal deformations of the inclusion i. In particular, this implies that the obstructions are naturally contained in $H^2(L')$.

Moreover, as in Section V.1.1, we can consider the following morphism of DGLAs

$$\boldsymbol{i} : L' \longrightarrow K' = \left\{ f \in \operatorname{Htp}\left(\ker(\partial), \frac{A_Y}{\partial A_Y}\right) \mid f(I_X \cap \ker(\partial)) \subset \frac{I_X}{I_X \cap \partial A_Y} \right\}.$$

Then we get the following corollary, whose proof is essentially contained in Manetti [**24**, Corollary 9.2].

Corollary V.4.5. *Let $i : X \longrightarrow Y$ be the inclusion of a submanifold X in a compact Kähler manifold Y. Let $p = dimY - dimX$. Then the obstruction space to the infinitesimal deformations of i is contained in the kernel of the map*

$$\sigma : H^2(L') \longrightarrow H^{p-1,p+2}(Y, I_X),$$

where $I_X = \ker i^* \subset A_Y^{*,*}$ is the subcomplex of differential forms vanishing on X.

Proof. It follows from Corollary V.3.4, by recalling that $L' \subset \{a \in A_Y^{0,*}(\Theta_Y) | \ \boldsymbol{i}_a(I_X) \subset I_X\}$.

□

Bibliography

[1] M. Artin, *Deformations of Singularities,* Tata Institute of Foundamental Research, Bombay, (1976).

[2] M.F. Atiyah, I.G. MacDonald, *Introduction to Commutative Algebra,* Addison-Wesley Publishing Company, (1969).

[3] S. Bloch, *Semi-regularity and de Rham cohomology,* Invent. Math **17**, (1972), 51-66.

[4] F. Catanese, *Moduli of algebraic surfaces,* Theory of moduli (Montecatini Terme,1985), Lecture Notes in Mathematics, **1337**, Springer-Verlag, New York/Berlin, (1988), 1-83.

[5] H. Clemens, *Geometry of formal Kuranishi theory,* Adv. Math., **198** (No. 1), (2005), 311-365.

[6] B. Fantechi, M. Manetti, *Obstruction calculus for functors of Artin rings, I,* J. Algebra, **202**, (1998), 541-576.

[7] D. Fiorenza, M. Manetti, *L_∞ structures on mapping cones.* Preprint `arXiv:math.QA/0601312`.

[8] K. Fukaya, *Deformation theory, homological algebra and mirror symmetry,* Geometry and physics of branes (Como, 2001), Ser. High Energy Phys. Cosmol. Gravit., IOP Bristol, (2003), 121-209. Electronic version available at `http://www.math.kyoto-u.ac.jp/%7Efukaya/como.dvi`

[9] R. Godement, *Topologie algébrique et théorie des faisceaux,*Actualités scientifiques et industrielles, **1252**, Hermann, Paris, (1958).

[10] W.M. Goldman, J.J. Millson, *The homotopy invariance of the Kuranishi space,* Ill. J. Math., **34**, (1990), 337-367.

[11] Ph. Griffiths, J. Harris, *Principles of algebraic geometry,* Wiley Classics Library, Wiley, New York, (1978).

[12] A. Grothendieck, *Technique de Descente et théorèmes d'existence en géométrie algébrique. II. Le théorèmes d'existence en théorie formelle des modules,* Séminaire Bourbaki, t. 12, Exp. no. **195**, (1959-1960).

[13] R. Hartshorne, *Algebraic Geometry,* Graduate texts in mathematics, **52**, Springer-Verlag, New York/Berlin, (1977).

[14] E. Horikawa, *On deformations of holomorphic maps I, II,* J. Math. Soc. Japan, **25** (No.3), (1973), 372-396; **26** (No.4), (1974), 647-667.

[15] E. Horikawa, *On deformations of holomorphic maps III,* Math. Annalen, **222**, (1976), 275-282.

[16] K. Kodaira, *Complex Manifolds and Deformation of Complex Structures,* Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, **283**, Springer-Verlag, New York/Berlin, (1986).

[17] K. Kodaira, D.C. Spencer, *On Deformations of Complex Analytic Structures, II* Ann. of Math., **67** (2), (1958), 403-466.

[18] M. Kontsevich, *Deformation quantization of Poisson manifolds, I.* Letters in Mathematical Physics, **66**, (2003), 157-216; `arXiv:q-alg/9709040`.

[19] M. Kuranishi, *Deformations of compact complex manifolds,* Séminaire de Mathematiques Supérieures, No. **39** (Été 1969), Les Presses de l'Université de Montreal, Montreal, (1971).

[20] M. Manetti, *Deformation theory via differential graded Lie algebras, Seminari di Geometria Algebrica 1998-1999,* Scuola Normale Superiore (1999).

[21] M. Manetti, *Extended deformation functors.* Int. Math. Res. Not., **14**, (2002), 719-756; `arXiv:math.AG/9910071`.

[22] M. Manetti, *Cohomological constraint to deformations of compact Kähler manifolds.*, Adv. Math., **186**, (2004), 125-142; `arXiv:math.AG/0105175`.

[23] M. Manetti, *Lectures on deformations of complex manifolds,* Rend. Mat. Appl. (7), **24**, (2004), 1-183; `arXiv:math.AG/0507286`.

[24] M. Manetti, *Lie description of higher obstructions to deforming submanifolds,* Preprint `arXiv:math.AG/0507287` v2 7Oct2005.

[25] M. Manetti, *Deformations of products and new examples of obstructed irregular surfaces,* Preprint `arXiv:math.AG/0601169`.

[26] H Matsumura, *Commutative ring theory,* Cambridge Studies in Advanced Mathematics, **8**, Cambridge Univesity Press, Cambridge, (1986).

[27] M. Namba, *Families of meromorphic functions on compact Riemann surfaces,* Lecture Notes in Mathematics, **767**, Springer-Verlag, New York/Berlin, (1979).

[28] Z. Ran, *Deformations of maps*, Algebraic curves and projective geometry (Trento 1989), Lecture Notes in Mathematics, **1389**, Springer-Verlag, New York/Berlin, (1989), 246-253.

[29] Z. Ran, *Semiregularity, obstructions and deformations of Hodge classes*, Ann. Scuola Norm. Pisa Cl. Sci., **28** (No. 4), (1999), 809-821.

[30] Z. Ran, *Universal variations of Hodge structure an Calabi-Yau-Schottky relations*, Invent. Math., **138**, (1999), 425-449.

[31] E. Sernesi, *Deformations of Algebraic Schemes*, Grundlehren der mathematischen Wissenschaften, **334**, Springer-Verlag, New York/Berlin, 2006. Electronic version available at `http://www.mat.uniroma3.it/users/sernesi/defsch.html`

[32] M. Schlessinger, *Functors of Artin rings,* Trans. Amer. Math. Soc., **130**, (1968), 208-222.

[33] C. Voisin, *Théorie de Hodge et géométrie algébrique complexe I,* Société Mathématique de France, Paris, (2002).